

FIGURE 1

CCAATCGCCCGGTGCGGTGGTGCAGGGTCTCGGGCTAGTCATGGCGTCCCCGTCTCGGAGAC
TGCAGACTAAACCAGTCATTACTTGTTTCAAGAGCGTTCTGCTAATCTACACTTTTATTTTC
TGGATCACTGGCGTTATCCTTCTTGCAGTTGGCATTGTTGGGGCAAGGTGAGCCTGGAGAATTA
CTTTTCTCTTTTAAATGAGAAGGCCACCAATGTCCCCTTCGTGCTCATTGCTACTGGTACCG
TCATTATTCTTTTGGGCACCTTTGGTTGTTTTGCTACCTGCCGAGCTTCTGCATGGATGCTA
AAACTGTATGCAATGTTTCTGACTCTCGTTTTTTTTGGTCGAACTGGTCGCTGCCATCGTAGG
ATTTGTTTTTCAGACATGAGATTAAGAACAGCTTTAAGAATAATTATGAGAAGGCTTTGAAGC
AGTATAACTCTACAGGAGATTATAGAAGCCATGCAGTAGACAAGATCCAAAATACGTTGCAT
TGTTGTGGTGTCAACGATTATAGAGATTGGACAGATACTAATTATTACTCAGAAAAAGGATT
TCCTAAGAGTTGCTGTAAACTTGAAGATTGTACTCCACAGAGAGATGCAGACAAAGTAAACA
ATGAAGGTTGTTTTATAAAGGTGATGACCATTATAGAGTCAGAAATGGGAGTCGTTGCAGGA
ATTTCTTTTGGAGTTGCTTGCTTCCAACGATTGGAATCTTTCTCGCCTACTGCCWCTCTCG
TGCCATAACAAATAACCAGTATGAGATAGTGTAACCCAATGTATCTGTGGGCCTATTCCTCT
CTACCTTTAAGGACATTTAGGGTCCCCCTGTGAATTAGAAAGTTGCTTGGCTGGAGAACTG
ACAACACTACTTACTGATAGACCAAAAACTACACCAGTAGGTTGATTCAATCAAGATGTAT
GTAGACCTAAACTACACCAATAGGCTGATTCAATCAAGATCCGTGCTCGCAGTGGGCTGAT
TCAATCAAGATGTATGTTTGCTATGTTCTAAGTCCACCTTCTATCCCATTTCATGTTAGATCG
TTGAAACCCTGTATCCCTCTGAAACACTGGAAGAGCTAGTAAATTGTAAATGAAGT

FIGURE 2

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA19902
><subunit 1 of 1, 245 aa, 1 stop, 1 unknown
><MW: -1, pI: 8.36, NX(S/T): 1
MASPSRRLQTKPVITCFKSVLLIYTFIFWITGVILLAVGIWGKVSLENYFSLLEKATNVPF
VLIATGTVIILLGTFGCFATCRASAWMLKLYAMFLTFLVELVAAIVGFVFRHEIKNSFKN
NYEKALKQYNSTGDYRSHAVDKIQNTLHCCGVTDYRDWTDTNYYSEKGFPSCKLEDCTPQ
RDADKVNNEGCFIKVMTIIESEMGVVAGISFGVACFQLIGIFLAYCXSRITNNQYEIV
```

Important features of the protein:**Signal peptide:**

amino acids 1-42

Transmembrane domains:

amino acids 19-42, 61-83, 92-114, 209-230,

N-glycosylation site.

amino acids 134-138

Tyrosine kinase phosphorylation site.

amino acids 160-168, 160-169

N-myristoylation site.

amino acids 75-81, 78-84, 210-216, 214-220, 226-232

Prokaryotic membrane lipoprotein lipid attachment site.

amino acids 69-80, 211-222

FIGURE 3

CCCACGCGTCCGGCGCCGTGGCCTCGCGTCCATCTTTGCCGTTCTCTCGGACCTGTCACAAA
 GGAGTCGCGCCGCCGCCGCCGCCCTCCCTCCGGTGGGCCCGGGAGGTAGAGAAAGTCAGT
 GCCACAGCCCGACCGCGCTGCTCTGAGCCCTGGGCACGCGGAACGGGAGGGAGTCTGAGGGT
 TGGGGACGTCTGTGAGGGAGGGGAACAGCCGCTCGAGCCTGGGGCGGGCGGACCGGACTGGG
 GCCGGGGTAGGCTCTGGAAAGGGCCCCGGGAGAGAGGTGGCGTTGGTCAGAACCTGAGAAACA
 GCCGAGAGGTTTTCCACCGAGGCCCGCGCTTGAGGGATCTGAAGAGGTTCTTAGAAGAGGGT
 GTTCCCTCTTTCGGGGGTCTCACCAGAAGAGGTTCTTGGGGGTGCGCCTTCTGAGGAGGCT
 GCGGCTAACAGGGCCCAGAACTGCCATTGGATGTCCAGAATCCCCTGTAGTTGATAATGTTG
 GGAATAAGCTCTGCAACTTTCTTTGGCATTGAGTTGTTAAAAACAAATAGGATGCAAATTCC
 TCAACTCCAGGTTATGAAAACAGTACTTGGAAAACGAAAACCTAAATGATCGTCTTTG
 GTTGGGCCGTGTTCTTAGCGAGCAGAAGCCTTGGCCAGGGTCTGTTGTTGACTCTCGAAGAG
 CACATAGCCCACTTCCTAGGGACTGGAGGTGCCGCTACTACCATGGGTAATTCCTGTATCTG
 CCGAGATGACAGTGGAACAGATGACAGTGTTGACACCCAACAGCAACAGGCCGAGAACAGTG
 CAGTACCCACTGCTGACACAAGGAGCCAACCACGGGACCCTGTTTCGGCCACCAAGGAGGGGC
 CGAGGACCTCATGAGCCAAGGAGAAAGAAACAAAATGTGGATGGGCTAGTGTTGGACACACT
 GGCAGTAATACGGACTCTTGTAGATAAGTAAGTATCTGACTCACGGTCACCTCCAGTGGAAT
 GAAAAGTGTTCTGCCCGGAACCATGACTTTAGGACTCCTTCAGTTCCTTTAGGACATACTCG
 CCAAGCCTTGTGCTCACAGGGCAAAGGAGAATATTTAATGCTCCGCTGATGGCAGAGTAAA
 TGATAAGATTTGATGTTTTTGTCTGCTGTCATCTACTTTGTCTGGAAATGTCTAAATGTTTC
 TGTAGCAGAAAACACGATAAAGCTATGATCTTTATTAGAG

FIGURE 4

MIVFGWAVFLASRSLGQGLLLTLEEHIAHFLGTGGAATTMGNSCICRDDSGTDDSVDTQQQQ
AENSAVPTADTRSQPRDPVRPPRRGRGPHEPRRKKQNVDGLVLDTLAVIRTLVDKO

Signal peptide:

amino acids 1-16

Casein kinase II phosphorylation site.

amino acids 22-26, 50-54, 113-117

N-myristoylation site.

amino acids 18-24, 32-38, 34-40, 35-41, 51-57

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100

FIGURE 5

GGCACGAGGCGCTGTCCACCCGGGGGCGTGGGAGTGAGGTACCAGATTAGCCCATTTGGCC
CCGACGCCTCTGTTCTCGGAATCCGGGTGCTGCGGATTGAGGTCCCGGTTCCCTAACGGACTG
CAAGATGGAGGAAGGCGGGAACCTAGGAGGCCTGATTAAGATGGTCCATCTACTGGTCTTGT
CAGGTGCCTGGGGCATGCAAATGTGGGTGACCTTCGTCTCAGGCTTCCTGCTTTTCCGAAGC
CTTCCCCGACATACCTTCGGACTAGTGCAGAGCAAACCTCTTCCCCTTCTACTTCCACATCTC
CATGGGCTGTGCCTTCATCAACCTCTGCATCTTGGCTTCACAGCATGCTTGGGCTCAGCTCA
CATTCTGGGAGGCCAGCCAGCTTTACCTGCTGTTCCCTGAGCCTTACGCTGGCCACTGTCAAC
GCCCCTGGCTGGAACCCCGCACCACAGCTGCCATGTGGGCCCTGCAAACCGTGGAGAAGGA
GCGAGGCCTGGGTGGGGAGGTACCAGGCAGCCACCAGGGTCCCGATCCCTACCGCCAGCTGC
GAGAGAAGGACCCCAAGTACAGTGCTCTCCGCCAGAATTTCTTCCGCTACCATGGGCTGTCC
TCTCTTTGCAATCTGGGCTGCGTCCTGAGCAATGGGCTCTGTCTCGCTGGCCTTGCCCTGGA
AATAAGGAGCCTCTAGCATGGGCCCTGCATGCTAATAAATGCTTCTTCAGAAATGAAAAAA
AAAAAAAAAAAA

bioRxiv preprint doi: <https://doi.org/10.1101/000000>; this version posted January 1, 2014. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.

FIGURE 6

```
</usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA56107
```

<subunit 1 of 1, 231 aa, 1 stop

 $\langle NX(S/T) : 0$

MEEGGNLGGLIKMVHLLVLSGAWGMQMWVTFVSGFLLFRSLPRHTFGLVQSKLFPPFYFHISM
GCAFINLCILASQHAWAQLTFWEASQLYLLFLSLTLATVNARWLEPRTTAAMWALQTVEKER
GLGGEVPGSHQGPDPYRQLREKDPKYSALRQNFRRYHGLSSLCNLGCVLSNGLCLAGLALEIRSL

Signal peptide:

amino acids 1-24

Transmembrane domain:

amino acids 86-103, 60-75

Casein kinase II phosphorylation site.

amino acids 82-86

Tyrosine kinase phosphorylation site.

amino acids 144-151

N-myristoylation site.

amino acids 4-10, 5-11, 47-53, 170-176, 176-182

Prokaryotic membrane lipoprotein lipid attachment site.

amino acids 54-65

G-protein coupled receptors proteins.

amino acids 44-85

FIGURE 7

AATTCAGATTTTAAAGCCATTCTGCAGTGGAATTCATGAACTAGCAAGAGGACACCATCTT
 CTTGTATTATACAAGAAAGGAGTGTACCTATCACACACAGGGGGAAAA**ATG**CTCTTTTGGGT
 GCTAGGCCTCCTAATCCTCTGTGGTTTTCTGTGGACTCGTAAAGGAAAATAAGATTGAAG
 ACATCACTGATAAGTACATTTTTATCACTGGATGTGACTCGGGCTTTGGAACTTGGCAGCC
 AGAACTTTTGATAAAAAGGGATTTTCATGTAATCGCTGCCTGTCTGACTGAATCAGGATCAAC
 AGCTTTAAAGGCAGAAACCTCAGAGAGACTTCGTACTGTGCTTCTGGATGTGACCGACCCAG
 AGAATGTCAAGAGGACTGCCCAGTGGGTGAAGAACCAAGTTGGGGAGAAAGGTCTCTGGGGT
 CTGATCAATAATGCTGGTGTTCCTGGCTGCCACTGACTGGCTGACACTAGAGGA
 CTACAGAGAACCTATTGAAGTGAACCTGTTTGGACTCATCAGTGTGACACTAAATATGCTTC
 CTTTGGTCAAGAAAGCTCAAGGGAGAGTTATTAATGTCTCCAGTGTGGAGGTCGCCCTTGCA
 ATCGTTGGAGGGGGCTATACTCCATCCAAATATGCAGTGGAAGGTTTCAATGACAGCTTAAG
 ACGGGACATGAAAGCTTTTGGTGTGCACGTCTCATGCATTGAACCAGGATTGTTCAAAACAA
 ACTTGGCAGATCCAGTAAAGGTAATTGAAAAAAACTCGCCATTTGGGAGCAGCTGTCTCCA
 GACATCAAACAACAATATGGAGAAGGTTACATTGAAAAAAGTCTAGACAACTGAAAGGCAA
 TAAATCCTATGTGAACATGGACCTCTCTCCGGTGGTAGAGTGCATGGACCACGCTCTAACAA
 GTCTCTTCCCTAAGACTCATTATGCCGCTGGAAAAGATGCCAAAATTTTCTGGATACCTCTG
 TCTCACATGCCAGCAGCTTTGCAAGACTTTTTATTGTTGAAACAGAAAGCAGAGCTGGCTAA
 TCCCAAGGCAGTGT**TGA**CTCAGCTAACCAAAATGTCTCCTCCAGGCTATGAAATTGGCCGAT
 TTCAAGAACACATCTCCTTTTCAACCCATTCTTATCTGCTCCAACCTGGACTCATTTAGA
 TCGTGCTTATTTGGATTGCAAAAGGGAGTCCCACCATCGCTGGTGGTATCCCAGGGTCCCTG
 CTCAAGTTTTCTTTGAAAAGGAGGGCTGGAATGGTACATCACATAGGCAAGTCCTGCCCTGT
 ATTTAGGCTTTGCCTGCTTGGTGTGATGTAAGGGAAATTGAAAGACTTGCCCATTCAAAATG
 ATCTTTACCGTGGCCTGCCCCATGCTTATGGTCCCCAGCATTTACAGTAACTTGTGAATGTT
 AAGTATCATCTCTTATCTAAATATTAAAAGATAAGTCAACCCAAAAAAAAAAAAAAAAAAAA
 AAAAAAAAAAAAAAAAAAAAA

1000
 900
 800
 700
 600
 500
 400
 300
 200
 100
 0

FIGURE 8

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA56406
><subunit 1 of 1, 319 aa, 1 stop
><MW: 35227, pI: 8.97, NX(S/T): 3
MLFWVLGLLILCGFLWTRKGKLIKIEDITDKYIFITGCDSGFGNLAARTFDKKGFHVIAACLT
ESGSTALKAETSERLRTVLLDVTDPENVKRTAQWVKVQVGEKGLWGLINNAGVPGVLAPTDW
LTLEDYREPIEVNLFGLISVTLNMLPLVKKAQGRVINVSSVGGRLAIVGGGYTPSKYAVEGF
NDSLRRDMKAFGVHVSCIEPGLFKTNLADPVKVIEKKLAIWEQLSPDIKQQYGEGYIEKSLD
KLKGNKSYVNMDLSPVVECMDHALTSLFPKTHYAAGKDAKIFWIPLSHMPAALQDFLLKQK
AELANPKAV
```

Important features of the protein:**Signal peptide:**

amino acids 1-17

Transmembrane domain:

amino acids 136-152

N-glycosylation sites.

amino acids 161-163, 187-190 and 253-256

Glycosaminoglycan attachment site.

amino acids 39-42

N-myristoylation sites.

amino acids 36-41, 42-47, 108-113, 166-171, 198-203 and 207-212

FIGURE 9

GCGGGCTGTTGACGGCGCTGCG**ATG**GCTGCCTGCGAGGGCAGGAGAAGCGGAGCTCTCGGTT
 CCTCTCAGTCGGACTTCCTGACGCCGCCAGTGGGCGGGGCCCCTTGGGCCGTGCGCACCACT
 GTAGTCATGTACCCACCGCCGCCGCCGCCCTCATCGGGACTTCATCTCGGTGACGCTGAG
 CTTTGGCGAGAGCTATGACAACAGCAAGAGTTGGCGGGCGGCGCTCGTGCTGGAGGAAATGGA
 AGCAACTGTCGAGATTGCAGCGGAATATGATTCTCTTCCTCCTTGCCCTTTCTGCTTTTCTGT
 GGACTCCTCTTCTACATCAACTTGGCTGACCATTGGAAAGCTCTGGCTTTTCAGGCTAGAGGA
 AGAGCAGAAGATGAGGCCAGAAATTGCTGGGTAAAACCAGCAAATCCACCCGTCTTACCAG
 CTCCTCAGAAGGCGGACACCGACCCTGAGAACTTACCTGAGATTTTCGTACAGAAGACACAA
 AGACACATCCAGCGGGGACCACCTCACCTGCAGATTAGACCCCCAAGCCAAGACCTGAAGGA
 TGGGACCCAGGAGGAGGCCACAAAAGGCAAGAAGCCCTGTGGATCCCCGCCCCGGAAGGAG
 ATCCGCAGAGGACAGTCATCAGCTGGAGGGGAGCGGTGATCGAGCCTGAGCAGGGCACCGAG
 CTCCCTTCAAGAAGAGCAGAAAGTGCCACCAAGCCTCCCCTGCCACCGGCCAGGACACAGGG
 CACACCAGTGCACTCTGAAGTATCGCCAGAAGGGCGTGATTGACGTCTTCCTGCATGCATGGA
 AAGGATACCGCAAGTTTGCATGGGGCCATGACGAGCTGAAGCCTGTGTCCAGGTCTTTCAGT
 GAGTGGTTTGGCCTCGGTCTCACACTGATCGACGCGCTGGACACCATGTGGATCTTGGGTCT
 GAGGAAAGAATTTGAGGAAGCCAGGAAGTGGGTGTGGAAGAAGTTACACTTTGAAAAGGACG
 TGGACGTCAACCTGTTTGGAGAGCACGATCCGCATCCTGGGGGGGCTCCTGAGTGCCTACCAC
 CTGTCTGGGGACAGCCTCTTCCTGAGGAAAGCTGAGGATTTTGGAAATCGGCTAATGCCTGC
 CTTCAGAACACCATCCAAGATTCCTTACTCGGATGTGAACATCGGTACTGGAGTTGCCACC
 CGCCACGGTGGACCTCCGACAGCACTGTGGCCGAGGTGACCAGCATTACAGCTGGAGTTCGGG
 GAGCTCTCCCGTCTCACAGGGGATAAGAAGTTTCAGGAGGCAGTGGAGAAGGTGACACAGCA
 CATCCACGGCCTGTCTGGGAAGAAGGATGGGCTGGTGCCCATGTTCAATACCCACAGTG
 GCCTCTTCACCCACCTGGGCGTATTACGCTGGGCGCCAGGGCCGACAGCTACTATGAGTAC
 CTGCTGAAGCAGTGGATCCAGGGCGGGGAAGCAGGAGACACAGCTGCTGGAAGACTACGTGGA
 AGCCATCGAGGGTGTGAGAACGCACCTGCTGCGGCACTCCGAGCCCAGTAAGCTCACCTTTG
 TGGGGGAGCTTGCCACCGGCCGCTTCAGTGCCAAAGATGGACCACCTGGTGTGCTTCTGCCA
 GGGACGCTGGCTCTGGGCGTCTACCACGGCCTGCCCGCCAGCCACATGGAGCTGGCCAGGA
 GCTCATGGAGACTTGTTACCAGATGAACCGGCAGATGGAGACGGGGCTGAGTCCCGAGATCG
 TGCACTTCAACCTTTACCCCCAGCCGGGCCGTCGGGACGTGGAGGTCAAGCCAGCAGACAGG
 CACAACCTGCTGCGGCCAGAGACCGTGGAGAGCCTGTTCTACCTGTACCGCGTCACAGGGGA
 CCGCAAATACCAGGACTGGGGCTGGGAGATTCTGCAGAGCTTCAGCCGATTACACGGGTCC
 CCTCGGGTGGCTATTCTTCCATCAACAATGTCCAGGATCCTCAGAAGCCCAGCCTAGGGAC
 AAGATGGAGAGCTTCTTCCTGGGGGAGACGCTCAAGTATCTGTTCTTGCTCTTCTCCGATGA
 CCCAAACCTGCTCAGCCTGGACGCCTACGTGTTCAACACCGAAGCCCACCCTCTGCCTATCT
 GGACCCCTGCC**TAG**GGTGGATGGCTGCTGGTGTGGGGACTTCGGGTGGGCAGAGGCACCTTG
 CTGGGTCTGTGGCATTTTCCAAGGGCCACGTAGCACCGGCAACCGCCAAGTGGCCCAGGCT
 CTGAACTGGCTCTGGGCTCCTCCTCGTCTCTGCTTTAATCAGGACACCGTGAGGACAAGTGA
 GGCCGTCAGTCTTGGTGTGATGCGGGGTGGGCTGGGCCGCTGGAGCCTCCGCCTGCTTCTC
 CAGAAGACACGAATCATGACTCACGATTGCTGAAGCCTGAGCAGGTCTCTGTGGGCCGACCA
 GAGGGGGGCTTCGAGGTGGTCCCTGGTACTGGGGTGACCGAGTGGACAGCCCAGGGTGCAGC
 TCTGCCCGGGCTCGTGAAGCCTCAGATGTCCCAATCCAAGGGTCTGGAGGGGCTGCCGTGA
 CTCCAGAGGCCTGAGGCTCCAGGGCTGGCTCTGGTGTTTACAAGCTGGACTCAGGGATCCTC
 CTGGCCGCCCCGACAGGGGGCTTGAGGGGCTGGACGGCAAGTCCGTCTAGCTCACGGGCCCCCT
 CCAGTGGAATGGGTCTTTTCGGTGGAGATAAAAGTTGATTGTCTTAACCGCAA

FIGURE 10

></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA56529

><subunit 1 of 1, 699 aa, 1 stop

><MW: 79553, pI: 7.83, NX(S/T): 0

MAACEGRRSGALGSSQSDFLTTPVGGAPWAVATTVVMYPPPPPPPHRDFISVTLSFGESYDN
 SKSWRRRSCWRKWKQLSRLQQRNMILFLLAFLLFCGLLFYINLADHWKALAFRLEEEQKMRPE
 IAGLKPANPPVLPAPQKADTDPENLPEISSQKTQRHIQRGPPHLQIRPPSQDLKDGTQEEAT
 KRQEAPVDPRPEGDPQRTVISWRGAVIEPEQGTELPSRRAEVPTKPPLPPARTQGTPVHLNY
 RQKGVIDVFLHAWKGYRKFAWGHDELKPVSRSFSEWFGLGLTLIDALDTMWILGLRKEFEEA
 RKWVSKKLHFEKDVDVNLFESTITIRILGGLLSAYHLSGDSLFLRKAEDFGNRLMPAFRTPSKI
 PYSDVNIGTGVAHPPRWTSDSTVAEVTISIQLFRELRSRLTGDKKFQEAVEKVTQHIHGLSGK
 KDGLVPMFINTHSGLFTHLGVFTLGARADSYEYLLKQWIQGGKQETQLLEDYVEAIEGVRT
 HLLRHSEPSKLT FVGELAHGRFSAKMDHLVCFLPGTLALGVYHGLPASHMELAQELMETCYQ
 MNRQMETGLSPEIVHFNLYPQPGRRDVEVKPADRHNLRLPETVESLFYLYRVTGDRKYQDWG
 WEILQSFSRFRTRVPSGGYSSINNVQDPQKPEPRDKMESFFLGETLKYLFLLFSDDPNLLSLD
 AYVFNTEAHPLPIWTPA

Important features of the protein:

Transmembrane domain:

amino acids 21-40 and 84-105 (type II)

FIGURE 11

GGCGCCGCGTAGGCCCGGGAGGCCGGGCCGGGCTGCGAGCGCCTGCCCCATGCGCCGC
CGCCTCTCCGCACG**ATG**TTCCCCTCGCGGAGGAAAGCGGCGCAGCTGCCCTGGGAGGACGGC
AGGTCCGGGTTGCTCTCCGGCGGCCTCCCTCGGAAGTGTTCCGTCTTCCACCTGTTTCGTGGC
CTGCCTCTCGCTGGGCTTCTTCTCCCTACTCTGGCTGCAGCTCAGCTGCTCTGGGGACGTGG
CCCGGGCAGTCAGGGGACAAGGGCAGGAGACCTCGGGCCCTCCCCGTGCCTGCCCCCAGAG
CCGCCCCCTGAGCACTGGGAAGAAGACGCATCCTGGGGCCCCCACC GCCTGGCAGTGCTGGT
GCCCTTCCGCGAACGCTTCGAGGAGCTCCTGGTCTTCGTGCCCCACATGCGCCGCTTCCTGA
GCAGGAAGAAGATCCGGCACCACATCTACGTGCTCAACCAGGTGGACCACTTCAGGTTCAAC
CGGGCAGCGCTCATCAACGTGGGCTTCCCTGGAGAGCAGCAACAGCACGGACTACATTGCCAT
GCACGACGTTGACCTGCTCCCTCTCAACGAGGAGCTGGACTATGGCTTTCCTGAGGCTGGGC
CCTTCCACGTGGCCTCCCCGGAGCTCCACCCTCTCTACCACTACAAGACCTATGTCGGCGGC
ATCCTGCTGCTCTCCAAGCAGCACTACCGGCTGTGCAATGGGATGTCCAACCGCTTCTGGGG
CTGGGGCCGCGAGGACGACGAGTTCTACCGGCGCATTAAGGGAGCTGGGCTCCAGCTTTTCC
GCCCCCTCGGGAATCACAACCTGGGTACAAGACATTTCGCCACCTGCATGACCCAGCCTGGCGG
AAGAGGGACCAGAAGCGCATCGCAGCTCAAAAACAGGAGCAGTTCAAGGTGGACAGGGAGGG
AGGCCTGAACACTGTGAAGTACCATGTGGCTTCCCGCACTGCCCTGTCTGTGGGCGGGGCCC
CCTGCACTGTCTCAACATCATGTTGGACTGTGACAAGACCGCCACACCCTGGTGCACATTC
AGCT**TGA**GCTGGATGGACAGTGAGGAAGCCTGTACCTACAGGCCATATTGCTCAGGCTCAGGA
CAAGGCCTCAGGTCGTGGGCCCAGCTCTGACAGGATGTGGAGTGGCCAGGACCAAGACAGCA
AGCTACGCAATTGCAGCCACCCGGCCGCAAGGCAGGCTTGGGCTGGGCCAGGACACGTGGG
GTGCCTGGGACGCTGCTTGCCATGCACAGTGATCAGAGAGAGGCTGGGGTGTGTCTGTCCG
GGACCCCCCTGCCTTCCTGCTCACCTACTCTGACCTCCTTCACGTGCCCAGGCCTGTGGG
TAGTGGGGAGGGCTGAACAGGACAACCTCTCATCACCTACTCTGACCTCCTTCACGTGCCC
AGGCCTGTGGGTAGTGGGGAGGGCTGAACAGGACAACCTCTCATCACCCCCAAAAAAAAAAAA
AAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA

FIGURE 12

></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA56531

><subunit 1 of 1, 327 aa, 1 stop

><MW: 37406, pI: 9.30, NX(S/T): 1

MFPSRRKAAQLPWEDGRSGLLSGGLPRKCSVFHLEFVACLSLGFFSLLWLQLSCSGDVARAVR
GQGQETSGPPRACPPPEPPPEHWEEDASWGPHRLAVLVPFRERFEELLVFVPHMRRFLSRKKI
RHHIYVLNQVDHFRFNRAALINVGFLSSNSTDYIAMHDVDLLPLNEELDYGFPEAGPFHVA
SPELHPLYHYKTYVGGILLLSKQHYRLCNGMSNRFWGWGREDDFYRRIKGAGLQLFRPSGI
TTGYKTRHLDPAWRKRDQKRIAAQKQEQFKVDREGGLNTVKYHVASRTALSVGGAPCTVL
NIMLDCDKTATPWCTFS

Signal peptide:

amino acids 1-42

Transmembrane domain:

amino acids 29-49 (type II)

N-glycosylation site.

amino acids 154-158

cAMP- and cGMP-dependent protein kinase phosphorylation site.

amino acids 27-31

Tyrosine kinase phosphorylation site.

amino acids 226-233

N-myristoylation site.

amino acids 19-25, 65-71, 247-253, 285-291, 303-309, 304-310

FIGURE 13

CAATGTTTGCCTATCCACCTCCCCCAAGCCCCTTTACCT**ATG**CTGCTGCTAACGCTGCTGCT
GCTGCTGCTGCTGCTTAAAGGCTCATGCTTGGAGTGGGGACTGGTCGGTGCCCAGAAAGTCT
CTTCTGCCACTGACGCCCCCATCAGGGATTGGGCCTTCTTTCCCCCTTCCTTTCTGTGTCTC
CTGCCTCATCGGCCTGCCATGACCTGCAGCCAAGCCCAGCCCCGTGGGGAAGGGGAGAAAGT
GGGGGATGGC**TAA**GAAAGCTGGGAGATAGGGAACAGAAGAGGGTAGTGGGTGGGCTAGGGGG
GCTGCCTTATTTAAAGTGTTGTTTATGATTCTTATACTAATTTATACAAAGATATTAAGGC
CCTGTTTCATTAAGAAATTGTTCCCTTCCCCTGTGTTCAATGTTTGTAAAGATTGTTCTGTGT
AAATATGTCTTTATAATAAACAGTTAAAAGCTGAAAAAAAAAAAAAAAAAAAAAAAAAAAAA

1000 900 800 700 600 500 400 300 200 100 0

FIGURE 14

</usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA56862

<subunit 1 of 1, 73 aa, 1 stop

<MW: 7879, pI: 7.21, NX(S/T): 0

MLLLTLLLLLLLLLLKGSCLEWGLVGAQKVSSATDAPIRDWAFFPPSFLCLLPHRPAMTCSQAQ
PRGEGEKVGDG

Important features:**Signal peptide:**

amino acids 1-15

Growth factor and cytokines receptors family:

amino acids 3-18

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100

FIGURE 15

GGGACCCATGCGGCCGTGACCCCCGGCTCCCTAGAGGCCAGCGCAGCCGCAGCGGACAAAG
 GAGCATGTCCGCGCCGGGAAGGCCCGTCTCCGGCCGCCATAAGGCTCCGGTCGCCGCTGG
 GCCCGCGCCGCGCTCCTGCCCGCCGGGCTCCGGGGCGGCCCGCTAGGCCAGTGCGCCGCCG
 CTCGCCCCGCAGGCCCGGCCCGCAGC**ATG**GAGCCACCCGGACGCCGGCGGGGCCGCGCGCA
 GCCGCCGCTGTTGCTGCCGCTCTCGCTGTTAGCGCTGCTCGCGCTGCTGGGAGGCGGGCGCG
 GCGGCGGCGCCGCGGCGCTGCCCGCCGGCTGCAAGCACGATGGGCGGCCCGAGGGGCTGGC
 AGGGCGGCGGGCGCCGCCGAGGGCAAGGTGGTGTGCAGCAGCCTGGAACCTCGCGCAGGTCCT
 GCCCCCAGATACTCTGCCCAACCGCACGGTCACCCTGATTCTGAGTAACAATAAGATATCCG
 AGCTGAAGAATGGCTCATTTTCTGGGTAAAGTCTCCTTGAAAGATTGGACCTCCGAAACAAT
 CTTATTAGTAGTATAGATCCAGGTGCCTTCTGGGGACTGTCATCTCTAAAAAGATTGGATCT
 GACAAACAATCGAATAGGATGTCTGAATGCAGACATATTTGAGGACTCACCAATCTGGTTC
 GGCTAAACCTTTTCGGGAATTTGTTTTCTTCATTATCTCAAGGAACCTTTTGATTATCTTGCG
 TCATTACGGTCTTTGGAATTCAGACTGAGTATCTTTTGTGTGACTGTAACATACTGTGGAT
 GCATCGCTGGGTAAAGGAGAAGAACATCACGGTACGGGATACCAGGTGTGTTTATCCTAAGT
 CACTGCAGGCCCAACCAGTCACAGGCGTGAAGCAGGAGCTGTTGACATGCGACCCTCCGCTT
 GAATTGCCGTCTTTCTACATGACTCCATCTCATCGCCAAGTTGTGTTTGAAGGAGACAGCCT
 TCCTTTCCAGTGCATGGCTTCATATATTGATCAGGACATGCAAGTGTTGTGGTATCAGGATG
 GGAGAATAGTTGAAACCGATGAATCGCAAGGTATTTTGTGTTGAAAAGAACATGATTCACAAC
 TGCTCCTTGATTGCAAGTGCCCTAACCATTTCTAATATTCAGGCTGGATCTACTGGAAATTG
 GGGCTGTCATGTCCAGACCAAACGTGGGAATAATACGAGGACTGTGGATATTGTGGTATTAG
 AGAGTTCTGCACAGTACTGTCTCCAGAGAGGGTGGTAAACAACAAAGGTGACTTCAGATGG
 CCCAGAACATTGGCAGGCATTACTGCATATCTGCAGTGTACGCGGAACACCCATGGCAGTGG
 GATATATCCCGGAAACCCACAGGATGAGAGAAAAGCTTGGCGCAGATGTGATAGAGGTGGCT
 TTTGGGCAGATGATGATTATTCTCGCTGTCAGTATGCAAATGATGTCACTAGAGTTCTTTAT
 ATGTTTAATCAGATGCCCTCAATCTTACCAATGCCGTGGCAACAGCTCGACAGTTACTGGC
 TTACACTGTGGAAGCAGCCAACCTTTCTGACAAAATGGATGTTATATTTGTGGCAGAAATGA
 TTGAAAAATTTGGAAGATTTACCAAGGAGGAAAAATCAAAAGAGCTAGGTGACGTGATGGTT
 GACATTGCAAGTAACATCATGTTGGCTGATGAACGTGTCTGTGGCTGGCGCAGAGGGAAGC
 TAAAGCCTGCAGTAGGATTGTGCAGTGTCTTCAGCGCATTGCTACCTACCGGCTAGCCGGTG
 GAGCTCACGTTTATTCAACATATTCACCCAATATTGCTCTGGAAGCTTATGTCATCAAGTCT
 ACTGGCTTCACGGGGATGACCTGTACCGTGTTCAGAAAGTGGCAGCCTCTGATCGTACAGG
 ACTTTTCGGATTATGGGAGGCGGGATCCAGAGGGGAAACCTGGATAAGCAGCTGAGCTTTAAGT
 GCAATGTTTCAAATACATTTTCGAGTCTGGCACTAAAGGTATGTTACATTCTGCAATCATTT
 AAGACTATTTACAGT**TAA**ATTAGAATGCTCCAAATGTTCTGCTTCGCAAAATAACCTTATTA
 AAAGATTTTTTTTTTGCAAGGAAGATAGGTATTATTGCTTTTGCTACTGTTTTAAAGAAAATA
 ACCAGGAAGAACTGCATTACGACTTTCAAGGGCCCTAGGCATTTTTGCCTTTGATTCCCTTT
 CTTACATAAAAAATATCAGAAATTACATTTTATAACTGCAGTGGTATAAATGCAAATATACT
 ATTGTTACATGTGAAAAAATTTTATTTGACTTAAAAGTTTATTTATTTGTTTTTTTTGCTCCT
 GATTTTAAAGACAATAAGATGTTTTTCATGGGCCCTAAAAGTATCATGAGCCTTTGGCACTGC
 GCCTGCCAAGCCTAGTGGAGAAGTCAACCCTGAGACCAGGTGTTTAATCAAGCAAGCTGTAT
 ATCAAAATTTTTTGGCAGAAAAACAAATATGTCATATATCTTTTTTTTAAAAAAAGTATTTCA
 TTGAAGCAAGCAAAATGAAAGCATTTTTTACTGATTTTTTAAATTTGGTGCTTTAGATATATTT
 GACTACACTGTATTGAAGCAAAATAGAGGAGGCACAACTCCAGCACCCCTAATGGAACCACATT
 TTTTTTCACTTAGCTTTTCTGTGGGCATGTGTAATTGTATTCTCTGCGGTTTTTAAATCTCACAG
 TACTTTATTTCTGTCTTGTCCCTCAATAATATCACAAACAATATTCAGTCATTTTAAATGGC
 TGCATAATAACTGATCCAACAGGTGTTAGGTGTTCTGGTTTGTGAGCACTCAATAAATA
 TTGAATGAATGAACGAAAAAAAAAAAAAAAAA

FIGURE 16

MEPPGRRRRGRAQPPLLLPLSLLALLALLGGGGGGGAAALPAGCKHDGRPRGAGRAAGAAEGK
 VVCSSLELAQVLPPDTLPNRTVTILSNNKISELKNGSFSGLSLLERLDLRNNLISSIDPGA
 FWGLSSLKRLDLTNNRIGCLNADIFRGLTNLVRLNLSGNLFSSLSQGTFDYLASLRSLEFQT
 EYLLCDCNILWMHRWVKEKNITVRDTRCVYPKSLQAQPVTGVKQELLTCDPPELPSFYMTP
 SHRQVVFEGLSLPFQCMASYIDQDMQVLWYQDGRIVETDESQGFVEKNMIHNCSLIASALT
 ISNIQAGSTGNWGCHVQTKRGNNTRTVDIVVLESSAQYCPPERVVNNKGDFRWPRTLAGITA
 YLQCTRNTHGSGIYPGNPQDERKAWRRCDRGGFWADDDYSRCQYANDVTRVLYMFNQMPNLN
 TNAVATARQLLAYTVEAANFSDKMDVIFVAEMIEKFGRFTKEEKSKELGDMVMDIASNIMLA
 DERVLWLAQREAKACSRIVQCLQRIATYRLAGGAHVYSTYSPNIALEAYVIKSTGFTGMTCT
 VFQKVAASDRGTGLSDYGRRDPEGNLDKQLSFKCNVSNTFSSLALKVCYILQSFKTIYS

Signal peptide:

amino acids 1-33

Transmembrane domain:

amino acids 13-40 (type II)

N-glycosylation site.

amino acids 81-85, 98-102, 159-163, 206-210, 301-305, 332-336,
 433-437, 453-457, 592-596

N-myristoylation site.

amino acids 29-35, 30-36, 31-37, 32-38, 33-39, 34-40, 51-57,
 57-63, 99-105, 123-129, 142-148, 162-168, 317-323, 320-326,
 384-390, 403-409, 554-560

FIGURE 17

GCGTGGGGATGTCTAGGAGCTCGAAGGTGGTGTGGGCCTCTCGGTGCTGCTGACGGCGGCC
ACAGTGGCCGGCGTACATGTGAAGCAGCAGTGGGACCAGCAGAGGCTTCGTGACGGAGTTAT
CAGAGACATTGAGAGGCAAATTCGGAAAAAGAAAACATTCGTCTTTTGGGAGAACAGATTA
TTTTGACTGAGCAACTTGAAGCAGAAAGAGAGAAGATGTTATTGGCAAAGGATCTCAAAAA
TCATTGACTTGAATGTGAAATATCTGTTGGACAGACAACACGAGTTTGTGTGTGTGTGTTGAT
GGAGAGTAGCTTAGTAGTATCTTCATCTTTTTTTTTTGGTCACTGTCCTTTTAAACTTGATCA
AATAAAGGACAGTGGGTCATATAAGTTACTGCTTTCAGGGTCCCTTATATCTGAATAAAGGA
GTGTGGGCAGACACTTTTTGGAAGAGTCTGTCTGGGTGATCCTGGTAGAAGCCCCATTAGGG
TCACTGTCCAGTGCTTAGGGTTGTTACTGAGAAGCACTGCCGAGCTTGTGAGAAGGAAGGGA
TGGATAGTAGCATCCACCTGAGTAGTCTGATCAGTCGGCATGATGACGAAGCCACGAGAACA
TCGACCTCAGAAGGACTGGAGGAAGGTGAAGTGAGGGGAGAGACGCTCCTGATCGTCGAATCC

1000 900 800 700 600 500 400 300 200 100 0

FIGURE 18

MSRSSKVVLGLSVLLTAATVAGVHVKQQWDQQRLRDGVIRDIERQIRKKENIRLLGEQIILT
EQLEAEREKMLLAKGSQKS

Signal peptide:

amino acids 1-21

MSRSSKVVLGLSVLLTAATVAGVHVKQQWDQQRLRDGVIRDIERQIRKKENIRLLGEQIILT
EQLEAEREKMLLAKGSQKS

FIGURE 19

CTGTCGCTCTTTGCTTTCAGCCGCACTGCGCTGGCTGCCTGAGGTGCTCTTACAGCCTGTTC
CAAGTGTGGCTTAATCCGTCTCCACCACCAGATCTTTCTCCGTGGATTCCCTCTGCTAAGACC
GCTGCCATGCCAGTGACGGTAACCCGCACCACCATCACAAACCACCACGACGTCACTTCGGG
CCTGGGGTCCCCCATGATCGTGGGGTCCCCTCGGGCCCTGACACAGCCCCTGGGTCTCCTTCGC
CTGCTGCAGCTGGTGTCTACCTGCGTGGCCTTCTCGCTGGTGGCTAGCGTGGGCGCCTGGAC
GGGGTCCATGGGCAACTGGTCCATGTTTACCTGGTGTCTCTGCTTCTCCGTGACCCCTGATCA
TCCTCATCGTGGAGCTGTGCGGGCTCCAGGCCCGCTTCCCCCTGTCTTGGCGCAACTTCCCC
ATCACCTTCGCCTGCTATGCGGCCCTCTTCTGCCTCTCGGCCTCCATCATCTACCCACCAC
CTATGTCCAGTTCCTGTCCACGGCCGTTTCGCGGGACCACGCCATCGCCGCCACCTTCTTCT
CCTGCATCGCGTGTGTGGCTTACGCCACCGAAGTGGCCTGGACCCGGGCCCCGGCCCCGGCGAG
ATCACTGGCTATATGGCCACCGTACCCGGGCTGCTGAAGGTGCTGGAGACCTTCGTTGCCTG
CATCATCTTCGCGTTCATCAGCGACCCCAACCTGTACCAGCACCAGCCGGCCCTGGAGTGGT
GCGTGGCGGTGTACGCCATCTGCTTCATCCTAGCGGCCATCGCCATCCTGCTGAACCTGGGG
GAGTGCACCAACGTGCTACCCATCCCCTTCCCCAGCTTCCTGTGGGGCTGGCCTTGCTGTC
TGTCTCTCTATGCCACCGCCCTTGTTCTCTGGCCCCCTTACCAGTTCGATGAGAAGTATG
GCGGCCAGCCTCGGCGCTCGAGAGATGTAAGCTGCAGCCGCAGCCATGCCTACTACGTGTGT
GCCTGGGACCGCCGACTGGCTGTGGCCATCCTGACGGCCATCAACCTACTGGCGTATGTGGC
TGACCTGGTGCACCTCTGCCACCTGGTTTTTGTCAAGGTCTTAAGACTCTCCCAAGAGGCTCC
CGTTCCCTCTCCAACCTCTTTGTTCTTCTTGCCCGAGTTTTCTTTATGGAGTACTTCTTTCC
TCCGCCTTTCCTCTGTTTTCTCTTCCCTGTCTCCCCCTCCCTCCCACCTTTTTCTTTCCTTCC
CAATTCCCTTGCACTCTAACCAGTTCTTGGATGCATCTTCTTCCCTTCCCTTTCCTCTTGCTGT
TTCCTTCCCTGTGTTGTTTTGTTGCCACATCCTGTTTTACCCCTGAGCTGTTTCTCTTTTT
CTTTTCTTTCTTTTTTTTTTTTTTTTTTTAAGACGGATTCTCACTCTGTGGCCCAGGCTGGAG
TGAGTGGTGCATCTCAGCTCACTGCAACCCCCGCCTCCTGGGTTCAGCGATTCTCCTCC
CCCAGCCTCCCAAGTAGCTGGGAGGACAGGTGTGAGCTGCCGCACCCAGCCTGTTTCTCTTT
TTCCACTCTTCTTTTTTCTCATCTCTTTTCTGGGTGTCCTGTGCGCTTCTTATCTGCCTGT
TTTGCAAGCACCTTCTCCTGTGTCCTTGGGAGCCCTGAGACTTCTTTCTCTCCTTGCCTCCA
CCCACCTCCAAAGGTGCTGAGCTCACATCCACACCCCTTGAGCCGTCCATGCCACAGCCCC
CCAAGGGGGCCCCATTGCCAAAGCATGCCTGCCACCCCTCGCTGTGCCTTAGTCAGTGTGTAC
GTGTGTGTGTGTGTGTGTTGGGGGGTGGGGGGTGGGTAGCTGGGGATTGGGCCCTCTTTCT
CCCAGTGGAGGAAGGTGTGAGTGTACTTCCCCTTTAAATTAAAAAACATATATATATATAT
ATTTGGAGGTCAGTAATTTCCAATGGGCGGGAGGCATTAAGCACCGACCTGGGTCCCTAGG
CCCCGCTGGCACTCAGCCTTGCCAGAGATTGGCTCCAGAATTTTGGCAGGCTTACAGAACAC
CCTACTGCCTAGAGGCCATCTTAAAGGAAGCAGGGGCTGGATGCCTTTCATCCCAACTATTCT
CTGTGGTATGAAAAAG

FIGURE 20

<MW: 35274, pI: 8.57, NX(S/T): 1

MPVTVTRTTITTTTTSSSGLGSPMIVGSPRALTQPLGLLRLLQLVSTCVAFSLVASVGAWTG
SMGNWSMFTWCFCFSVTLLIILIVELCGLQARFPLSWRNFPIITFACYAALFCLSASIIYPTTY
VQFLSHGRSRDHAIAATFFSCIACVAYATEVAWTRARPGEITGYMATVPGLLKVLETFVACI
IFAFISDPNLYQHQPALWCVAVYAICFILAAIAILLNLGECTNVLPPIFPSPFLSGLALLSV
LLYATALVLWPLYQFDEKYGGQPRRSRDVSCSRSHAYYVCAWDRRLAVAILTAINLLAYVAD
LVHSAHLVFEVKV

Transmembrane domains:

amino acids 41-60 (type II), 66-85, 101-120, 137-153, 171-192, 205-226, 235-255 and 294-312

amino acids 66-69

amino acids 18-21

[illegible]

FIGURE 22

MFLATLSFLLPFAHPFGTVSCEYMLGSPLSSLAQVNLSPFSPKPVHMDPNYCHPSTSLHLCS
LAWSFTRLHPPLSPGISQVVKDHVTKPTAMAQGRVAHLIEWKGWSKPSDSPAALESASFSSY
SDLSEGEQEARFAAGVAEQFAIAEAKLRAWSSVDGEDSTDDSYDEDFAGGMDTDMAGQLPLG
PHLQDLFTGHRFSRPVRQGSVEPESDCSQTVSPDTLCSSSLCSLEDGLLGSPARLASQLLGDE
LLLAKLPSPRESAFRSLGPLEAQDSLINSPLTESCLSPAEEEEPAPCKDCQPLCPPLTGSWER
QRQASDLASSGVVSLDEDEAEPEEQ

Signal peptide:

amino acids 1-15

Casein kinase II phosphorylation site.

amino acids 123-127, 128-132, 155-159, 162-166, 166-170, 228-232,
285-289, 324-328

Tyrosine kinase phosphorylation site.

amino acids 44-52

N-myristoylation site.

amino acids 17-23, 26-32, 173-179

Prokaryotic membrane lipoprotein lipid attachment site.

amino acids 11-22

FIGURE 23

GGTTCCTGGGCGCTCTGTTACACAAGCAAGATACAGCCAGCCCCACCTAATTTTGTTCCT
GGCACCCCTCCTGCTCAGTGCACACTTAACCCATCTGTTTTCTCTAATGCACGA
CAGATTCCCTTTCAGACAGGACAACCTGTGATATTTTCAGTTCCTGATTGTAAATACCTCCTAAG
CCTGAAGCTTCTGTTACTAGCCATTGTGAGCTTCAGTTTCTTCATCTGCAAAATGGGCATAA
TACAATCTATTCTTGCCACATCAAGGGATTGTTATTCCCTTTAAAAAAAACCAATACCAAAG
AAGCCTACA**ATG**TTGGCCTTAGCCAAAATTCTGTTGATTTCAACGTTGTTTTATTCACTTCT
ATCGGGGAGCCATGGAAAAGAAAATCAAGACATAAACACAACACAGAACATTGCAGAAGTTT
TTAAAACAATGGAAAATAAACCTATTTCTTTGGAAAGTGAAGCAAACCTTAACTCAGATAAA
GAAAATATAACCACCTCAAATCTCAAGGCGAGTCATTCCCCTCCTTTGAATCTACCCAACAA
CAGCCACGGAATAACAGATTTCTCCAGTAACTCATCAGCAGAGCATTCTTTGGGCAGTCTAA
AACCACATCTACCATTTCCACAAGCCCTCCCTTGATCCATAGCTTTGTTTCTAAAGTGCCT
TGGAATGCACCTATAGCAGATGAAGATCTTTTGCCCATCTCAGCACATCCCAATGCTACACC
TGCTCTGTCTTCAGAAAACCTTCACTTGGTCTTTGGTCAATGACACCGTGAAAACCTCCTGATA
ACAGTTCCATTACAGTTAGCATCCTCTCTTCAGAACCAACTTCTCCATCTGTGACCCCTTG
ATAGTGAACCAAGTGGATGGCTTACCACAAACAGTGATAGCTTCACTGGGTTTACCCCTTA
TCAAGAAAAACAACCTCTACAGCCTACCTTAAAATTACCAATAATTCAAACTCTTTCCAA
ATACGTCAGATCCCCAAAAAGAAAATAGAAATACAGGAATAGTATTCGGGGCCATTTTAGGT
GCTATTCTGGGTGTCTCATTTGCTTACTCTTGTGGGCTACTTGTTGTGTGGAAAAGGAAAAC
GGATTCATTTTCCCATCGGCGACTTTATGACGACAGAAATGAACCAGTTCTGCGATTAGACA
ATGCACCGGAACCTTATGATGTGAGTTTTGGGAATTCTAGCTACTACAATCCAACCTTTGAAT
GATTACAGCCATGCCAGAAAGTGAAGAAAATGCACGTGATGGCATTCCCTATGGATGACATACC
TCCACTTCGTACTTCTGTAT**TAGA**ACTAACAGCAAAAAGGCGTTAAACAGCAAGTGTCTCTA
CATCCTAGCCTTTTGACAAATTCATCTTTCAAAGGTTACACAAAATTACTGTCACGTGGAT
TTTGTCAAGGAGAATCATAAAAGCAGGAGACCAGTAGCAGAAATGTAGACAGGATGTATCAT
CCAAAGGTTTTCTTTCTTACAATTTTGGCCATCCTGAGGCATTTACTAAGTAGCCTTAATT
TGTATTTTAGTAGTATTTTCTTAGTAGAAAATATTTGTGGAATCAGATAAACTAAAAGATT
TCACCATTACAGCCCTGCCTCATAACTAAATAATAAAAATTATTCCACCAAAAAATTCTAAA
ACAATGAAGATGACTCTTTACTGCTCTGCCTGAAGCCCTAGTACCATAATTCAAGATTGCAT
TTTCTTAAATGAAAATTGAAAGGGTGCTTTTTTAAAGAAAATTGACTTAAAGCTAAAAAGAG
GACATAGCCCAGAGTTTCTGTTATTGGGAAATTGAGGCAATAGAAATGACAGACCTGTATTC
TAGTACGTTATAATTTCTAGATCAGCACACATGATCAGCCCACTGAGTTATGAAGCTGA
CAATGACTGCATTCAACGGGGCCATGGCAGGAAAGCTGACCCTACCCAGGAAAGTAATAGCT
TCTTTAAAGTCTTCAAAGGTTTTGGGAATTTTAACTTGTCTTAATATATCTTAGGCTTCAA
TTATTTGGGTGCCTTAAAACTCAATGAGAATCATGGT

FIGURE 24

></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA58732

><subunit 1 of 1, 334 aa, 1 stop

><MW: 36294, pI: 4.98, NX(S/T): 13

MLALAKILLISTLFYSLLSGSHGKENQDINTTQNI AEVFKT MENKPISLESEANLNSDKENI
TTSNLKASHSPPLNLPNN SHGITDFSSNSSAEHSLGSLKPTSTISTSPPLIHSEFVSKVPWNA
PIADEDLLPISAHPNATPALSS ENFTWSLVNDTVKTPDNSSITVSILSSEPTSPSVTPLIVE
PSGWLTTNSDSFTGFTPYQEKTTLQPTLKFTNNSKLFNPNTSDPQKENRNTGIVFGAILGAIL
GVSLTLVGYLLCGKRKTDSFSHRRLYDDRNEPVLRLDNAPEPYDVSFGNSSYYNPTLNDSA
MP ESEENARDGIPMDDIPPLRTSV

Signal peptide:

amino acids 1-23

Transmembrane domain:

amino acids 235-262

N-glycosylation site.

amino acids 30-34, 61-65, 79-83, 90-94, 148-152, 155-159,
163-167, 218-222, 225-229, 298-302, 307-311

FIGURE 25

AACAGGATCTCCTCTTGCAGTCTGCAGCCCAGGACGCTGATTCCAGCAGCGCCTTACCGCGC
 AGCCCGAAGATTCACT**ATG**GTGAAAATCGCCTTCAATACCCCTACCGCCGTGCAAAAGGAGG
 AGGCGCGGCAAGACGTGGAGGCCCTCCTGAGCCGCACGGTCAGAACTCAGATACTGACCGGC
 AAGGAGCTCCGAGTTGCCACCCAGGAAAAAGAGGGCTCCTCTGGGAGATGTATGCTTACTCT
 CTTAGGCCTTTTCATTCATCTTGGCAGGACTTATTGTTGGTGGAGCCTGCATTTACAAGTACT
 TCATGCCCAAGAGCACCATTACCCTGGAGAGATGTGCTTTTTTTGATTCTGAGGATCCTGCA
 AATTCCCTTCGTGGAGGAGAGCCTAACTTCCTGCCTGTGACTGAGGAGGCTGACATTCGTGA
 GGATGACAACATTGCAATCATTGATGTGCCTGTCCCCAGTTTCTCTGATAGTGACCCCTGCAG
 CAATTATTCATGACTTTGAAAAGGGAATGACTGCTTACCTGGACTTGTTGCTGGGGAACCTGC
 TATCTGATGCCCCCTCAATACTTCTATTGTTATGCCTCCAAAAAATCTGGTAGAGCTCTTTGG
 CAAACTGGCGAGTGGCAGATATCTGCCTCAAACCTTATGTGGTTCGAGAAGACCTAGTTGCTG
 TGGAGGAAATTCGTGATGTTAGTAACCTTGGCATCTTTATTTACCAACTTTGCAATAACAGA
 AAGTCCTTCCGCCTTCGTGCGCAGAGACCTCTTGCTGGGTTTCAACAAACGTGCCATTGATAA
 ATGCTGGAAGATTAGACACTTCCCCAACGAATTTATTGTTGAGACCAAGATCTGTCAAGAG**T**
AAGAGGCAACAGATAGAGTGTCTTGTAATAAGAAGTCAGAGATTTACAATATGACTTTAA
 CATTAAGGTTTATGGGATACTCAAGATATTTACTCATGCATTTACTCTATTGCTTATGCTTT
 AAAAAAAGGAAAAAAAAAAAAAAAACTACTAACCCTGCAAGCTCTTGTCAAATTTTAGTTTAAAT
 TGGCATTGCTTGTTTTTTTGAACTGAAATTACATGAGTTTCATTTTTTCTTGCATTTATAG
 GGTTTAGATTTCTGAAAGCAGCATGAATATATCACCTAACATCCTGACAATAAATTCATCC
 GTTGTTTTTTTTGTTTGTTTGTTTTTCTTTTCCTTTAAGTAAGCTCTTTATTCATCTTATG
 GTGGAGCAATTTTAAATTTGAAATATTTTAAATTGTTTTTGAACTTTTTGTGTAAAATATA
 TCAGATCTCAACATTGTTGGTTTCTTTTGTTTTTTCATTTTGTACAACTTTCTTGAATTTAGA
 AATTACATCTTTGCAGTTCTGTTAGGTGCTCTGTAATTAACCTGACTTATATGTGAACAATT
 TTCATGAGACAGTCATTTTTAACTAATGCAGTGATTCTTTCTCACTACTATCTGTATTGTGG
 AATGCACAAAATTGTGTAGGTGCTGAATGCTGTAAGGAGTTTAGGTTGTATGAATTCTACAA
 CCCTATAATAAATTTTACTCTATACAAAAAAAAAAAAAAAAAAAA

FIGURE 26

MVKIAFNTPTAVQKEEARQDVEALLSRTVRTQILTGKELRVATQEKEGSSGRCLTLLGLSF
ILAGLIVGGACIYKYFMPKSTIYRGEMCFFDSEDPANSLRGGEPNFLPVTEEADIREDDNIA
IIDVPVPVSFSDSDPAAIHDFEKGMTAYLDLLLGNCYLMPLNTSIVMPKKNLVELFGKLASG
RYLPQTYVVREDLVAVEEIRDVSNLGIFIYQLCNNRKSFRLRRDLLLGFNKRAIDKCWKIR
HFPNEFIVETKICQE

amino acids 53-75

amino acids 166-170

amino acids 35-39, 132-136, 134-138

amino acids 66-72, 103-109

amino acids 63-74

FIGURE 27

GGAGGAGGGAGGGCGGGCAGGCCAGCCAGAGCAGCCCCGGGCACCCAGCACGGACTCTCT
CTTCCAGCCCAGGTGCCCCCCTCTCGCTCCATTCGGCGGGAGCACCCAGTCCTGTACGCC
AAGGAACTGGTCCTGGGGGCACCA**ATG**GTTTTCGGCGGCAGCCCCCAGCCTCCTCATCCTTCTG
TTGCTGCTGCTGGGGTCTGTGCTGCTACCGACGCCCGCTCTGTGCCCTGAAGGCCACGTT
CCTGGAGGATGTGGCGGGTAGTGGGGAGGCCGAGGGCTCGTCGGCCTCCTCCCCGAGCCTCC
CGCCACCCTGGACCCCGGCCCTCAGCCCCACATCGATGGGGCCCCAGCCCACAACCCTGGGG
GGCCCATCACCCCCACCAACTTCCTGGATGGGATAGTGGACTTCTTCCGCCAGTACGTGAT
GCTGATTGCTGTGGTGGGCTCCCTGGCCTTTCTGCTGATGTTTCATCGTCTGTGCCGCGGTCA
TCACCCGGCAGAAGCAGAAGGCCTCGGCCTATTACCCATCGTCCTTCCCCAAGAAGAAGTAC
GTGGACCAGAGTGACCGGGCCGGGGGCCCCCGGGCCTTCAGTGAGGTCCCCGACAGAGCCCC
CGACAGCAGGCCCCGAGGAAGCCCTGGATTCTCCCGGCAGCTCCAGGCCGACATCTTGGCCG
CCACCCAGAACCTCAAGTCCCCCACCAGGGCTGCACTGGGCGGTGGGGACGGAGCCAGGATG
GTGGAGGGCAGGGGCGCAGAGGAAGAGGAGAAGGGCAGCCAGGAGGGGGACAGGAAGTCCA
GGGACATGGGGTCCCAGTGGAGACACCAGAGGCGCAGGAGGAGCCGTGCTCAGGGGTCCCTTG
AGGGGGCTGTGGTGGCCGGTGAGGGCCAAGGGGAGCTGGAAGGGTCTCTCTTGTAGCCAG
GAAGCCAGGGACCAGTGGGTCCCCCGAAAGCCCCTGTGCTTGCAGCAGTGTCCACCCAG
TGTC**TAA**CAGTCCTCCCGGGCTGCCAGCCCTGACTGTCGGGCCCCCAAGTGGTCACCTCCCC
GTGTATGAAAAGGCCTTCAGCCCTGACTGCTTCCTGACACTCCCTCCTTGGCCTCCCTGTGG
TGCCAATCCCAGCATGTGCTGATTCTACAGCAGGCAGAAATGCTGGTCCCCGGTGCCCCGGA
GGAATCTTACCAAGTGCCATCATCCTTCACCTCAGCAGCCCCAAAGGGCTACATCCTACAGC
ACAGCTCCCCTGACAAAGTGAGGGAGGGCACGTGTCCCTGTGACAGCCAGGATAAAACATCC
CCCAAAGTGCTGGGATTACAGGCGTGAGCCACCGTGCCCGGCCCAAACCTACTTTTTTAAACA
GCTACAGGGTAAAATCCTGCAGCACCCACTCTGGAAAATACTGCTCTTAATTTTCCTGAAGG
TGGCCCCCTGTTTCTAGTTGGTCCAGGATTAGGGATGTGGGGTATAGGGCATTTAAATCCTC
TCAAGCGCTCTCCAAGCACCCCCGGCCTGGGGGTGAGTTTCTCATCCCGCTACTGCTGCTGG
GATCAGGTTGAATGAATGGA**ACT**CCTCCTGTCTGGCCTCCAAGCAGCCTAGAAGCTGAGGG
GCTGTGTTTGAGGGGACCTCCACCCTGGGGAAGTCCGAGGGGCTGGGGAAGGGTTTCTGACG
CCCAGCCTGGAGCAGGGGGGCCCTGGCCACCCCTGTTGCTCACACATTGTCTGGCAGCCTG
TGTCACAATATTCGT**CAGT**CCTCGACAGGGAGCCTGGGCTCCGTCTGCTTTAGGGAGGCT
CTGGCAGGAGGTCTCTCCCCATCCCTCCATCTGGGGCTCCCCAACCTCTGCACAGCTCT
CCAGGTGCTGAGATATAATGCACCAGCACAAATAAACCTTTATTCCGGCCTGAAAAAAAAAAAA
AAA

FIGURE 28

></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA58852

><subunit 1 of 1, 283 aa, 1 stop

><MW: 29191, pI: 4.52, NX(S/T): 0

MVSAAAPSLILLILLLLLLGSPATDARSVPLKATFLEDVAGSGEAGSSASSPSLPPPWTPAL
SPTSMGPQPTTLGGPSPPTNFLDGIVDFFRQYVMLIAVVGSLAFLLMFIVCAAVITRQKQKA
SAYYPSSFPKKKYVDQSDRAGGPRAFSEVPDRAPDSRPEEALDSSRQLQADILAATQNLKSP
TRAALGGGDGARMVEGRGAEEEEKGSQEGDQEVQGHGVPVETPEAQEEPCSGVLEGAVVAGE
GQGELEGSLLLAQEAQGPVGPPEPCACSSVHPSV

Signal peptide:

amino acids 1-25

Transmembrane domain:

amino acids 94-118

N-myristoylation site.

amino acids 18-24, 40-46, 46-52, 145-151, 192-198, 193-199,
211-217, 238-244, 242-248

FIGURE 29

GTGGACTCTGAGAAGCCCAGGCAGTTGAGGACAGGAGAGAGAAGGCTGCAGACCCAGAGGGA
 GGGAGGACAGGGAGTTCGGAAGGAGGAGGACAGAGGAGGGCACAGAGACGCAGAGCAAGGGCG
 GCAAGGAGGAGACCCTGGTGGGAGGAAGACACTCTGGAGAGAGAGGGGGCTGGGCAGAG**ATG**
 AAGTTCCAGGGGCCCTGGCCTGCCTCCTGCTGGCCCTCTGCCTGGGCAGTGGGGAGGCTGG
 CCCCCTGCAGAGCGGAGAGGAAAGCACTGGGACAAATATTGGGGAGGCCCTTGGACATGGCC
 TGGGAGACGCCCTGAGCGAAGGGGTGGGAAAGGCCATTGGCAAAGAGGCCGGAGGGGCAGCT
 GGCTCTAAAGTCAGTGAGGCCCTTGGCCAAGGGACCAGAGAAGCAGTTGGCACTGGAGTCAG
 GCAGGTTCCAGGCTTTGGCGCAGCAGATGCTTTGGGCAACAGGGTCGGGGAAGCAGCCCATG
 CTCTGGGAAACACTGGGCACGAGATTGGCAGACAGGCAGAAGATGTCATTTCGACACGGAGCA
 GATGCTGTCCGCGGCTCCTGGCAGGGGGTGCCTGGCCACAGTGGTGCTTGGGAACTTCTGG
 AGGCCATGGCATCTTTGGCTCTCAAGGTGGCCTTGGAGGCCAGGGCCAGGGCAATCCTGGAG
 GTCTGGGGACTCCGTGGGTCCACGGATACCCCGGAAACTCAGCAGGCAGCTTTGGAATGAAT
 CCTCAGGGAGCTCCCTGGGGTCAAGGAGGCAATGGAGGGCCACCAAACCTTTGGGACCAACAC
 TCAGGGAGCTGTGGCCCAGCCTGGCTATGGTTCAGTGAGAGCCAGCAACCAGAATGAAGGGT
 GCACGAATCCCCCACCATCTGGCTCAGGTGGAGGCTCCAGCAACTCTGGGGGAGGCAGCGGC
 TCACAGTCGGGCAGCAGTGGCAGTGGCAGCAATGGTGACAACAACAATGGCAGCAGCAGTGG
 TGGCAGCAGCAGTGGCAGCAGCAGTGGCAGCAGCAGTGGCGGCAGCAGTGGCGGCAGCAGTG
 GTGGCAGCAGTGGCAACAGTGGTGGCAGCAGAGGTGACAGCGGCAGTGAGTCCTCCTGGGGA
 TCCAGCACCGGCTCCTCCTCCGGCAACCACGGTGGGAGCGGCGGAGGAAATGGACATAAACC
 CGGGTGTGAAAAGCCAGGGAATGAAGCCCGCGGGAGCGGGGAATCTGGGATTGAGGGCTTCA
 GAGGACAGGGAGTTTCCAGCAACATGAGGGGAAATAAGCAAAGAGGGCAATCGCCTCCTTGA
 GGCTCTGGAGACAATTATCGGGGGCAAGGGTCGAGCTGGGGCAGTGGAGGAGGTGACGCTGT
 TGGTGGAGTCAATACTGTGAACTCTGAGACGTCTCCTGGGATGTTTAACTTTGACACTTTCT
 GGAAGAATTTTAAATCCAAGCTGGGTTTCATCAACTGGGATGCCATAAACAAGGACCAGAGA
 AGCTCTCGCATCCCG**TGA**CCTCCAGACAAGGAGCCACCAGATTGGATGGGAGCCCCCAGACT
 CCCTCCTTAAAACACCACCTCTCATCACTAATCTCAGCCCTTGCCCTTGAAATAAACCTTA
 GCTGCCCCACAAA
 AA

1000
 900
 800
 700
 600
 500
 400
 300
 200
 100
 0

FIGURE 30

></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA59212

><subunit 1 of 1, 440 aa, 1 stop

><MW: 42208, pI: 6.36, NX(S/T): 1

MKFQGPLACLLALCLGSGEAGPLQSGEESTGTNIGEALGHGLGDALSEG VGKAIGKEAGGA
 AGSKVSEALGQGTREAVGTGVRQVPFGAADALGNRVGEAAHALGNTGHEIGRQAEDVIRHG
 ADAVRGSWQGVPGHSGAWETSGGHGIFGSQGGGLGGQGNPGGLGTPWVHGYPGNSAGSFGM
 NPQGAPWQGGNGGPPNFGTNTQGAVAQPGYGSVRASNQNEGCTNPPPSGSGGGSSNSGGGS
 GSQSGSSGSGSNGDNNNGSSSGSSSGSSSGSSSGSSSGSSSGSSGNSGGSRGDSGSESSW
 GSSTGSSSGNHGGSGGGNGHKPGCEKPGNEARGSGESGIQGFRGQGVSSNMREISKEGNRLL
 GGSGDNYRGQGSWSGGGDAVGGVNTVNSETSPGMFNFDTFWKNFKSKLGFINWDAINKDQ
 RSSRIP

Signal peptide:

amino acids 1-21

N-glycosylation site.

amino acids 265-269

Glycosaminoglycan attachment site.

amino acids 235-239, 237-241, 244-248, 255-259, 324-328, 388-392

Casein kinase II phosphorylation site.

amino acids 26-30, 109-113, 259-263, 300-304, 304-308

N-myristoylation site.

amino acids 17-23, 32-38, 42-48, 50-56, 60-66, 61-67, 64-70,
 74-80, 90-96, 96-102, 130-136, 140-146, 149-155, 152-158,
 155-161, 159-165, 163-169, 178-184, 190-196, 194-200, 199-205,
 218-224, 236-242, 238-244, 239-245, 240-246, 245-251, 246-252,
 249-252, 253-259, 256-262, 266-272, 270-276, 271-277, 275-281,
 279-285, 283-289, 284-290, 287-293, 288-294, 291-297, 292-298,
 295-301, 298-304, 305-311, 311-317, 315-321, 319-325, 322-328,
 323-329, 325-331, 343-349, 354-360, 356-362, 374-380, 381-387,
 383-389, 387-393, 389-395, 395-401

Cell attachment sequence.

amino acids 301-304

ACCGCGTCCCTCCGGTCTCTGGATGTGCGGACTCTGCTGCAGCGAGGGCTGCAGGCCCGCCGGGCGGTGCTCACCC
 TGCCCTGGCTGGTGGAGTTTCTCTCCTTTGCTGACCATGTGTTCCTTGCTGGAATATTACCGGGACATCTTCA
 CTCTCCTGCTGCGCCTGCACCGGAGCTTGGTGTTGTGCGAGGAGAGTGAGGGGAAGATGTTGTTTCTTCTGGAAGAGG
 TGCTGCTACTTGTCTGTCTGGGCTGGCTTTTCCAGATTCCACAGTCCCTGAGGACTTGTCTTTCTGGAAGAGG
 GTCCCTCATATGCTTTGAGTGGACACAGTAGCCCCAGACATGGCTTGACCAATGCGCCTGTGGTGGACAGCAG
 AGCTGCTCTACACCTGCTGCCCTACATCGGAGAGCTCCGGAACATGCTCGCTTCGTGGGTGTACGGCAGTAGTG
 GACGGAGTGGGGGCTTCATGAGGAAAAATCACCCCCACCACTACCACCAGCCTGGGAGCCAGCCTTCCCAGACCA
 GCCAGGGGCTGCAGGCACAGCTCGCCAGGCCTTTTTCACAACCAGCCGCCCTCCTTGCGCCGGACCGTAGAGT
 TCGTGGCAGAAAAGAAATTGGATCAAACCTGTGTCAAACATATCAAGGCTACACTGGTGGCAGATCTGGTGCGCCAG
 CAGAGTCACTTCTCAAGAGCAGCCTGGTGACACAGGGAGAGGAAGGGGGAGACCAGCCAGCTGTTGGAGATCT
 TGTGTTCCAGCTGTGCCCTCAGCGGGCCAGGCATTGCCCTTGGGCGGGAGTTCGTCTCAAGGAAGAGCCCTC
 GGGCTGTGCGGGCGCTGCTTCCAGAGGAGACCCCGGACGCCGTTCTGAGCAGTGCAGAGAACATTGCTGTGGGGC
 TTGCAACAGAGAAAAGCCTGTGCTTGGCTGTGAGCCAACATCACAGCACTGATCAGGAGGGGAGGTGAAAGCAGCAG
 TGAGTCGCACACTTCGAGCCAGGGTCTGAACCTGCTGCCCGGGGGGAGCGGAGGGGCTGCTCCCGCGCTCGAC
 GTGCTCTCCTTGCCCGTGGGGCCACGGGACCCCTGACGAGGGAGTCTCCCCAGAGCATCTGGAACAGCTCCTAGGC
 CAGCTGGGCCAGACGCTGCGGTGCCGCCAGTTCTGTGCCACCTGCTGAGCAGCATCTGGCAAAGTGCTCTGTG
 GAGTTAGCTTCCCTCCTCGTTGCAGATCAAATTCCTATCTAGGGCCCCCGGCACAGTACAGGCTGGAGAGAGGG
 CAGGCTCGAAGGCTTCTGCAATGCTGCTTTCTGTGGAAGAAAGCTTCAGGGGCGGTTCCGCTGCAGCTG
 CTGCTGAGCCCAAGAAATGTGGGGCTTCTGGCAGACACAAGGCCAAGGGAGTGGAAGTCTGCTGCTATTCTTGCTA
 CGGGAGCTGGTGGAAGGGTCTGATGGGACGGATGGAGATAGAGGCTGCCCTGGGCAGCCTCCACCAGGCCAG
 TGGCCAGGGGACTTTGCTGAAGAATTAGCAACACTGTCTAATCTGTTTCTAGCCGAGCCCCACCTGCCAGAACCC
 CAGCTAAGAGCCTGTGAGTTGGTGACGCCAAAACCGGGGCACTGTGCTGGCCCAGAGCTAGGGCTGAGAAGTGGCC
 TGTCTTGGGCATTGCACCAGAACCTTGGACCCCGCTCAGGAGGAGGCCAAGTGCCAATGCACACCTCAC
 TGGTTGGGGTGAGCTGGGTCTACAGTCAAGTCTGCTGCTTAAGGGTGCTCACTGCTGGCATCCACCACGCGA
 ATCTAGAGGAAGGAGAGTTGGCCTGATTTGGGATTATGGCAGAAAAGTCCAGAGATGCCAGTCTTGAGTAGAA
 GAGGTGGTGTTTGTATTCTCTTGATACTAAATGAAATGAGGTGTGTGGGCTTGTCAACACAGAAATCAAGCCT
 CATTTGCTATCCAGCATCTCTTAAACTTTGTAGTCTTGGAATTATGACAGAGGCAAAATGACTCCTGCTTAAC
 TTATGAAGAAAGTTAAACCATGAATCTTTGGGAGTCTACATTTTCTTATCACCAGGAGCTGGACTGCCATCTCCTT
 ATAAATGCTTAACACAGACCGGGTCTGGTGGCTCATGCTGTAATCCAGCACTTTGAGAGGCTGAGGTGGCG
 GACTGCCTGAGTGAGGAAATCAAGACAGCCCTGGCCAACATGGCAAAACCCCATCTCTACTAAAAATAAAAAA
 TTATTAGCTGGGCATGGTGGTGTGTGCTGTAAATCCCAGCTACTCAGGAGGATGAGGCAGGAGACCTGCTTGAAC
 CTGGAGGTGGAGGTTGCAGTGAGCCGAGGTGCACCACTGCACTCCAGTCTGGGTAAACAGAGCGAGACTTTCTAG
 AAAAAAGCTTAACAAACAGATAAGGTAGGACTCAACCAACTGAAACCTGACTTTCCCCCTGTACCTTCAGCCCTG
 TGCAGGTAGTAACCTCTTGAGACCTCTCCCTGACCGGACCAAGCAGAGGCGATTTAGAGCTTTTTAGAATAAA
 CTGTTTCTTTTAAAAAAGGCTTTTATTAATAATCTCCCCACAGTAGGCTCTGCAATCTGCCACAGCTC
 TGGGGCGTGTCTGTAGGGAAAGGCCCTGTTTTCTTGGAGCGGGGCTGGGCTTGTCCATGGGTCCGCGGAGCTG
 GCCGTGCTTGGCGCCTGGCGTGTGTCTAGCTGCTTCTTGCCGGGCACAGAGCTGCGGGGTCTGGGGGCACCGG
 AGCTAAGAGCAGGCTCTGGTGACGGGGTGGAGGCCTGTCTTAACCGACACCTGAGGTGCTCCTGAGATGCTG
 GGTCCACCTTGTGTCACGGGGAGCAGCTGTGGCCGGTGCTCCTTCTAGGCCAGTCTGGGGAAGCTAAGCTC
 GGGCCCTTCTTTGCAAGACCGAGATGGGTGGGTGGGGACTCATGGGAATGGCTTGAGGAGTACGTGT
 GAAGAGGGCGCCGGTTTGTGGCTGCAGCGGCCTGGAGCGCCTCTCTCCTGAGCCTCAGTTTCCCTTTCCGTCTA
 ATGAAGAACATGCCGTCTCGGTGTCTCAGGGCTATTAGGACTTGCCTCAGGAAGTGGCCTTGGACGAGCGTCT
 GTTATTTTCACAACTGTCTGCGACGTTGGCTGGGCACGTCTGGAATGGCCCATGTCTCTGCTGCGTGGAC
 TCTCGCGTGGGAGTGCGCAGCCAGAGGCGGGGCCAGACGTGCGCCTGGGGGTGAGGGGAGGCGCCCCGGGAGGG
 CCTCACAGGAAGTTGGCTCCCGCACCAAGGCGGGCGGGCTCCCGCCGCCCGCCACCAACCGTCCAGG
 GGCCGCTAGACAAAGTGAAGTCGCGCTTGGGCTGCGTGGCGCAGCAGGTAGCCCTTGATGCAGTGCGGCAGCGG
 TCGTCCGCCAGCTGGAAGCAGCGCCCGTCCACAGCACGAACAGCCGTTGCGCT

FIGURE 32

MCFLNKLLLLLAVLGWLFQIPTVPEDLFFLEEGPSYAFEVDTVAPEHGLDNAPVVDQQLLYTC
CPYIGELRKLLASWVSGSSGRSGGFMRKITPTTTTSLGAQPSQTSQGLQAQLAQAFFHNQPP
SLRRTVEFVAERIGSNCVKHIKATLVADLVRQAESLLQEQLVTQGEEGGDPAQLLEILCSQL
CPHGAQALALGREFCQRKSPGAVRALLPEETPAAVLSSAENI AVGLATEKACAWLSANITAL
IRREVKAAVSRTLRAQGPEPAARGERRGCSRA

Signal peptide:

amino acids 1-18

N-glycosylation site.

amino acids 244-248

cAMP- and cGMP-dependent protein kinase phosphorylation site.

amino acids 89-93

Casein kinase II phosphorylation site.

amino acids 21-25, 167-171, 223-227

N-myristoylation site.

amino acids 100-106, 172-178, 207-213

Microbodies C-terminal targeting signal.

amino acids 278-282

FIGURE 33

TCCCTTGACAGGTCTGGTGGCTGGTTCGGGGTCTACTGAAGGCTGTCTTGATCAGGAACTG
 AAGACTCTCTGCTTTTGCCACAGCAGTTCCTGCAGCTTCCTTGAGGTGTGAACCCACATCCC
 TGCCCCCAGGGCCACCTGCAGGACGCCGACACCTACCCCTCAGCAGACGCCGGAGAGAA**ATG**
 AGTAGCAACAAAGAGCAGCGGTGAGCAGTGTTTCGTGATCCTCTTTGCCCTCATCACCATCCT
 CATCCTCTACAGCTCCAACAGTGCCAATGAGGTCTTCCATTACGGCTCCCTGCGGGGCCGTA
 GCCGCCGACCTGTCAACCTCAAGAAGTGGAGCATCACTGACGGCTATGTCCCCATTCTCGGC
 AACAAAGACACTGCCCTCTCGGTGCCACCAGTGTGTGATTGTGAGCAGCTCCAGCCACCTGCT
 GGGCACCAAGCTGGGCCCTGAGATCGAGCGGGCTGAGTGTACAATCCGCATGAATGATGCAC
 CCACCACTGGCTACTCAGCTGATGTGGGCAACAAGACCACCTACCGCGTCGTGGCCCATTC
 AGTGTGTTCCGCGTGCTGAGGAGGCCCCAGGAGTTTGTCAACCGGACCCCTGAAACCGTGTT
 CATCTTCTGGGGGGCCCCGAGCAAGATGCAGAAAGCCCCAGGGCAGCCTCGTGCGTGTGATCC
 AGCGAGCGGGCCTGGTGTTCGCCAACATGGAAGCATATGCCGTCTCTCCCGGCCGCATGCGG
 CAATTTGACGACCTCTTCCGGGGTGAGACGGGCAAGGACAGGGAGAAGTCTCATTCGTGGTT
 GAGCACAGGCTGGTTTACCATGGTGATCGCGGTGGAGTTGTGTGACCACGTGCATGTCTATG
 GCATGGTCCCCCCCCAACTACTGCAGCCAGCGGCCCGCCTCCAGCGCATGCCCTACCACTAC
 TACGAGCCCAAGGGGGCCGGACGAATGTGTACCTACATCCAGAATGAGCACAGTCGCAAGGG
 CAACCACCACCGCTTCATCACCGAGAAAAGGGTCTTCTCATCGTGGGCCCAGCTGTATGGCA
 TCACCTTCTCCACCCCTCCTGGACCT**TAG**GCCACCCAGCCTGTGGGACCTCAGGAGGGTCAG
 AGGAGAAGCAGCCTCCGCCCAGCCGCTAGGCCAGGGACCATCTTCTGGCCAATCAAGGCTTG
 CTGGAGTGTCTCCAGCCAATCAGGGCCTTGAGGAGGATGTATCCTCCAGCCAATCAGGGCC
 TGGGGAATCTGTTGGCGAATCAGGGATTTGGGAGTCTATGTGGTTAATCAGGGGTGTCTTTC
 TTGTGCAGTCAGGGTCTGCGCACAGTCAATCAGGGTAGAGGGGGTATTTCTGAGTCAATCTG
 AGGCTAAGGACATGTCTTTCCCATGAGGCCTTGTTTCAGAGCCCCAGGAATGGACCCCCCA
 ATCACTCCCCACTCTGCTGGGATAATGGGGTCTGTCCCAGGAGCTGGGAACCTTGGTGTG
 CCCCCTCAATTTCCAGCACCAAGAGAGATTGTGTGGGGGTAGAAGCTGTCTGGAGGCC
 GGCCAGAGAATTTGTGGGGTTGTGGAGGTTGTGGGGGCGGTGGGGAGGTCCCAGAGGTGGGA
 GGCTGGCATCCAGGTCTTGGCTCTGCCCTGAGACCTTGGACAAACCCTTCCCCCTCTCTGGG
 CACCCTTCTGCCCACACCAAGTTTCCAGTGCGGAGTCTGAGACCCCTTTCCACCTCCCCTACAA
 GTGCCCTCGGGTCTGTCTCCCCGTCTGGACCTCCCAGCCACTATCCCTTGCTGGAAGGCT
 CAGCTCTTTGGGGGGTCTGGGGTGACCTCCCCACCTCCTGGAAAACCTTTAGGGTATTTTTGC
 GCAAACCTCCTTCAGGGTTGGGGGACTCTGAAGGAAACGGGACAAAACCTTAAGCTGTTTTCT
 TAGCCCCTCAGCCAGCTGCCATTAGCTTGGCTCTTAAAGGGCCAGGCCTCCTTTTCTGCCCT
 CTAGCAGGGAGGTTTTCCAACCTGTTGGAGGCGCCTTTGGGGCTGCCCTTTGTCTGGAGTCA
 CTGGGGGCTTCCGAGGGTCTCCCTCGACCTCTGTCTGCTGGGATGGCTGTGCGGAGCTGT
 ATCACCTGGGTCTGTCCCCCTGGCTCTGTATCAGGCACTTTATTAAAGCTGGGCCCTCAGTGG
 GGTGTGTTTGTCTCCTGCTCTTCTGGAGCCTGGAAGGAAAGGGCTTCAGGAGGAGGCTGTGA
 GGCTGGAGGGACCAGATGGAGGAGGCCAGCAGCTAGCCATTGCACACTGGGGTGATGGGTGG
 GGGCGGTGACTGCCCCAGACTTGGTTTTGTAAATGATTTGTACAGGAATAAACACACCTACGC
 TCCGGAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA

FIGURE 34

MSSNKEQRSVAVFVILFALITILILYSSNSANEVFHYGSLRGRSRRPVNLKKWSITDGYVPIL
GNKTLPSRCHQCVIVSSSSHLLGTKLGPEIERAECTIRMNDAPTTGYSADVGNKTTYRVVAH
SSVFRVLRRPQEFVNRTPETVFIFWGPPSKMQKPQGSLVRVIQRAGLVFPNMEAYAVSPGRM
RQFDDLFRGETGKDREKSHSWLSTGWFTMVIARELCHVHVYGMVPPNYCSQRPRLQRMPIH
YYEPKGPDECVTYIQNEHSRKGNNHFRFITEKRVFSSWAQLYGITFSHPSWT

Signal peptide:

amino acids 1-29

Transmembrane domain:

amino acids 9-31 (type II)

N-glycosylation site.

amino acids 64-68, 115-119

cAMP- and cGMP-dependent protein kinase phosphorylation site.

amino acids 50-54

Casein kinase II phosphorylation site.

amino acids 3-7, 29-33, 53-57, 197-201

Tyrosine kinase phosphorylation site.

amino acids 253-262

N-myristoylation site.

amino acids 37-43, 114-120, 290-294

GTTCTCATAGTTGGCGTCTCTAAAGGAAAAACACTAAAATGAGGAACCTCAGCGGACCGGGAGCGACGACGCTT
 GAGGGAAGCATCCCTAGCTGTTGGCGCAGAGGGGCGAGGCTGAAGCCGAGTGGCCCCAGGTTGTCTGAGGGGCTGG
 CGCAAAGGTGAAGAGTTTCAAGAACAGCTTCTTCCGAAACCCATGACCCATGAAGTCTTGTGCAGATTTATACCGT
 TCTAGGGGTAGCAGCTCGAAACTAGAAAGAGTGGAGTGTGGCAGGGACGGCAGTATCTCTTTGTGTGACCCCTGGC
 GGCCATATGGGACGTTGGCTTCAGACCTTTTGTGATACACCATGCTGCGTGGGACGATGACGGCGTGGAGAGGAATG
 AGGCCGTAGGTTACACTGGCTTGGCTCCTCCTAGCCACAGCAGGCTGCCTTGTCTGACTTGAACGAGGTCCCTCAG
 GTCACCCGTCCAGCCTGCGCTCCACCGTCCGAAAGCCGGAGGCATGTGATCTTGGGCTGCGTGGTGGAACTCCCA
 AGGATGAATGTAACCTGGCGCTGAATGGAAAGGAGCTGAATGGCTCGGATGATGCTCTGGGTGTCCTCATCACC
 CACGGGACCCCTCGTCACTCACTGCCCCTTAAACAACCACACTGTGGGACGGTACCAGTGTGTGGCCCCGGATGCCCTGCC
 GGGGCTGTGGCCAGCGTGCCAGCCACTGTGACACTAGCCAATCTCAGGACTTCAAGTTAGATGTGCAGCAGCT
 ATTGAAGTGGATGAGGGAAACACAGCAGTCATTGGCTGCCACCTGCCCTGAGAGGCCACCCCCAAGGCCAGGTCCGG
 TACAGCGTCAACAAGAGTGGCTGGAGGCTCCAGAGGTAACCTGATCATGCCCTCAGGGAACCTCCAGATT
 GTGAATGCCAGCCAGGAGGACGAGGCGCATGTACAAGTGTGCAGCCTACAACCAGTGAACCCAGGAAGTGAAACC
 TCCGGCTCCAGCGACAGGCTACGTGTGCGCCGCTCCACCGCTGAGGCTGCCCGCATCATCTACCCCCCAGAGGCC
 CAAACCATCATCGTCAACAAAGGCCAGAGTCTCATTTCTGGAGTGTGTGGCCAGTGGAAATCCCCACCCCCACGGGTC
 ACCTGGGCGAAGGATGGGTCCAGTGTACCCGGCTACAACAAGACGCGCTCTCTGCTGAGCAACCTCTCATCGAC
 ACCACGAGCGAGGAGGACTCAGGCACCTACCGCTGCATGGCCGACAATGGGGTTGGGACGCCCGGGGCGAGCGGTC
 ATCCTCTACAATGTCCAGGTGTTTGAACCCCTGAGGTCACCATGGAGCTATCCCAGCTGGTTCATCCCTGGGGC
 CAGATGCCCCAAGCTTACCTGTGAGGTGCGTGGGAACCCCCCGCCCTCCGTGCTGTGGCTGAGGAATGCTGTGCC
 CTCACTCCAGCCAGCCCTCCGGCTCTCCCGCAGGGCCCTGCGCGTGCTCAGCATGGGGCTGAGGACGAAGG
 GTCTACCACTGCATGGCCGAGAAGTGTGGAGCGCCCATGCGGTAGTCCAGCTGCGGACCTCCAGGCCAAGC
 ATAACCCCAAGGCTATGGCAGGATGCTGAGCTGGCTACTGGCACACCTCCTGTATCACCTCCAAACTCGGCAAC
 CCTGAGCAGATGCTGAGGGGGCAACCGGCGCTCCCGAGACCCCCAACGTCAGTGGGGCGCTGCTTCCCCGAAGTGT
 CCAGGAGAGAAGGGGCGGGGGCTCCCGCGAGGCTCCCATCATCTCAGCTCGCCCGGCACCTCCAAGACAGAC
 TCATATGAATGGTGTGGCGGCTCGGCATGAGGGCAGTGGCCGGGCGCCAATCCTCTACTATGTGGTGAACAC
 CGCAAGCAGGTCACAAATTCCTCTGACGATTGGACCATCTCTGGCATTCCAGCCAACCAGCACCGCCTGACCCCT
 ACCAGCTATGACCCCGGGAGCTTGTATGAAGTGGAGATGGCAGCTTACAACCTGTGCGGGAGAGGGGCGAGACGCC
 ATGGTCACCTTCCGAACCTGGACGGCGGCCAAACCAGATCATGGCCAGCAAGAGCAGCAGATCCAGAGAGAC
 GACCTGGAGCCGATCCCCAGAGCAGCAGCCAGCCAGACACCGGCCGCTCTCCCCCCAGAAGCTCCCGACAGG
 CCCACCATCTCCACGGCCTCCGAGACCTCAGTGTACGTGACCTGGATTCCCCGTGGGAATGGTGGGTTCCCAATC
 CAGTCCCTCCGTGTGGAGTACAAGAGAGCTAAAGAAAGTGGGAGACTGGATTCTGGCCACCAAGGCCATCCCCCA
 TCGCGCTGTCCTGGAGATCAGGGGCTAGAGAAAGGCACCTCCTACAAGTTTTCAGTCCGGGCTCTGAACATG
 CTGGGGAGAGCGAGGCCAGCCAGCGCCCCCTCTCGGCCCTACGTTGGTGTGGGCTACAGCGGTGCGGTGTACGAGG
 CCCGTGGCAGGTCTTATATCACCTTTCACGGATGCGGTCAATGAGACCACCATCATGCTCAAGTGGATGTACAT
 CCAGCAAGTTAACACAACACCCCAATCCATGGCTTTTATATCTATTATCGACCCACAGACAGTGACAATGATAGT
 GACTACAAGAAGGATATGGTGAAGGGGACAAGTGGCACTCCATCAGCCACCTCGAGCCAGAGACCTCTTAC
 GACATTAAGATGTCAGTGCCTTCAATGAAGGAGGGGAGAGCGAGTTACGAACGTCGATGATCTGTGAGACCAAGCT
 CGGAAGTCTTCTGGCCAGCCTGGTCGACTGCCACCCCCAACTCTGGCCCCACCACAGCCGCCCCCTTCTGAAACC
 ATAGAGCGGGCGGTGGGCACTGGGGCCATGGTGGCTCGCTCCAGCGACCTGCCCTATCTGATTGTGGGGTCTGT
 CTGGGCTCCATCGTTCTCATCATCTGTCACCTTTCATCCCTTCTGCTTGTGGAGGGCTGGTCTAAGCAAAAACAT
 ACAACAGACCTGGGTTTCTCTCGAAGTGGCTTCCACCCCTCGTGGCGTATACTATGGTGCCATTGGGAGGACTC
 CCAGGCCACCAGGCCAGTGGACAGCCCTACCTCAGTGGCATCAGTGGACGGGCTGTGCTAATGGGATCCACATG
 AATAGGGGCTGCCCCCTCGGCTGCACTGGGTACCCGGGATGAAGCCCCAGCAGCAGTCCCCAGGCGAGCTTCAG
 CAGCAGAGTGACACCAGCAGCTGCTGAGGACAGCCCATCTTGGCAATGGATGATGACCCCCAAGCTCACCATG
 ACAGGGGTGCCAAGTCTAGCCCGGACGAGGGCTCTTTCTTATACACACTGCCGACGACTCCACTCACCAGCTG
 CTGCAGCCCCATCAGACTGCTGCCAACGCCAGGAGCAGCCTGCTGCTGTGGGCCAGTCAGGGGTGAGGAGAGCC
 CCGACAGTCTGTCTGGAAGCAGTGTGGACCCCTCCATTCACTCAGGCGCCCATGCTGCTTGGGCTTGTG
 CCAGTTGAAGAGTGGACAGTCTGACTCTGCCAAGTGAAGTGGAGGAGCTGGTGTCCCGACACCCCGTAGGG
 GCCTACGTAGGACAGCACTGGAATGCAGTCTCCCGGGGCGACTGTCGCTGTGCTTTTGAACACCACCT
 CTCACAATTTAGGCAGAAGCTGATATCCAGAAAAGACTATATATTGTTTTTTTTTTAAAAAAGAAAGAAAA
 AGAGACAGAGAAAAATTGGTATTTTATTTTCTATTATAGCCATATTTATATATTTATGGACCTGTGAATAAATGTA
 TATGTTTTATAATTCTGGAGAGACATAAGGAGTCTACCCGTTGAGGTTGGAGAGGAAAAAAGAAGAGCTGCCA
 CCTAACAGGAGTCAACCCAGGAAAGCACCGCAGAGGCTGGCGCGGGACAGACTCTTAACCTGGGGCTCTGCAGTG
 GCAGGCGAGGCTGCAGGAGGCCACAGATAAGCTGGCAAGAGGAAGGATCCAGGCACATGGTTTCATCACGAGCA
 TGAGGGAACAGCAAGGGGACGGTATCACAGCCTGGAGACACCCACAGATGGCTGGATCCGGTGTCTACGGGAA
 ACATTTTCTTAAGATGCCATGAGAACAGCAACCAAGATGTGTACAGCACATGAGCATTAATAAACCTTCCAGAA
 CAATAATCCGTGGCAACATATCTCTGTAAAAACAAACACTGTAACTCTAAATAAATGTTTGTAGTCTTCCCTGTAAA

FIGURE 36

MLRGTMTAWRGMRPEVTLACLLLATAGCFADLNEVPQVTVQPASTVQKPGGTVILGCVVEPP
 RMNVTWRLNGKELNGSDDALGVLITHGTLVITALNNHTVGRYQCVARMPAGAVASVPATVTL
 ANLQDFKLDVQHVEVDEGNTAVIACHLPESHHPKAQVRYSVKQEWLEASRGNYLIMPSGNLQ
 IVNASQEDEGMYKCAAYNPVTQEVKTSGSSDRLRVRRSTAEAAARI IYPPEAQTTIIVTKGQSL
 ILECVASGIPPPRVTWAKDGSSVTGYNKTRFLLSNLLIDTTSEEDSGTYRCMADNGVGQPGA
 AVILYNVQVFEPPPEVTMELSQLVIPWGS AKLTCEVRGNPPPSVLWLRNAVPLISSQRLRLS
 RRALRVLSMGPEDEGVYQCMAENEVGSAAHAVVQLRTSRPSITPRLWQDAELATGTPPVSPSK
 LGNPEQMLRGQPALPRPPTSVGPASPKCPGEGKGQAPAEAPIILSSPRTSKTDSYELVWRPR
 HEGSGRAPILYYVVKHRKQVTNSSDDWTISGIPANQHRLTLTRLDPGSLYEVEMAAYNCAGE
 GQTAMVTFRTGRRPKPEIMASKEQQIQRDDPGASPQSSSQPDHGRLSPPEAPDRPTISTASE
 TSVYVTWIPRGNGGFPIQSFVEYKKLKKVGDWILATSAIPPSRLSVEITGLEKGTSYKFRV
 RALNMLGESEPSAPSRPYVVS GYSGRVYERP VAGPYITFTDAVNETTIMLKWMI PASNNNT
 PIHGFIYYRPTDSDNDSYKKDMVEGDKYWHSISHLQPETS YDIKMQC FNEGGESEFSNVM
 ICETKARKSSGQPGRLPPPTLAPPQPLPETIERPVGTGAMVARSSDLPLYLIVGVVLGSIVL
 IIVTFIPFCLWRAW SKQKHTTDLGFPR SALPPSCPYTMVPLGGLPGHQASGQPYLSGISGRA
 CANGIHMNRGCPSAAVGYPGMKPQQHC PGELQQQSDTSSLLRQTHLGNGYDPQSHQITRGPK
 SSPDEGSFLYTL PDDSTHQLLQPHHDCCQRQE QPAAVGQSGVRRAPDSPVLEAVWDPPFHSG
 PPCCLGLVPVEEVDSPDSCQVSGGDWCPQHVP GAYVGQEPGMQLSPGPLVRV SFETPPLTI

Signal peptide:

amino acids 1-30

Transmembrane domain:

amino acids 16-30 (type II), 854-879

FIGURE 37

CGGGAGGCTGGGTTCGTCATGATCCGGACCCCATTTGTCGGCCTCTGCCCATCGCCTGCTCCTC
 CCAGGCTCCCGCGGCCGACCCCGCGCAAC**ATG**CAGCCCACGGGCCGCGAGGGTTCCCGCGC
 GCTCAGCCGGCGGTATCTGCGGCGTCTGCTGCTCCTGCTACTGCTGCTGCTGCTGCGGCAGC
 CCGTAACCCGCGCGGAGACCACGCCGGGCGCCCCCAGAGCCCTCTCCACGCTGGGGCTCCCC
 AGCCTCTTCACCACGCCGGGTGTCCCCAGCGCCCTCACTACCCAGGCCTCACTACGCCAGG
 CACCCCCAAAACCTGGACCTTCGGGGTTCGCGCGCAGGCCCTGATGCGGAGTTTCCCACTCG
 TGGACGGCCACAATGACCTGCCCCAGGTCCTGAGACAGCGTTACAAGAATGTGCTTCAGGAT
 GTTAACCTGCGAAATTTACGCCATGGTCAGACCAGCCTGGACAGGCTTAGAGACGGCCTCGT
 GGGTGGCCAGTTCTGGTCAGCCTCCGTCTCATGCCAGTCCCAGGACCAGACTGCCGTGCGCC
 TCGCCCTGGAGCAGATTGACCTCATTCACCGCATGTGTGCCTCCTACTCTGAACCTCGAGCTT
 GTGACCTCAGCTGAAGGTCTGAACAGCTCTCAAAGCTGGCCTGCCTCATTGGCGTGNAGGG
 TGGTCACTCACTGGACAGCAGCCTCTCTGTGCTGCGCAGTTTCTATGTGCTGGGGGTGCGCT
 ACCTGACACTTACCTTCACCTGCAGTACACCATGGGCAGAGAGTTCCACCAAGTTCAGACAC
 CACATGTACACCAACGTCAGCGGATTGACAAGCTTTGGTGAGAAAGTAGTAGAGGAGTTGAA
 CCGCCTGGGCATGATGATAGATTTGTCTATGCATCGGACACCTTGATAAGAAGGGTCCTGG
 AAGTGTCTCAGGCTCCTGTGATCTTCTCCCACTCAGCTGCCAGAGCTGTGTGTGACAATTTG
 TTGAATGTTCCCGATGATATCCTGCAGCTTCTGAAGAACGGTGGCATCGTGATGGTGACACT
 GTCCATGGGGGTGCTGCAGTGCAACCTGCTTGCTAACGTGTCCACTGTGGCAGATCACTTTG
 ACCACATCAGGGCAGTCATTGGATCTGAGTTCATCGGGATTGGTGGAAATTATGACGGGACT
 GGCCGGTTCCCTCAGGGGCTGGAGGATGTGTCCACATACCCAGTCCTGATAGAGGAGTTGCT
 GAGTCGTASCTGGAGCGAGGAAGAGCTTCAAGGTGTCCTTCGTGGAAACCTGCTGCGGGTCT
 TCAGACAAGTGGAAGAGGTGAGAGAGGAGAGCAGGGCGCAGAGCCCCGTGGAGGCTGAGTTT
 CCATATGGGCAACTGAGCACATCCTGCCACTCCCACCTCGTGCCCTCAGAATGGACACCAGGC
 TACTCATCTGGAGGTGACCAAGCAGCCAACCAATCGGGTCCCCTGGAGGTCCTCAAATGCCT
 CCCCATACCTTGTTCCAGGCCTTGTGGCTGCTGCCACCATCCCAACCTTCACCCAGTGGCTC
 TGCT**TGA**CACAGTCGGTCCCCGCAGAGGTCAGTGTGGCAAAGCCTCACAAAGCCCCCTCTCCT
 AGTTCATTACAAAGCATATGCTGAGAATAAACATGTTACACATGGAAAA

FIGURE 38

></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA59817

><subunit 1 of 1, 487 aa, 1 stop, 2 unknown

><MW: 53569.32, pI: 7.68, NX(S/T): 5

MQPTGREGSRALSRRYLRRLLLLLLLLLLLLRQPVTRAETTPGAPRALSTLGSPSLFTTPGVPS
 ALTTPGLTTPGTPKTLDLRGRAQALMRSFPLVDGHNDLPQVLRQRYKNVLQDVNLRNFSHGQ
 TSLDRLRDGLVGAQFWASVSCQSQDQTAVRLALEQIDLIHRMCASYSELELV TSAEGLNSS
 QKLACLIGVXGGHSLDSSLSVLRSFYVLGVRYLTLTFTCSTPWAESSTKFRHHMYTNVSGLT
 SFGEKVVEELNRLGMMIDLSYASDTLIRRVLEVSQAPVIFSHSAARAVCDNLLNVPDDILQL
 LKNGGIVMVTLSMGVLQCNLLANVSTVADHFDHIRAVIGSEFIGIGGNYDGTGRFPQGLEDV
 STYPVLIEELLSRXWSEELQGVLRGNLLRVFRQVEKVREESRAQSPVEAEFPYGQLSTSCH
 SHLVPQNGHQATHLEVTKQPTNRVPWRSSNASPYLVPGLVAAATIPTFTQWLC

Important features of the protein:

Signal peptide:

amino acids 1-36

Transmembrane domain:

amino acids 313-331

N-glycosylation sites.

amino acids 119-122, 184-187, 243-246 and 333-336

N-myristoylation sites.

amino acids 41-46, 59-64, 73-78, 133-138, 182-187, 194-199, 324-329, 354-359, 357-362, 394-399, 427-432 and 472-477.

Prokaryotic membrane lipoprotein lipid attachment site.

amino acids 136-146

FIGURE 39

TGCTAGGCTCTGTCCCACAATGCACCCGAGAGCAGGAGCTGAAAGCCTCTAACACCCACAGA
TCCCTCTATGACTGCAATGTGAGGTGTCCGGCTTTGCTGGCCCAGCAAGCCTGATAAGC**ATG**
AAGCTCTTATCTTTGGTGGCTGTGGTCTGGGTGTTTGCTGGTGCCCCAGCTGAAGCCAACAA
GAGTTCTGAAGATATCCGGTGCAAATGCATCTGTCCACCTTATAGAAACATCAGTGGGCACA
TTTACAACCAGAATGTATCCCAGAAGGACTGCAACTGCCTGCACGTGGTGGAGCCCATGCCA
GTGCCTGGCCATGACGTGGAGGCCTACTGCCTGCTGTGCGAGTGCAGGTACGAGGAGCGCAG
CACCACCACCATCAAGGTCATCATTTGTCATCTACCTGTCCGTGGTGGGTGCCCTGTTGCTCT
ACATGGCCTTCCTGATGCTGGTGGACCTCTGATCCGAAAGCCGGATGCATACACTGAGCAA
CTGCACAATGAGGAGGAGAATGAGGATGCTCGCTCTATGGCAGCAGCTGCTGCATCCCTCGG
GGGACCCCGAGCAAACACAGTCCTGGAGCGTGTGGAAGGTGCCCAGCAGCGGTGGAAGCTGC
AGGTGCAGGAGCAGCGGAAGACAGTCTTCGATCGGCACAAGATGCTCAGC**TAGAT**GGGCTGG
TGTGGTTGGGTCAAGGCCCAACACCATGGCTGCCAGCTTCCAGGCTGGACAAAGCAGGGGG
CTACTTCTCCCTTCCCTCGGTTCCAGTCTTCCCTTTAAAGCCTGTGGCATTTTTCTCCTT
CTCCCTAACTTTAGAAATGTTGTACTTGGCTATTTTGATTAGGGAAGAGGGATGTGGTCTCT
GATCTCTGTTGTCTTCTTGGGTCTTTGGGGTTGAAGGGAGGGGGAAGGCAGGCCAGAAGGGA
ATGGAGACATTCGAGGCGGCCTCAGGAGTGGATGCGATCTGTCTCTCCTGGCTCCACTCTTG
CCGCCTTCCAGCTCTGAGTCTTGGGAATGTTGTTACCTTTGGAAGATAAAGCTGGGTCTTCA
GGAACTCAGTGTCTGGGAGGAAAGCATGGCCCAGCATTCAGCATGTGTTCCCTTTCTGCAGTG
GTTCTTATCACCACTCCCTCCCAGCCCCGGCGCCTCAGCCCCAGCCCCAGCTCCAGCCCTG
AGGACAGCTCTGATGGGAGAGCTGGGCCCCCTGAGCCCACTGGGTCTTCAGGGTGCCTGGA
AGCTGGTGTTCGCTGTCCCCTGTGCACTTCTCGCACTGGGGCATGGAGTGCCCATGCATACT
CTGCTGCCGGTCCCCTCACCTGCACTTGAGGGGTCTGGGCAGTCCCTCCTCTCCCCAGTGTC
CACAGTCACTGAGCCAGACGGTCGGTTGGAACATGAGACTCGAGGCTGAGCGTGGATCTGAA
CACCACAGCCCCTGTACTTGGGTGCCTCTTGTCCTGAACTTCGTTGTACCAGTGCATGGA
GAGAAAATTTTGTCTCTTGTCTTAGAGTTGTGTGTAAATCAAGGAAGCCATCATTAATTG
TTTTATTTCTCTCA

FIGURE 40

```
</usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA60278
<subunit 1 of 1, 183 aa, 1 stop
<MW: 20574, pI: 6.60, NX(S/T): 3
MKLLSLVAVVGCLLVPPAEANKSSEDIRCKCICPPYRNISGHIYNQNVSQKDCNCLHVVEPM
PVPGHDVEAYCLLCECRYEERSTTTIKVIIIVYLSVVGALLLYMAFLMLVDPLIRKPDAYTE
QLHNEEENEDARSMAAAAASLGGPRANTVLERVEGAQQRWKLQVQEQRKTVFDRHKMLS
```

Important features:**Signal peptide:**

amino acids 1-20

Transmembrane domain:

amino acids 90-112

N-glycosylation sites.

amino acids 21-24, 38-41 and 47-50

1000 800 600 400 200 0

FIGURE 41

AGCGGGTCTCGCTTGGGTTCCGCTAATTTCTGTCTTGAGGCGTGAGACTGAGTTCATAGGGTCTGGGTCCCCGA
 ACCAGGAAGGGTTGAGGGAACACAATCTGCAAGCCCCGCGACCCAAGTGAGGGGCCCCGTGTTGGGGTCTCC
 TCCCTTTGCATTCCCACCCCTCCGGGCTTTGCGTCTTCTGGGGACCCCTCGCCGGGAGATGCGCCGCGTTGATG
 CGGAGCAAGGATTCGTCTGCTGCTGCTCCTACTGGCCGCGGTGCTGATGGTGGAGAGCTCACAGATCGGCAGT
 TCGCGGGCCAACTCAACTCCATCAAGTCTCTCTGGGCGGGGAGACGCCTGGTCAGGCCGCCAATCGATCTGCG
 GGCATGTACCAAGGACTGGCATTCGGCGGCAGTAAGAAGGGCAAAAACCTGGGGCAGGCCTACCCTTGTAGCAGT
 GATAAGGAGTGTGAAGTTGGGAGGTATTGCCACAGTCCCCACCAAGGATCATCGGCCTGCATGGTGTGTCGGAGA
 AAAAAAGAGCGCTGCCACCGAGATGGCATGTGCTGCCCCAGTACCCGCTGCAATAATGGCATCTGTATCCCAGTT
 ACTGAAAGCATCTTAACCCCTCACATCCCGGCTCTGGATGGTACTCGGCACAGAGATCGAAACCACGGTCATTAC
 TCAAACCATGACTTGGGATGGCAGAATCTAGGAAGACCACACACTAAGATGTCACATATAAAGGGCATGAAGGA
 GACCCCTGCCTACGATCATCAGACTGCATTGAAGGGTTTTGCTGTGCTCGTCATTTCTGGACCAAAATCTGCAAA
 CCAGTGTCTCCATCAGGGGGAAGTCTGTACCAACAACGCAAGAAGGGTTCTCATGGGCTGGAAATTTTCCAGCGT
 TGGCATGTGCGAAGGGCCTGTCTTGCAAAGTATGGAAAGATGCCACCTACTCCTCCAAAGCCAGACTCCATGTG
 TGTGAGAAAATTTGATCACCATTGAGGAACATCATCAATTGCAGACTGTGAAGTTGTGATTTAATGCATTATAG
 CATGGTGGAAAATAAGGTTGAGATGCAGAAGAATGGCTAAAATAAGAAACGTGATAAGAATATAGATGATCACAA
 AAAGGGGAGAAAGAAAACATGAAGTGAATAGATTAGAATGGGTGACAAATGCAGTGCAGCCAGTGTTCATTATG
 CAACTTGTCTATGTAAATAATGTACACATTTGTGGAAAATGCTATTATTAAGAGAACAGCACACAGTGGAAATT
 ACTGATGAGTAGCATGTGACTTTTCCAAGAGTTTAGGTTGTGCTGGAGGAGAGGTTTCTTCAGATTGCTGATTGC
 TTATACAAATAACCTACATGCCAGATTTCTATTCAACGTTAGAGTTTAAACAAAATACTCCTAGAATAACTTGTTA
 TACAATAGGTTCTAAAATAAAAATTGCTAAACAAGAAATGAAAACATGGAGCATTTGTTAATTTACAACAGAAAAT
 TACCTTTTGAATTTGTAACACTACTTCTGCTGTTCAATCAAGAGTCTTGGTAGATAAGAAAAAAATCAGTCAATAT
 TTCAAATAAATTGCAAAATAATGGCCAGTTGTTTAGGAAGGCCTTTAGGAAGACAAATAAATAACAAACAAACAG
 CCACAAATACTTTTTTTTTTCAAATTTTGTAGTTTACCTGTAATTAATAAGAACTGATACAAGACAAAAACAGTTCC
 TTCAGATTCTACGGAATGACAGTATATCTCTCTTATCCTATGTGATTCTGCTCTGAATGCATTATATTTTCCA
 AACTATACCCATAAATTGTGACTAGTAAAATACTTACACAGAGCAGAAATTTTACAGATGGCAAAAAATTTAA
 GATGTCCAATATATGTGGGAAAAGAGCTAACAGAGAGATCATTATTTCTTAAAGATTGGCCATAACCTATATTTT
 GATAGAATTAGATTGGTAAATACATGTATTACATACACTCTGTGGTAATAGAGACTTAAGCTGGATCTGTACTG
 CACTGGAGTAAGCAAGAAAATTGGGAAAACTTTTTCGTTTGTTCAGGTTTGGCAACACATAGATCATATGTCTG
 AGGCACAAGTTGGCTGTTTCATCTTTGAAACCAGGGGATGCACAGTCTAAATGAATATCTGCATGGGATTTGCTAT
 CATAATATTTACTATGCAGATGAATTCAAGTGTGAGGTCCTGTGTCCGTACTATCCTCAAATTTATTTATTTATAG
 TGCTGAGATCCTCAAATAATCTCAATTTAGGAGGTTTCACAAAATGTACTCCTGAAGTAGACAGAGTAGTGAGG
 TTTTATTGCCCTCTATAAGCTTCTGACTAGCCAATGGCATCATCCAATTTTCTTCCCAACCTCTGCAGCATCTG
 CTTTATTGCCAAAGGGCTAGTTTTCGGTTTTCTGCAGCCATTGCGGTAAAAAATATAAGTAGGATAACTTGTAAA
 ACCTGCATATTGCTAATCTATAGACACCACAGTTTCTAAATTTCTTGAACCACTTTACTACTTTTTTTAACTT
 AACTCAGTTCTAAATACTTTGTCTGGAGCACAAAACAATAAAGGTTATCTTATAGTCGTGACTTTAACTTTTG
 TAGACCACAATTCACTTTTTAGTTTTCTTTACTTAAATCCCATCTGCAGTCTCAAATTTAAGTTCTCCCAGTAG
 AGATTGAGTTTGAGCCTGTATATCTATTAATAAATTCAACTTCCACATATATTTACTAAGATGATTAAGACTTA
 CATTTTCTGCACAGGTCTGCAAAAACAAAATTATAAACTAGTCCATCCAAGAACCAGTTTGTATAAACAGGT
 TGCTATAAGCTTGTGAAATGAAAATGGAACATTTCAATCAAACATTTCCATATATAACAATTATTTATATTTACAAT
 TTGGTTTTCTGCAATATTTTTCTTATGTCCACCCTTTTAAAAATTATTTTGAAGTAATTTATTTACAGGAAATG
 TTAATGAGATGATTTTTCTTATAGAGATATTTCTTACAGAAAGCTTTGTAGCAGAATATATTTGCAGCTATTGAC
 TTTGTAATTTAGGAAAAATGTATAATAAGATAAAATCTATTAAATTTTTCTCCTCTAAAAACTGAAAAA
 AAAAAAAAAAAAAAAAAA

FIGURE 42

MAALMRSKDSSCCLLLLLAAVLMVESSQIGSSRAKLNSIKSSLGGETPGQAANRSAGMYQGLA
FGGSKKGKNLQAYPCSSDKECEVGRYCHSPHQSSACMVCRRKKKRCHRDGMCCPSTRCNN
GICIPVTESILTPHIPALDGTRHRDRNHGHYSNHDLGWQNLGRPHTKMSHIKGHEGDPCLRS
SDCIEGFCCARHFWTKICKPVLHQGEVCTKQRKKGSHGLEIFQRCDCAKGLSCKVWKDATYS
SKARLHVCQKI

Signal peptide:

amino acids 1-25

1000
900
800
700
600
500
400
300
200
100
0

FIGURE 43

TCTCAATCTGCTGACCTCGTGGATCCGCGCTGACCTTGTAATCCACCTACCTTGGCCCTCCCAA
GTGTTGGGATTACAGGCGTGAGCCACCGCGCCCGGCCAACATCACGTTTTTAAAAATTGATT
TCTTCAAATTCATGGCAAATATTTCCCTTCCCTTTAACTTCTTATGTGAGAATGAGGAAGGA
TAGCTGCATTTATTTAGTCAGTTTTTCATTGCATAGTAATATTTTCATGTAGTATTTTCTAAG
TTATATTTTAGTAATTCATATGTTTTAGATTATAGGTTTTAACATACTTGTGAAAATACTTG
ATGTGTTTTTAAAGCCTTGGGCAGAAATTCTGTATTGTTGAGGATTTGTTCTTTTATCCCCCT
TTTAAAGTCATCCGTCCTTGGCTCAGGATTTGGAGAGCTTGCACCACCAAAAATGGCAAACA
TCACCAGCTCCCGAGATTTTGGACCAGTTGAAAGCTCCGAGTTTGGGCCAGTTTACCACCACC
CCAAGTACACAGCAGAATAGTACAAGTCACCCTACAACACTACTACTTCTTGGGACCTCAAGCC
CCCAACATCCCAGTCCTCAGTCCTCAGTCATCTTGACTTCAAATCTCAACCTGAGCCATCCC
CAGTTCTTAGCCAGTTGAGCCAGCGACAACAGCACCAGAGCCAGGCAGTCACTGTTCTCTCT
CCTGGTTTTGGAGTCCTTTCTTCCCAGGCCAAAACCTTCGAGAATCAACACCTGGAGACAGTCC
CTCCACTGTGAACAAGCTTTTGCAGCTTCCCAGCACGACCATTGAAAATATCTCTGTGTCTG
TCCACCAGCCACAGCCCAAACACATCAAACCTTGCTAAGCGGCGGATACCCCCAGCTTCTAAG
ATCCCAGCTTCTGCAGTGGAAATGCCTGGTTCAGCAGATGTCACAGGATTAAATGTGCAGTT
TGGGGCTCTGGAATTTGGGTCAGAACCTTCTCTCTCTGAATTTGGATCAGCTCCAAGCAGTG
AAAATAGTAATCAGATTCCCATCAGCTTGTATTCGAAGTCTTTAAGTGAGCCTTTGAATACA
TCTTTATCAATGACCAGTGCAGTACAGAACTCCACATATACAACCTTCCGTCATTACCTCCTG
CAGTCTGACAAGCTCATCACTGAATTCTGCTAGTCCAGTAGCAATGTCTTCCTCTTATGACC
AGAGTTCTGTGCATAACAGGATCCCATAACAAAGCCCTGTGAGTTCATCAGAGTCAGCTCCA
GGAACCATCATGAATGGACATGGTGGTGGTGAAGTCAGCAGACACTAGACAGTAAGTATAG
CAGCAAGCTACTCTTGTGTCATGGCTGGTGCCAACCAAACAGAGGAAGAGGATAGCTCACGTGA
TGTGGAAAACACCAGTTGGTCAATGGCTCATTTCGT**TAA**AAAGCAGCCCTTTTGCTTTTTTTGT
TTTTGGACCAGGTGTTGGCTGTGGTGTATTAGAAATGTCTTAACCACAGCAAGAAGGAGGT
GGTGGTCTCATATTCCTTCTGCCCTAATCAGACTGCACCACAAGTGCAGCATAAGTATGCAT
TTTAAAGATGCTTGGGCCAGGCGGGGTGGCTGATGCCCATAAATCCCAGTGCTTTGGGGGGCC
AAGGCAGGCAGATTGCCCAAGCTCAGGAGTTTGAGACCAACCTGGGCAACATGGTGAAACTC
TGTCTCTACTAAAATACGAAAACTAGCCGGGTGTGGTGGCGGCGCGTGCCTGTAATCCAG
CTACTTGGGAGGCTGAGGCACAAGAATCGCTTGAGCCAGCTTGGGCTACAAAGTGAGACTCC
GTCTGAAAAGA

FIGURE 44

MCFKALGRNSVLLRICSFIPLLKSSVLGSGFGELAPPKMANITSSQILDQLKAPSLGQFTTT
PSTQQNSTSHPTTTTTSWDLKPPTSQSSVLSHLDFKSQPEPSPVLSQLSQRQQHQSQAVTVPP
PGLESFPSQAKLRESTPGDSPSTVNKLLQLPSTTIENISVSVHQPPKHIKLAKRRIPPASK
IPASAVEMPGSADVTGLNVQFGALEFGSEPSLSEFGSAPSSSENSNQIPISLYSKSLSEPLNT
SLSMTSAVQNSTYTTTSVITSCSLTSSSLNSASPVAMSSSYDQSSVHNRIPIYQSPVSSSESAP
GTIMNGHGGGRSQQTLDISKYSSKLLLSWLVP TKQRKRIAHVMWKTPVGQWLIR

Signal peptide:

amino acids 1-24

1000 900 800 700 600 500 400 300 200 100 0

FIGURE 45

GCCGAGTGGGACAAAGCCTGGGGCTGGGGCGGGGCC**ATG**GCGCTGCCATCCCGAATCCTGCT
 TTGGAAACTTGTGCTTCTGCAGAGCTCTGCTGTTCTCCTGCACTCAGCGGTGGAGGAGACGG
 ACGCGGGGCTGTACACCTGCAACCTGCACCATCACTACTGCCACCTCTACGAGAGCCTGGCC
 GTCCGCCTGGAGGTACCCGACGGCCCCCGGCCACCCCGCCTACTGGGACGGCGAGAAGGA
 GGTGCTGGCGGTGGCGCGGGCGCACCCGCGCTTCTGACCTGCGTGAACCGCGGGCACGTGT
 GGACCGACCGGCACGTGGAGGAGGCTCAACAGGTGGTGCCTGCGGACCGGCAGCCGCCCGGG
 GTCCCGCACGACCGCGCGGACCGCCTGCTGGACCTCTACGCGTCGGGCGAGCGCCGCGCCTA
 CGGGCCCCCTTTTTCTGCGCGACCGCGTGGCTGTGGGCGCGGATGCCTTTGAGCGCGGTGACT
 TCTCACTGCGTATCGAGCCGCTGGAGGTGCGCGACGAGGGCACCTACTCCTGCCACCTGCAC
 CACCATTACTGTGGCCTGCACGAACGCCGCGTCTTCCACCTGACGGTCGCCGAACCCACGC
 GGAGCCGCCCCCGGGGCTCTCCGGGCAACGGCTCCAGCCACAGCGGCGCCCCAGGCCCAG
 ACCCCACACTGGCGCGCGGCCACAACGTATCAATGTCATCGTCCCCGAGAGCCGAGCCAC
 TTCTTCCAGCAGCTGGGCTACGTGCTGGCCACGCTGCTGCTCTTCATCCTGCTACTGGTCAC
 TGTCTCCTGGCCGCCCGCAGGCGCCGCGGAGGCTACGAATACTCGGACCAGAAGTCGGGAA
 AGTCAAAGGGGAAGGATGTTAACTTGGCGGAGTTCGCTGTGGCTGCAGGGGACCAGATGCTT
 TACAGGAGTGAGGACATCCAGCTAGATTACAAAAACAACATCCTGAAGGAGAGGGCGGAGCT
 GGCCACAGCCCCCTGCCTGCCAAGTACATCGACCTAGACAAAGGGTTCCGGAAGGAGAACT
 GCAA**ATAG**GGAGGCCCTGGGCTCCTGGCTGGGCCAGCAGCTGCACCTCTCCTGTCTGTGCTC
 CTCGGGGCATCTCCTGATGCTCCGGGGCTCACCCCCCTTCCAGCGGCTGGTCCCGCTTTCCT
 GGAATTTGGCCTGGGCGTATGCAGAGGCCGCTCCACACCCCTCCCCAGGGGCTTGGTGGC
 AGCATAGCCCCACCCCTGCGGCCTTTGCTCACGGGTGGCCCTGCCACCCCTGGCACAACC
 AAAATCCCCTGATGCCCATCATGCCCTCAGACCCCTTCTGGGCTCTGCCCGCTGGGGGCCTG
 AAGACATTCTTGGAGGACACTCCCATCAGAACCTGGCAGCCCCAAA**ACT**GGGGTCAGCCTCA
 GGGCAGGAGTCCCCTCCTCCAGGGCTCTGCTCGTCCGGGGCTGGGAGATGTTCTTGGAGGA
 GGACACTCCCATCAGAACTTGGCAGCCTTGAAGTTGGGGTCAGCCTCGGCAGGAGTCCCCT
 CCTCCTGGGGTGTGCTGCCTGCCACCAAGAGCTCCCCACCTGTACCACCATGTGGGACTCCAG
 GCACCATCTGTTCTCCCCAGGGACCTGCTGACTTGAATGCCAGCCCTTGCTCCTCTGTGTTG
 CTTTGGGCCACCTGGGGCTGCACCCCTGCCCTTTCTCTGCCCATCCCTACCCTAGCCTTG
 CTCTCAGCCACCTTGATAGTCACTGGGCTCCCTGTGACTTCTGACCCTGACACCCCTCCCTT
 GGACTCTGCCTGGGCTGGAGTCTAGGGCTGGGGCTACATTTGGCTTCTGTACTGGCTGAGGA
 CAGGGGAGGGAGTGAAGTTGGTTTGGGGTGGCCTGTGTTGCCACTCTCAGCACCCACATTT
 GCATCTGCTGGTGGACCTGCCACCATCACAATAAAGTCCCCATCTGATTTTTTAAAAAAAAA
 AAAAA

FIGURE 46

</usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA60618

<subunit 1 of 1, 341 aa, 1 stop

<MW: 38070, pI: 6.88, NX(S/T): 1

MALPSRILLWKLVLQSSAVLLHSAVEETDAGLYTCNLHHHYCHLYESLAVRLEVTDGPPAT
PAYWDGEKEVLAVARGAPALLTCVNRGHVWTD RHVEEAQQVVHWDRQPPGVPHDRADRLDL
YASGERRAYGPLEFLRDRVAVGADAFERGFSLRIEPLVADEGTYSCHLHHHYCGLHERRVF
HLTVAEPHAEPFPRGSPGNSSSHSGAPGPDPTLARGHNVINVIVPESRAHFFQQLGYVLATL
LLFILLLVTVLLAARRRRGGYEYSDQKSGKSKGKDVNLAEFAVAAGDQMLYRSEDIQLDYKN
NILKERAELAHSPLPAKYIDLDKGFRKENCK

Important features:**Signal peptide:**

amino acids 1-19

Transmembrane domain:

amino acids 237-262

N-glycosylation site.

amino acids 205-208

Cell attachment sequence.

amino acids 151-154

Coproporphyrinogen III oxidase proteins.

amino acids 115-140

FIGURE 47

CGCCGGAGGCAGCGGCGGCGTGGCGCAGCGGCGAC**ATG**GCCGTTGTCTCAGAGGACGACTTT
CAGCACAGTTCAAACTCCACCTACGGAACCACAAGCAGCAGTCTCCGAGCTGACCAGGAGGC
ACTGCTTGAGAAGCTGCTGGACCGCCCCGCCCCCTGGCCTGCAGAGGCCCGAGGACCGCTTCT
GTGGCACATACATCATCTTCTTCAGCCTGGGCATTGGCAGTCTACTGCCATGGAACCTTCTTT
ATCACTGCCAAGGAGTACTGGATGTTCAAACCTCCGCAACTCCTCCAGCCCAGCCACCGGGGA
GGACCCTGAGGGCTCAGACATCCTGAACTACTTTGAGAGCTACCTTGCCGTTGCCTCCACCG
TGCCCTCCATGCTGTGCTGGTGGCCAACTTCTGCTTGTCAACAGGGTGCAGTCCACATC
CGTGTCTTGGCCTCACTGACGGTCATCCTGGCCATCTTCATGGTGATAACTGCACTGGTGAA
GGTGGACACTTCTCCTGGACCCGTGGTTTTTTTTGCGGTCACCATTGTCTGCATGGTGATCC
TCAGCGGTGCCTCCACTGTCTTCAGCAGCAGCATCTACGGCATGACCGGCTCCTTTCCTATG
AGGAACTCCCAAGCACTGATATCAGGAGGAGCCATGGGCGGGACGGTCAGCGCCGTGGCCTC
ATTGGTGGACTTGGCTGCATCCAGTGATGTGAGGAACAGCGCCCTGGCCTTCTTCCTGACGG
CCACCATCTTCCTCGTGCTCTGCATGGGACTCTACCTGCTGCTGTCCAGGCTGGAGTATGCC
AGGTACTACATGAGGCCTGTTCTTGCGGCCCATGTGTTTTCTGGTGAAGAGGAGCTTCCCCA
GGACTCCCTCAGTGCCCCCTTCGGTGGCCTCCAGATTCAATTGATTCCCACACACCCCTCTCC
GCCCCATCCTGAAGAAGACGGCCAGCCTGGGCTTCTGTGTACCTACGTCTTCTTCATCACC
AGCCTCATCTACCCCGCCGTCTGCACCAACATCGAGTCCCTCAACAAGGGCTCGGGCTCACT
GTGGACCACCAAGTTTTTCATCCCCCTCACTACCTTCTCCTGTACAACCTTTGCTGACCTAT
GTGGCCGGCAGCTCACCGCCTGGATCCAGGTGCCAGGGCCCAACAGCAAGGCGCTCCCAGGG
TTCGTGCTCCTCCGGACCTGCCTCATCCCCCTCTTCGTGCTCTGTAACCTACCAGCCCCGCGT
CCACCTGAAGACTGTGGTCTTCCAGTCCGATGTGTACCCCGCACTCCTCAGCTCCCTGCTGG
GGCTCAGCAACGGCTACCTCAGCACCCCTGGCCCTCCTCTACGGGCCTAAGATTGTGCCCAGG
GAGCTGGCTGAGGGCCACGGGAGTGGTGATGTCCTTTTATGTGTGCTTGGGCTTAACACTGGG
CTCAGCCTGCTCTACCCCTCCTGGTGCACCTCATCT**AGAA**GGGAGGACACAAGGACATTGGTG
CTTCAGAGCCTTTGAAGATGAGAAGAGAGTGCAGGAGGGGCTGGGGGCCATGGAGGAAAGGCC
TAAAGTTTCACTTGGGGACAGAGAGCAGAGCACACTCGGGCCTCATCCCTCCCAAGATGCCA
GTGAGCCACGTCCATGCCCATTCGCTGCAAGGCAGATATTCCAGTCATATTAACAGAACACT
CCTGAGACAGTTGAAGAAGAAATAGCACAAATCAGGGGTACTCCCTTCACAGCTGATGGTTA
ACATTCCACCTTCTTTCTAGCCCTTCAAAGATGCTGCCAGTGTTGCGCCCTAGAGTTATTACA
AAGCCAGTGCCAAAACCCAGCCATGGGCTCTTTGCAACCTCCCAGCTGCGCTCATTCAGCT
GACAGCGAGATGCAAGCAAATGCTCAGCTCTCCTTACCCTGAAGGGGTCTCCCTGGAATGGA
AGTCCCTGGCATGGTCAGTCCTCAGGCCCAAGACTCAAGTGTGCACAGACCCCTGTGTTCT
GCGGGTGAACAACTGCCCACTAACCAGACTGGAAAACCCAGAAAGATGGGCCTTCATGAAT
GCTTCATTCCAGAGGGACCAGAGGGCCTCCCTGTGCAAGGGATCAAGCATGTCTGGCCTGGG
TTTTCAAAAAAAGAGGGATCCTCATGACCTGGTGGTCTATGGCCTGGGTCAAGATGAGGGTC
TTTCAGTGTTCTGTTTACAACATGTCAAAGCCATTGGTTCAAGGGCGTAATAAATACTTGC
GTATTCAAAA

FIGURE 48

MAVVSEDDFQHSSNSTYGTSSSLRADQEALLEKLLDRPPPGLQRPEDRFCGTYIIFFSLGI
GSLLPWNFFITAKEYWMFKLRNSSSPATGEDPEGSDILNYFESYLAVASTVPSMLCLVANFL
LVNRVAVHIRVLASLTVILAIEMVITALVKVDTSWTRGFFAVTIVCMVILSGASTVFSSSI
YGMTGSFPMRNSQALISGGAMGGTVSAVASLVDLAASSDVRNSALAFFLTATIFLVLCMGLY
LLLSRLEYARYYMRPVLAHVFSGEEELPQDSLSAPSVASRFIDSHTPPLRPILKKTASLGF
CVTYVFFITSLLIYPAVCTNIESLNKSGSLWTTKFFIPLTTFLLYNFADLCGRQLTAWIQVP
GPNSKALPGFVLLRTCLIPLEFVLCNYQPRVHLKTVVFQSDVYPALLSSLLGLSNGYLSTLAL
LYGPKIVPRELAEATGVVMSFYVCLGLTLGSACSTLLVHLI

Transmembrane domain:

amino acids 50-74 (type II), 105-127, 135-153, 163-183, 228-252,
305-330, 448-472

48/249

FIGURE 49

GACAGTGGAGGGCAGTGGAGAGGACCGCGCTGTCTGCTGTCACCAAGAGCTGGAGACACCA
 TCTCCCACCGAGAGTCA**ATG**GCCCCATTGGCCCTGCACCTCCTCGTCCTCGTCCCCATCCTCC
 TCAGCCTGGTGGCCTCCCAGGACTGGAAGGCTGAACGCAGCCAAGACCCCTTCGAGAAATGC
 ATGCAGGATCCTGACTATGAGCAGCTGCTCAAGGTGGTGACCTGGGGGCTCAATCGGACCCT
 GAAGCCCCAGAGGGTGATTGTGGTTGGCGCTGGTGTGGCCGGGCTGGTGGCCGCCAAGGTGC
 TCAGCGATGCTGGACACAAGGTCACCATCCTGGAGGCAGATAACAGGATCGGGGGCCGCATC
 TTCACCTACCGGGACCAGAACACGGGCTGGATTGGGGAGCTGGGAGCCATGCGCATGCCAG
 CTCTCACAGGATCCTCCACAAGCTCTGCCAGGGCCTGGGGCTCAACCTGACCAAGTTCACCC
 AGTACGACAAGAACACGTGGACGGAGGTGCACGAAGTGAAGCTGCGCAACTATGTGGTGGAG
 AAGGTGCCCCGAGAAGCTGGGCTACGCCTTGCGTCCCCAGGAAAAGGGCCACTCGCCCGAAGA
 CATCTACCAGATGGCTCTCAACCAGGCCCTCAAAGACCTCAAGGCACTGGGCTGCAGAAAGG
 CGATGAAGAAGTTTGAAAGGCACACGCTCTTGGAATATCTTCTCGGGGAGGGGAACCTGAGC
 CGGCCGGCCGTGCAGCTTCTGGGAGACGTGATGTCCGAGGATGGCTTCTTCTATCTCAGCTT
 CGCCGAGGCCCTCCGGGCCCACAGCTGCCTCAGCGACAGACTCCAGTACAGCCGCATCGTGG
 GTGGCTGGGACCTGCTGCCGCGCGCGCTGCTGAGCTCGCTGTCCGGGCTTGTGCTGTTGAAC
 GCGCCCGTGGTGGCGATGACCCAGGGACCGCACGATGTGCACGTGCAGATCGAGACCTCTCC
 CCCGGCGCGGAATCTGAAGGTGCTGAAGGCCGACGTGGTGTGCTGACGGCGAGCGGACCGG
 CGGTGAAGCGCATCACCTTCTCGCCGCCGCTGCCCCGCCACATGCAGGAGGCGCTGCGGAGG
 CTGCACTACGTGCCGGCCACCAAGGTGTTCTTAAGCTTCCGCAGGCCCTTCTGGCGCGAGGA
 GCACATTGAAGGCGGCCACTCAAACACCGATCGCCCGTCGCGCATGATTTTCTACCCGCCGC
 CGCGCGAGGGCGCGCTGCTGCTGGCCTCGTACACGTGGTCGGACGCGGCGGCAGCGTTCCGC
 GGCTTGAGCCGGGAAGAGGCGTTGCGCTTGGCGCTCGACGACGTGGCGGCATTGCACGGGCC
 TGTCTGTGCGCCAGCTCTGGGACGGCACCGGCGTCGTCAAGCGTTGGGCGGAGGACCAGCACA
 GCCAGGGTGGCTTTGTGGTACAGCCGCCGGCGCTCTGGCAAACCGAAAAGGATGACTGGACG
 GTCCCTTATGGCCGCATCTACTTTGCCGGCGAGCACACCGCCTACCCGCACGGCTGGGTGGA
 GACGGCGGTCAAGTCGGCGCTGCGCGCCGCCATCAAGATCAACAGCCGGAAGGGGCCTGCAT
 CGGACACGGCCAGCCCCGAGGGGACGCATCTGACATGGAGGGGCAGGGGCATGTGCATGGG
 GTGGCCAGCAGCCCCTCGCATGACCTGGCAAAGGAAGAAGGCAGCCACCCTCCAGTCCAAGG
 CCAGTTATCTCTCCAAAACACGACCCACACGAGGACCTCGCAT**TAA**AGTATTTTCGGAAAAA
 AA

FIGURE 50

MAPLALHLLVLPILLSLVASQDWKAERSQDPFEKCMQDPDYEQLLKVVVTWGLNRTLKPQRV
IVVGAGVAGLVAAKVLSDAGHKVTILEADNRIGGRIFTYRDQNTGWIGELGAMRMPSSHRI
HKLCQGLGLNLTKFTQYDKNTWTEVHEVKLRNYVVEKVPEKLGYPALRPQEKGHS PEDIIYQMA
LNQALKDLKALGCRKAMKKFERHTLLEYLLGEGNLSRPAVQLLGDVMSEDGEFFYLSFAEALR
AHSCLSDRLQYSRIVGGWDL LPRALLSSLSGLVLLNAPVVAMTQGPHDVHVQIETSPPARNL
KVLKADVLLTASGPAVKRITFSPPLPRHMQEALRRLHYVPATKVFLSFRRPFWRREEHIEGG
HSNTDRPSRMIFYPPPREGALLASYTWSDAAAAFAGLSREEALRLALDDVAALHGPVVRQL
WDGTGVVKRWAEDQHSQGGFVVQPPALWQTEKDDWTVPYGRIYFAGEHTAYPHGWVETAVKS
ALRAAIKINSRKG PASDTASPEGHASDMEGQGHVHGVASSPSHDLAKEEGSHPPVQGQLSLQ
NTTHTRTSH

Signal peptide:

amino acids 1-21

bioRxiv preprint doi: <https://doi.org/10.1101/000000>; this version posted January 1, 2014. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.

FIGURE 51

CTGACATGGCCTGACTCGGGACAGCTCAGAGCAGGGCAGAACCTGGGGACACTCTGGGCCGGCCTTCTGCCTGCAT
 GGACGCTCTGAAGCCACCCTGTCTCTGGAGGAACCACGAGCGAGGGAAGAGGACAGGGACTCGTGTGGCAGGAA
 GAACTCAGAGCCGGGAAGCCCCATTCACTAGAAGCACTGAGAGATGCGGCCCTCGCAGGGTCTGAATTTCT
 GCTGCTGTTTACAAAGATGCTTTTTATCTTTAACTTTTTGTTTTCCCACTTCCGACCCCGGCGTTGATCTGCAT
 CCTGACATTTGGAGCTGCCATCTTCTTGTGGCTGATCACCAGACCTCAACCCGTCTTACCTCTTCTTGACCTGAA
 CAATCAGTCTGTGGGAATTGAGGGAGGAGCACGGAAGGGGGTTTCCAGAAGAACAATGACCTAACAAGTTGCTG
 CTCTCAGATGCCAAGACTATGTATGAGGTTTTCCAAAGAGGACTCGCTGTGTCTGACAATGGGCCCTGCTTGGG
 ATATAGAAAACCAACAGCCCTACAGATGGCTATCTTACAAACAGGTGTCTGATAGAGCAGAGTACCTGGGTTC
 CTGTCTCTTGATAAAGGTTATAAATCATCACCAGACAGTTTGTGCGCATCTTTGCTCAGAATAGGCCAGAGTG
 GATCATCTCCGAATTGGCTTGTTACACGTACTCTATGGTAGCTGTACCTCTGTATGACACCTTGGGACCAGAAGC
 CATCGTACATATTGTCAACAAGGCTGATATCGCCATGGTGATCTGTGACACACCCAAAAGGCATTGGTGCTGAT
 AGGGAATGTAGAGAAAGGCTTACCCCGAGCCTGAAGGTGATCATCCTTATGGACCCCTTTGATGATGACCTGAA
 GCAAAGAGGGGAGAAGAGTGAATTGAGATCTTATCCCTATATGATGCTGAGAACCTAGGCAAAGAGCACTTCAG
 AAAACCTGTGCCTCCTAGCCCAGAAGACCTGAGCGTCATCTGCTTACCAGTGGGACCACAGGTGACCCCAAAGG
 AGCCATGATAACCCATCAAAATATTGTTTCAAATGCTGCTGCCTTTCTCAAATGTGTGGAGCATGCTTATGAGCC
 CACTCCTGATGATGTGGCCATATCCTACCTCCCTCTGGCTCATATGTTTGAGAGGATTGTACAGGCTGTTGTGTA
 CAGCTGTGGAGCCAGAGTTGGATTCTTCCAAGGGGATATTGCGTTGCTGGCTGACGACATGAAGACTTTGAAGCC
 CACATTGTTTTCCCGCGGTGCCCTCGACTCCTTAAACAGGATCTACGATAAGGTACAAAATGAGGCCAAGACACCCCTT
 GAAGAAGTTCTTGTGAAGCTGGCTGTTTCCAGTAAATTCAAAGAGCTTCAAAAGGGTATCATCAGGCATGATAG
 TTTCTGGGACAAGCTCATCTTTGCAAAGATCCAGGACAGCCTGGGCGGAAGGGTTCGTGTAATTGTCACTGGAGC
 TGCCCCCATGTCCACTTCAGTCATGACATCTTCCGGGCAGCAATGGGATGTGAGGTGTATGAAGCTTATGGTCA
 AACAGAATGCACAGGTGGCTGTACATTTACATTACCTGGGGACTGGACATCAGGTACGTTGGGGTGGCCCTGGC
 TTGCAATTACGTGAAGCTGGAAGATGTGGCTGACATGAACCTACTTTACAGTGAATAATGAAGGAGAGGTCTGCAT
 CAAGGGTACAAACGTGTTCAAAGGATACCTGAAGGACCCCTGAGAAGACACAGGAAGCCCTGGACAGTGATGGCTG
 GCTTCACACAGGAGACATTGGTCGCTGGCTCCCGAATGGAACCTCTGAAGATCATCGACCGTAAAAAGAACATTTT
 CAAGCTGGCCCAAGGAGAATACATTGCACCAGAGAAGATAGAAAATATCTACAACAGGAGTCAACCAGTGTTACA
 AATTTTTGTACACGGGGAGAGCTTACGGTCATCCTTAGTAGGAGTGGTGGTTCTTGACACAGATGTACTTCCCTC
 ATTTGCAGCCAAGCTTGGGGTGAAGGGCTCCTTTGAGGAACTGTGCCAAAACCAAGTTGTAAGGGAAGCCATTTT
 AGAAGACTTGCAGAAAATTGGGAAAGAAAGTGGCTTAAAACTTTTGAACAGGTCAAAGCCATTTTTCTTCATCC
 AGAGCCATTTTCCATTGAAAATGGGCTCTTGACACCAACATTGAAAGCAAAGCGAGGAGAGCTTTCCAAATACTT
 TCGGACCCAAATTGACAGCCTGTATGAGCACATCCAGGATTAGGATAAGGTACTTAAGTACCTGCCGGCCCACTG
 TGCATGCTTGTGAGAAAATGGATTAAAAACTATTCTTACATTTGTTTTGCCCTTCTCCTATTTTTTTTAAACC
 TGTTAAACTCTAAAGCCATAGCTTTTGTTTTATATTGAGACATATAATGTGTAAACTTAGTTCCCAAATAAATCA
 ATCCTGTCTTTCCCATCTTCGATGTTGCTAATATTAAGGCTTCAGGGCTACTTTTATCAACATGCCTGTCTTCAA
 GATCCCACTTTTATGTTCTGTGCTCCTTCCCTCATGATTTCCAACCTTAATACTATTAGTAACCACAAGTTCAAGGGT
 CAAAGGGACCCCTCTGTGCCTTCTTCTTTGTTTTGTGATAAACATAACTTGCCAACAGTCTCTATGCTTATTTACA
 TCTTCTACTGTTCAAACCTAAGAGATTTTAAATCTGAAAAACTGCTTACAATTCATGTTTTCTAGCCACTCCAC
 AAACCACTAAAAATTTTAGTTTGTAGCCTATCACTCATGTCAATCATATCTATGAGACAAATGTCTCCGATGCTCTT
 CTGCGTAAATTAATTTGTGTACTGAAGGGAAAAAGTTTGTATACATAACCAACATTTTCTAAACTCTCTAGTTAGATA
 TCTGACTTGGGAGTATTAAAAATTTGGTCTATGACATACTGTCCAAAAGGAATGCTGTTCTTAAAGCATTATTTA
 CAGTAGGAACCTGGGGAGTAAATCTGTTCCCTACAGTTTGCTGCTGAGCTGGAAGCTGTGGGGGAAGGAGTTGACA
 GGTGGGCCCAGTGAACCTTTCCAGTAAATGAAGCAAGCACTGAATAAAAACCTCCTGAACCTGGGAACAAAGATCT
 ACAGGCAAGCAAGATGCCACACAACAGGCTTATTTTCTGTGAAGGAACCAACTGATCTCCCCACCCTTGATT
 AGAGTTCTCTGCTTACCTTACCACAGATAACACATGTTGTTTCTACTTGTAATGTAAAGTCTTTAAATAAAC
 TATTACAGATAAAAAA

FIGURE 52

</usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA60775

<subunit 1 of 1, 739 aa, 1 stop

<MW: 82263, pI: 7.55, NX(S/T): 3

MDALKPPCLWRNHERGKKDRDSCGRKNSEPGSPHSLEALRDAAPSQGLNFKLLFTKMLFIFN
FLFSPLPTPALICILTFGAAIFLWLITRPQPVLPLLDLNNQSVGIEGGARKGVSQKNNDLTS
CCFSDAKTMYEVFQRLAVSDNGPCLGYRKPNQPYRWLSYKQVSDRAEYLGSCLLHKGYKSS
PDQFVGIFAQNRPEWIISELACYTYSMVAVPLYDTLGPEAIVHIVNKADIAMVICDTPQKAL
VLIGNVEKGFTPSLKVIIILMDPFDDDLKQERGEKSGIEILSLYDAENLGKEHFRKPVPPSPED
LSVICFTSGTTGDPKGAMITHQNIVSNAAAFKLCVEHAYEPTPDDVAISYLPLAHMFERIVQ
AVVYSCGARVGFFQGDIRLLADDMKTLKPTLFPAPVRLNRIYDKVQNEAKTPLKKFLLKLA
VSSKFKELOKGIIRHDSFWDKLIFAKIQDSLGRVRVIVTGAAPMSTSVMTFFRAAMGCQVY
EAYGQTECTGGCTFTLPGDWTSGHVGVPACNYVKLEDVADMNYFTVNNEGEVCIKGTNVFK
GYLKDPEKTQEALDSGWLHTGDIGRWLPNGTLKIIDRKKNIFKLAQGEYIAPEKIENIYNR
SQPVLQIFVHGESLRSSLVGVVVPD TDVLP SFAAKLGVKGSFEELCQNQV VREAILEDLQKI
GKESGLKTFEQVKAI FLHPEPF SIENGLLTPTLKAKRGELSKYFRTQIDSLYEHIQD

Important features:**Type II transmembrane domain:**

amino acids 61-80

Putative AMP-binding domain signature.

amino acids 314-325

N-glycosylation site.

amino acids 102-105, 588-591 and 619-622

FIGURE 53

GGAGGCGGAGGCCGCGGCGAGCCGGGCCGAGCAGTGAGGGCCCTAGCGGGGGCCCGAGCGGGG
 CCCGGGGCCCCCTAAGCCATTCTGAAGTCATGGGCTGGCCAGGACATTGGTGACCCGCCAAT
 CCGGT**ATG**GACGACTGGAAGCCCAGCCCCCTCATCAAGCCCTTTGGGGCTCGGAAGAAGCGG
 AGCTGGTACCTTACCTGGAAGTATAAACTGACAAACCAGCGGGCCCTGCGGAGATTCTGTCA
 GACAGGGGCCGTGCTTTTCCTGCTGGTGAAGTGTCAATATCAAGTTGATCCTGGACA
 CTCGGCGAGCCATCAGTGAAGCCAATGAAGACCCAGAGCCAGAGCAAGACTATGATGAGGCC
 CTAGGCCGCCTGGAGCCCCACGGCGCAGAGGCAGTGGTCCCCGGCGGGTCTTGACGCTAGA
 GGTGTATTCAAGTCGCAGCAAAGTATATGTGGCAGTGGATGGCACCACGGTGTGAGGATG
 AGGCCCCGGGAGCAGGGCCGGGGCATCCATGTCAATTGTCTCAACCAGGCCACGGGCCACGTG
 ATGGCAAACGTGTGTTTGACACGTACTCACCTCATGAGGATGAGGCCATGGTGCTATTCCT
 CAACATGGTAGCGCCCGGCCGAGTGCTCATCTGCACTGTCAAGGATGAGGGCTCCTTCCACC
 TCAAGGACACAGCCAAGGCTCTGCTGAGGAGCCTGGGCAGCCAGGCTGGCCCTGCCCTGGGC
 TGGAGGGACACATGGGCCTTCGTGGGACGAAAAGGAGGTCTGTCTTCGGGGAGAAACATTC
 TAAGTCACCTGCCCTCTCTTCCTGGGGGACCCAGTCTGTGAAGACAGATGTGCCATTGA
 GCTCAGCAGAAGAGGCAGAGTGCCACTGGGCAGACACAGAGCTGAACCGTCGCCGCCGGCGC
 TTCTGCAGCAAAGTTGAGGGCTATGGAAGTGTATGCAGCTGCAAGGACCCACACCCATCGA
 GTTCAGCCCTGACCCACTCCCAGACAACAAGGTCCTCAATGTGCCTGTGGCTGTCAATTGCAG
 GGAACCGACCCAATTACCTGTACAGGATGCTGCGCTCTCTGCTTTCAGCCCAGGGGGTGTCT
 CCTCAGATGATAACAGTTTTTCATTGACGGCTACTATGAGGAACCCATGGATGTGGTGGCACT
 GTTTGGTCTGAGGGGCATCCAGCATACTCCCATCAGCATCAAGAAATGCCCGCGTGTCTCAGC
 ACTACAAGGCCAGCCTCACTGCCACTTTCAACCTGTTTCCGGAGGCCAAGTTTGCTGTGGTT
 CTGGAAGAGGACCTGGACATTGCTGTGGATTTTTTCAGTTTCCTGAGCCAATCCATCCACCT
 ACTGGAGGAGGATGACAGCCTGTACTGCATCTCTGCCTGGAATGACCAGGGGTATGAACACA
 CGGCTGAGGACCCAGCACTACTGTACCGTGTGGAGACCATGCCTGGGCTGGGCTGGGTGCTC
 AGGAGGTCTTGTACAAGGAGGAGCTTGAGCCCAAGTGGCCTACACCGGAAAAGCTCTGGGA
 TTGGGACATGTGGATGCGGATGCCTGAACAACGCCGGGGCCGAGAGTGATCATCCCTGACG
 TTTCCCGATCCTACCACTTTGGCATCGTCGGCCTCAACATGAATGGCTACTTTCACGAGGCC
 TACTTCAAGAAGCACAAGTTCAACACGGTTCCAGGTGTCCAGCTCAGGAATGTGGACAGTCT
 GAAGAAAGAAGCTTATGAAGTGAAGTTCACAGGCTGCTCAGTGAGGCTGAGGTTCTGGACC
 ACAGCAAGAACCCTTGTGAAGACTCTTTCCTGCCAGACACAGAGGGCCACACCTACGTGGCC
 TTTATTCGAATGGAGAAAGATGATGACTTCACCACCTGGACCCAGCTTGCCAAGTGCCTCCA
 TATCTGGGACCTGGATGTGCGTGGCAACCATCGGGGCCTGTGGAGATTGTTTCGGAAGAAGA
 ACCACTTCCTGGTGGTGGGGGTCCCCGGCTTCCCCCTACTCAGTGAAGAAGCCACCCTCAGTC
 ACCCAATTTTCCTGGAGCCACCCCCAAAGGAGGAGGGAGCCCCAGGAGCCCCAGAACAGAC
ATGAGACCTCCTCCAGGACCCTGCGGGGCTGGGTACTGTGTACCCCCAGGCTGGCTAGCCCT
 TCCCTCCATCCTGTAGGATTTTGTAGATGCTGGTAGGGGCTGGGGCTACCTTGTTTTTAACA
 TGAGACTTAATTACTAACTCCAAGGGGAGGGTTCCCCTGCTCCAACACCCCGTTCCCTGAGTT
 AAAAGTCTATTTATTTACTTCTTGTGGAGAAGGGCAGGAGTACCTGGGAATCATTACG
 ATCCCTAGCAGCTCATCTGCCCTTTGAATACCCTCACTTTCCAGGCCTGGCTCAGAATCTA
 ACCTATTTATTGACTGTCCTGAGGGCCTTGAAAACAGGCCGAACCTGGAGGGCCTGGATTTT
 TTTTGGGCTGGAATGCTGCCCTGAGGGTGGGGCTGGCTCTTACTCAGGAACTGCTGTGCC
 CAACCCATGGACAGGCCAGCTGGGGCCACATGCTGACACAGACTCACTCAGAGACCCTTA
 GACACTGGACCAGGCCTCCTCTCAGCCTTCTCTTGTCCAGATTTCCAAAGCTGGATAAGTT
 GGTCAATTGATTAAAAAAGGAGAAGCCCTCTGGGAAAAAAAAAAAAAAAAAAAAAAAAA

FIGURE 54

></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA61185

><subunit 1 of 1, 660 aa, 1 stop

><MW: 75220, pI: 6.76, NX(S/T): 0

MDDWKPSPLIKPFGARKKRSWYLTWKYKLTNQALRRFCQTGAVLFLLVTVIVNIKLILDTR
 RAISEANEDPEPEQDYDEALGRLEPPRRRGSGPRRVLDVEVYSSRSKVYVAVDGTTVLEDEA
 REQGRGIHVIVLNQATGHVMAKRVDFTYSPHEDEAMVLFNLMVAPGRVLICTVKDEGSFHLK
 DTAKALLRSLGSQAGPALGWRDTWAFVGRKGGPVFGEKHSKSPALSSWGDPVLLKTDVPLSS
 AEEAECHWADTELNRNRRRRFCFSKVEGYGSVCCKDPTPIEFSPDPLPDNKVLNVPVAVIAGN
 RPNYLYRMLRSLLSAQGVSPQMITVFIDGYEPMDDVVALFGLRGIQHTPISIKNARVSQHY
 KASLTATFNLFPEAKFAVVLEEDLDIAVDFFSFLSQSIHLLEEDDSLYCISAWNDQGYEHTA
 EDPALLYRVETMPGLGWVLRSLYKEELEPKWPTPEKLWDWDMWMMRMPEQRRGRECIIPDVS
 RSYHFGIVGLNMNGYFHEAYFKKHKFNTVPGVQLRNVDLKLKEAYEVEVHRLLSAEVLDHS
 KNPCEDSFLPDTEGHTYVAFIRMEKDDDDFTTWTQLAKCLHIWDLDVIRGNHRGLWRLFRKKNH
 FLVVGVPASPYSVKKPPSVTPIFLEPPPKEEGAPGAPEQT

Important features of the protein:

Transmembrane domain:

amino acids 38-55

Homologous region to Mouse GNT1

amino acids 229-660

FIGURE 55

CGGACGCGTGGGCTGCTGGTGGGAAGGCCTAAAGAACTGGAAAGCCCACTCTCTTGGAACCACCACAC
 CTGTTTAAAGAACCTAAGCACCATTAAAGCCACTGGAAATTTGTTGTCTAGTGGTTGTGGGTGAATA
 AAGGAGGGCAGAAATGGATGATTTTCATCTCCATTAGCCTGCTGTCTCTGGCTATGTTGGTGGGATGTTA
 CGTGGCCGGAATCATTCCCTTGGCTGTTAATTTCTCAGAGGAACGACTGAAGCTGGTGACTGTTTTGG
 GTGCTGGCCTTCTCTGTGGAACCTGCTCTGGCAGTCATCGTGCCTGAAGGAGTACATGCCCTTTATGAA
 GATATTCTTGAGGGAAAACACCACCAAGCAAGTGAAACACATAATGTGATTGCATCAGACAAAGCAGC
 AGAAAAATCAGTTGTCCATGAACATGAGCACAGCCACGACCACACAGCTGCATGCCCTATATTGGTG
 TTTCCCTCGTTCTGGGCTTCGTTTTTCATGTTGCTGGTGGACCAGATTGGTAACTCCCATGTGCATTCT
 ACTGACGATCCAGAAGCAGCAAGGTCTAGCAATTCCAAAATCACCACCACGCTGGGTCTGGTTGTCCA
 TGCTGCAGCTGATGGTGTTGCTTTGGGAGCAGCAGCATCTACTTCACAGACCAGTGTCCAGTTAATTG
 TGTTTGTGGCAATCATGCTACATAAGGCACCAGCTGCTTTTGGACTGGTTTCCTTCTTGATGCATGCT
 GGCTTAGAGCGGAATCGAATCAGAAAGCACTTGCTGGTCTTTGCATTGGCAGCACCAGTTATGTCCAT
 GGTGACATACTTAGGACTGAGTAAGAGCAGTAAAGAAGCCCTTTAGAGGTGAACGCCACGGGAGTGG
 CCATGCTTTTTCTCTGCCGGGACATTTCTTTATGTTGCCACAGTACATGTCTCCCTGAGGTGGGCGGA
 ATAGGGCACAGCCACAAGCCCGATGCCACGGGAGGGAGAGGCCTCAGCCGCTGGAAGTGGCAGCCCT
 GGTTCCTGGGTTCCTCATCCCTCTCATCCTGTCAGTAGGACACCAGCATTAATGTTCAAGGTCCAGC
 CTTGGTCCAGGGCCGTTTGCCATCCAGTGAGAACAGCCGGCACGTGACAGCTACTCACTTCCTCAGTC
 TCTTGTCTCACCTTGCGCATCTCTACATGTATTCCCTAGAGTCCAGAGGGGAGGTGAGGTTAAAACTG
 AGTAATGGAAAAGCTTTTAGAGTAGAAACACATTTACGTGTCAGTTAGCTATAGACATCCCATTGTGT
 TATCTTTTAAAAGGCCCTTGACATTTTTCGTTTTAATATTTCTCTTAACCCTATTCTCAGGGAAGATG
 GAATTTAGTTTTAAGGAAAAGAGGAGAACCTCATACTACAATGAAATAGTGATTATGAAAATACAGT
 GTTCTGTAATTAAGCTATGTCTCTTTCTTTAGTTTAGAGGCTCTGCTACTTTATCCATTGATTTTT
 AACATGGTTCACCACATGTAAGACTGGTGCTTTAGCATCTATGCCACATGCGTTGATGGAAGGTCATA
 GCACCCACTCACTTAGATGCTAAAGGTGATTCTAGTTAATCTGGGATTAGGGTCAGGAAAATGATAGC
 AAGACACATTGAAAGCTCTCTTTATACTCAAAGAGATATCCATTGAAAAGGGATGTCTAGAGGGATT
 TAAACAGCTCCTTTGGCAGCTGCCTCTCTGAATCCAGCCTGCCATTCCATCAAATGGAGCAGGAGAGG
 TGGGAGGAGCTTCTAAAGAGGTGACTGGTATTTTGTAGCATTCCTTGTCAAGTTCTCCTTTGCAGAA
 ACCTGTCTCCACATTCCTAGAGAGGAGCCAAGTTCTAGTAGTTTCAGTTCTAGGCTTTTCTTCAAGAA
 CAGTCAGATCACAAAGTGCTTTTGAAATTAAGGGATATTAAATTTTAAAGTGATTTTTTGGATGGTTAT
 TGATATCTTTGTAGTAGCTTTTTTTTAAAGACTACCAAATGTATGGTTGTCCCTTTTTTTTTGTTTT
 TTTTTTTTTAATTATTTCTCTTAGCAGATCAGCAATCCCTCTAGGGACCTAAATACTAGGTCAGCTTT
 GCGCAGACTGTGCTTCTCACATAACCACCTGTAGCAAGATGGATCATAAATGAGAAGTGTTTGCCTA
 TTGATTTAAAGCTTATTGGAATCATGTCTCTTGTCTCTTCGTCTTTCTTTGCTTTTCTTCTAACTTT
 TCCCTCTAGCCTCTCCTCGCCACAATTTGCTGCTTACTGCTGGTGTAAATATTTGTGTGGGATGAATT
 CTTATCAGGACAACCACTTCTCGAAGTGAATAATGAAGATAATAATATCTTTATTTATTTATCCCTT
 CAAAGAAATTACCTTTGTGTCAAATGCCGCTTTGTTGAGCCCTTAAATAACCACCTCCTCATGTGTAA
 ATTGACACAATCACTAATCTGGTAATTTAAACAATTGAGATAGCAAAAGTGTTTAAACAGACTAGGATA
 ATTTTTTTTTTTCATATTTGCCAAAATTTTTGTAAACCCTGTCTTGTCAAATAAGTGATAATATTGTAT
 TATTAATTTATTTTTACTTTCTATACCATTTCAAACACATTACACTAAGGGGGAACCAAGACTAGTT
 TCTTCAGGGCAGTGGACGTAGTAGTTTGTAAAAACGTTTTCTATGACGCATAAGCTAGCATGCCTATG
 ATTTATTTCTTCATGAATTTGTCACTGGATCAGCAGCTGTGGAAATAAAGCTTGTGAGCCCTCTGCT
 GGCCACAGTGAGGAAAGTAGCACAAATAGGATACAGTTGTATGTAGTCATTGGCAACAATTGCATACA
 ATTTTACTACCAAGAGAAGGTATAGTATGGAAAGTCCAAATGACTTCCTTGATTGGATGTTAACAGCT
 GACTGGTGTGAGACTTGAGGTTTTCATCTAGTCCTTCAAACCTATATGGTTGCCTAGATTCTCTCTGGA
 AACTGACTTTGTCAAATAAATAGCAGATTGTAGTGTCAAAAAA

FIGURE 56

MDDFISISLLSLAMLVGCYVAGIIPLAVNFSEERLKLVTVLGAGLLCGTALAVIVPEGVHAL
YEDILEGKHHQASETHNVIASDKAAEKSVVHEHEHSHDHTQLHAYIGVSLVLGFVFMLLVDQ
IGNSHVHSTDDPEAARSSNSKITTTTLGLVVHAAADGVALGAAASTSQTSVQLIVFVAIMLHK
APAAFGLVSFLMHAGLERNRIRKHLV FALAAPVMSMVTYLGLSKSSKEALSEVNATGVAML
FSAGTFLYVATVHVLPEVGGIGHSHKPDATGGRGLSRLEVAALVLGCLIPILISVGHQH

Signal peptide:

amino acids 1-18

Transmembrane domain:

amino acids 37-56, 106-122, 211-230, 240-260, 288-304



FIGURE 57


GCTCGAGGCCGGCGGCGGGAGAGCGACCCGGGCGGCCTCGTAGCGGGGCCCCGGATCCC
 CGAGTGGCGGCCGGAGCCTCGAAAAGAGATTCTCAGCGCTGATTTTGAGATGATGGGCTTGG
 GAAACGGGCGTCGCAGCATGAAGTCGCCGCCCTCGTGCTGGCCGCCCTGGTGGCCTGCATC
 ATCGTCTTGGGCTTCAACTACTGGATTGCGAGCTCCCGGAGCGTGACCTCCAGACACGGAT
 CATGGAGCTGGAAGGCAGGGTCCGCAGGGCGGCTGCAGAGAGAGGCGCCGTGGAGCTGAAGA
 AGAACGAGTTCCAGGGAGAGCTGGAGAAGCAGCGGGAGCAGCTTGACAAAATCCAGTCCAGC
 CACAACCTCCAGCTGGAGAGCGTCAACAAGCTGTACCAGGACGAAAAGGCGGTTTTGGTGAA
 TAACATCACCACAGGTGAGAGGCTCATCCGAGTGCTGCAAGACCAGTTAAAGACCCTGCAGA
 GGAATTACGGCAGGCTGCAGCAGGATGTCCTCCAGTTTCAGAAGAACCAGACCAACCTGGAG
 AGGAAGTTCTCCTACGACCTGAGCCAGTGCATCAATCAGATGAAGGAGGTGAAGGAACAGTG
 TGAGGAGCGAATAGAAGAGGTCACCAAAAAGGGGAATGAAGCTGTAGCTTCCAGAGACCTGA
 GTGAAAACAACGACCAGAGACAGCAGCTCCAAGCCCTCAGTGAGCCTCAGCCCAGGCTGCAG
 GCAGCAGGCCTGCCACACACAGAGGTGCCACAAGGGAAGGGAAACGTGCTTGGTAACAGCAA
 GTCCCAGACACCAGCCCCCAGTTCCGAAGTGGTTTTGGATTCAAAGAGACAAGTTGAGAAAG
 AGGAAACCAATGAGATCCAGGTGGTGAATGAGGAGCCTCAGAGGGACAGGCTGCCGCAGGAG
 CCAGGCCGGGAGCAGGTGGTGAAGACAGACCTGTAGGTGGAAGAGGCTTCGGGGGAGCCGG
 AGAACTGGGCCAGACCCACAGGTGCAGGCTGCCCTGTCAGTGAGCCAGGAAAATCCAGAGA
 TGGAGGGCCCTGAGCGAGACCAGCTTGTCATCCCCGACGGACAGGAGGAGGAGCAGGAAGCT
 GCCGGGGAAGGGAGAAAACCAGCAGAACTGAGAGGAGAAGATGACTACAACATGGATGAAAA
 TGAAGCAGAATCTGAGACAGACAAGCAAGCAGCCCTGGCAGGGAATGACAGAAACATAGATG
 TTTTAAATGTTGAAGATCAGAAAAGAGACACCATAAATTTACTTGATCAGCGTGAAAAGCGG
 AATCATACACTCTGAATTGAACTGGAATCACATATTTACAACAGGGCCGAAGAGATGACTA
 TAAAATGTTTCATGAGGGACTGAATACTGAAAACGTGAAATGTACTAAATAAAATGTACATCTGA

FIGURE 58

MMGLGNRRSMKSPPLVLAALVACIIVLGFNYWIIASSRSVDLQTRIMELEGRVRRAAAERGA
VELKKNEFQGELEKQREQLDKIQSSHNFQLESVNKLYQDEKAVLVNNITTGERLIRVLQDQL
KTLQRNYGRLQQDVLQFQKNQTNLERKFSYDLSQCINQMKEVKEQCEERIEEVTKKGNEAVA
SRDLSENNDQRQQQLQALSEPQPRLQAAGLPHTTEVPQGKGNVLGNSKSQTPAPSSEVVLDSCR
QVEKEETNEIQVVNEEPQRDRLPQEPGREQVVEDRPVGGRGFGGAGELGQTPQVQAALSVSQ
ENPEMEGPERDQLVIPDGQEEEQEAAGEGRNQKLRGEDDYNMDENEAESETDKQAALAGND
RNIDVFNVEDQKRDTINLLDQREKRNHTL

Signal peptide:

amino acids 1-29



GGATGTCAGAAAGCCTCAGTGTCTCTCTCTGGCTGGGTCTGCTTCTCTTACGCTGGCATTGCCCTCTTCA
CCAGTGGCTTCTGCTCACCCTGTTTGGAGCTACCAACCATAGCAGCTGCCAAGAGCCCCAGGCCCTGGGTCCC
TGCCATGGGGGAGCCAAGGGAAACCTGGGGCCTGCTGGATGGCTTCCCGATTTTCGCGGGTTGTGTTGGTGTCTGA
TAGATGCTCTGCGATTTGACTTCGCCAGCCCCAGCATTACACAGTGCCTAGAGAGCCTCCTGTCTCCCTACCCT
TCTGGGCAAACCTAAGCTCCTTGCAGAGGATCCTGGAGATTACAGCCCCACCATGCCCGGCTCTACCGATCTCAGG
TTGACCCTCCTACCACCACCATGCAGCGCCTCAAGGCCCTCACCCTGGCTACTGCTACCTTATTATGATGCTG
GTAGTAACTTCGCCAGCCACGCCATAGTGAAGACAATCTCATTAAGCAGCTCACCAGTGCAGGAAGGCGTGTAG
TCTTCATGGGAGATGATACCTGGAAAGACCTTTTCCCTGGTGTCTTCTCCAAAGCTTCTTCTTCCCATCCTTCA
ATGTCTCAGAGACCTAGACACAGTGGACAATGGCATCCTGGAACACCTCTACCCCACCATGGACAGTGGTGAATGGG
ACGTGCTGATTGCTCACTTCTCTGGGTGTGGACCACTGTGGCCACAAGCATGGCCCTCACCACCCCTGAAATGGCCA
AGAACTTAGCCAGATGGACAGGTGATCCAGGACTGTGGAGCTGTGGAGCGTCTGGAGAATGACACACTGCTGGTAGTGG
CTGGGGACCATGGGATGACCACAAATGGAGACCATGGAGGGGACAGTGAGCTGGAGGTCTCAGCTGCTCTCTTTCT
TGTATAGCCCCACAGCAGTCTTCCCCAGCACCCCCACCAGAGGAGCCAGAGGTGATTCTCAAGTTAGCCTTGTGC
CCACGCTGGCCCTGCTGCTGGGCCTGCCCATCCATTTGGGAATATCGGGGAAGTGATGGCTGAGCTATTCTCAG
GGGGTGAGGACTCCCAGCCCCACTCCTCTGCTTTAGCCCAAGCCTCAGCTCTCCATCTCAATGCTCAGCAGGTGT
CCCGATTTCTTCATACCTACTCAGCTGCTACTCAGGACCTTCAAGCTAAGGAGCTTCATCAGCTGCAGAACCTCT
TCTCAAGGCCCTCTGCTGACTACAGTGGCTTCTCCAGAGCCCCAAGGGGGCTGAGGCGACACTGCCGACTGTGA
TTGCTGAGCTGCAGAGTTCTCGGGGGAGCTCGGGCCATGTGCATCGATCTGTTGGCTCGTTTCTCTCTGGTCC
GCATGGCGGGGGGTACTGCTCTCTGGCTGCTTCTGCTTTATCTGCCTGCTGGCATCTCAGTGGGCAATATCCC
CAGGCTTTCCATTTCTGCCCTCTACTCCTGACACCTGTGGCCTGGGGCCTGGTTGGGGCCATAGCGTATGCTGGAC
TCCTGGGAACCTATTGAGCTGAAGCTAGATCTAGTGCTTCTAGGGGCTGTGGCTGCAGTGAGCTCATTCCTCCCTT
TTCTGTGGAAGCCTGGGCTGGCTGGGGTCCAAAGAGGCCCTGGCAACCCTGTTTCCCATCCCTGGGCCCGTCC
TGTTACTGCTGCTGTTTTCGTTGGCTGTGTTCTTCTCTGATAGATTTTGTGTAGCTGAGGCGAGGGCCACCCCT
TCCTTTTGGGCTCATTTCATCCTGCTCCTGGTTGTCCAGTTCTCACTGGGAGGGCCAGCTGCTCCACCTAAGCTAC
TCACAATGCCCCGCCCTTGGCACTTCAGCCACAACAAACCCCCACGGCAACAATGGTGATATGCCCTGAGGCTTG
GAATTGGGTTGCTTTTATGTACAAGGCTAGCTGGGCTTTTTCATCGTTGCCCTGAAGAGACACCTGTTTGCCACT
CCTCTCCCTGGCTGAGTCTCTGGCATCCATGGTGGGTGGTGCAGCCAAGAATTTATGGTATGGAGCTTGTGTGG
CGGCGCTGGTGGCCCTGTTAGCTGCCGTGCGCTTGTGGCTTCGCCGCTATGGTAATCTCAAGAGCCCCGAGCCAC
CCATGCTCTTTGTGCGCTGGGACTGCCCTAATGGCATTTGGTACTGCTGCTACTGGGCATTTGGCGTCCGGGG
CAGATGAGGCTCCCCCGCTCTCCGGGCTCTGGTCTCTGGGGCATTCATGGTGCTGCTCCTGGGTGTAGCAGGGC
TGGCTGCTTCAGGGCTCGCGCTGCTGCTCTGGAAGCCTGTGACAGTGCTGGTGAAGGCTGGGGCAGGCGCTCCAA
GGACCAGGACTGTCTCTCACTCCCTTCTCAGGCCCCCCCCACTTCTCAAGCTGACTTGGATTATGTGGTCCCTCAAA
TCTACCGACACATGCAGGAGGAGTTCGGGGGCCGGTTAGAGAGGACCAAATCTCAGGGTCCCCTGACTGTGGCTG
CTTATCAGTTGGGGAGTGTCTACTCAGCTGCTATGGTCAAGCCCTCACCCTGTTGGCCTTCCCACCTTCTGCTGT
TGATCGGGAGCGCATCAGCTTGTGTTCTGTTCTGTTTCTGAGAGCTTCTTCTCTACATCTGCTTGTCTG
CTGGGATAACCGCTCACCACCCCTGGTCTTTTACTGTGCCATGGCAGGCAGTCTCGGCTTGGGCCCTCATGGCCA
CACAGACCTTCTACTCCACAGGCCACCAGCCTGTCTTTCCAGCCATCCATTGGCATGCAGCCTTCGTGGGATTCC
CAGAGGGTTCATGGCTCCTGTACTTGGCTGCCTGCTTTGCTAGTGGGAGCCAACACCTTTGCCTCCCACCTCCTCT
TTGCAGTAGGTTGCCCACTGCTCCTGCTCTGGCCTTTCCTGTGTGAGAGTCAAGGGCTGCGGAAGAGACAGCAGC
CCCCAGGGAATGAAGCTGATGCCAGAGTCAGACCCGAGGAGGAAGAGGAGCCACTGATGGAGATGCGGCTCCGGG
ATGCGCTTCAGCATTCTATGCAGCACTGTGCAGCTGGGCTCAAGTACCTTTTATCCTTGGTATTCAGATTCT
TGGCCTGTGCCTTGGCAGCCTCCATCCTTCGCGAGGCATCTCATGCTTGGAAGTGTTTGGCCCTCAAGTTATAT
TTGAGGCTGTGGGCTTCATTGTGAGCAGCGTGGGACTTCTCCTGGGCATAGCTTTGGTGATGAGAGTGGATGGTG
CTGTGAGCTCCTGGTTCAGGCAGCTATTTCTGGCCCAGCAGAGGTAGCCTAGTCTGTGATTACTGGCACTTGGCT
ACAGAGAGTGTGGAGAACAGTGTAGCCTGGCCTGTACAGGTACTGGATGATCTGCAAGACAGGCTCAGCCATAC
TCTTACTATCATGCAGCCAGGGGCCGCTGACATCTAGGACTTCATTATTCTATAATTACAGGACCAGTGGAGTA
GTATCCCTACTCTGATTTGGATGCATCTGAGGACAAGGGGGCGGTCTCCGAAGTGAATAAAAATAGGCCGG
GCGTGGTGACTTGCACCTATAATCCCAGCACTTTGGGAGGCAGAGTGGGAGGATTGCTTGGTCCCAGGAGTTCA
AGACCAGCCTGTGGAACATAACAAGACCCCGTCTCTACTATTTAAAAAAAAGTGAATAAAAATGATAATAT

FIGURE 60

</usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA62809

<subunit 1 of 1, 1089 aa, 1 stop

<MW: 118699, pI: 8.49, NX(S/T): 2

MQKASVLLFLAWVCFLFYAGIALETSGLLTRLLELTNHSSCQEPPGPGSLPWGSQGKPGACW
 MASRFSRVVLVLIDALRFDAQPQSHVPREPPVSLPFLGKLSSLQRILEIQPHHARLYRSQ
 VDPPTTTMQRLKALTGSLPTFIDAGSNFASHAIVEDNLIKQLTSAGRRVVFMDGDDTWKDLF
 PGAFSKAFFFPNVRDLDTVDNGILEHLYPTMDSGEWDVLIHFLGVDHCGHKHGPHHP
 AKKLSQMDQVIQGLVERLENDTLLVVAGDHGTTNGDHGGDSELEVSAALFLYSPTAVFPST
 PPEEPEVIPQVSLVPTLALLLGLPIPFNGIGEVMAELEFSGGEDSQPHSSALAQASALHLNAQ
 QVSRFLHTYSAATQDLQAKELHQLQNLFSKASADYQWLLQSPKGAEATLPTVIAELQQFLRG
 ARAMCIESWARFSLVRMAGGTALLAASCFICLLASQWAI SPGFPCPLLLTPVAWGLVGAIA
 YAGLLGTIELKLDLVLGAVAAVSSFLPFLWKAWAGWGSKRPLATLFPIPGPVLLLLLFRLA
 VFFSDSFVVAEARATPFLGSGFILLLVQLHWEGQLLPKLLTMPRLGTSATTNPPRHNGAY
 ALRLGIGLLLCTRLAGLFHRCPEETPVCHSSPWLSPPLASMVGGRAKNLWYGACVAALVALLA
 AVRLWLRRYGNLKSPEPPMLFVRWGLPLMALGTAAWALASGADEAPPRLRVLVSGASMLVP
 RAVAGLAASGLALLLWKPVTVLVKAGAGAPRTRTVLTPFSGPPTSQADLDYVVPQIYRHMQE
 EFRGRLERTKSQGPLTVAAYQLGSVYSAAMVTALTLLAFPLLLLHAERISLVFLLLFLQSFL
 LLHLLAAGIPVTTGPGFTVPWQAVSAWALMATQTFYSTGHQPVFPAIHWHA AFVGFPEGHGS
 CTWLPALLVGANTFASHLLFAVGCPLLLLWPFLLCESQGLRKRQPPGNEADARVRPEEEEEEP
 LMEMRLRDAPQHFYAALLQLGLKYLFI LGIQILACALAASILRRHLMVWKVFAPKFI FEAVG
 FIVSSVGLLLGIALVMRVDGAVSSWFERQLFLAQQR

Important features:**Signal peptide:**

amino acids 1-16

Transmembrane domains:

amino acids 317-341, 451-470, 481-500, 510-527, 538-555, 831-850,
 1016-1034, 1052-1070

Leucine zipper pattern.

amino acids 843-864

N-glycosylation sites.

amino acids 37-40, 268-271

FIGURE 61

TGCCGCTGCCGCGCTGCTGCTGTTGCTCCTGGCGGCGCCTTGGGGACGGGCAGTTCCTGT
GTCTCTGGTGGTTTGCCTAAACCTGCAAACATCACCTTCTTATCCATCAACATGAAGAATG
CCTACAATGGACTCCACCAGAGGGTCTTCAAGGAGTTAAAGTTACTTACACTGTGCAGTATT
TCATCACAAATTGGCCACCAGAGGTGGCACTGACTACAGATGAGAAGTCCATTTCTGTTGT
CCTGACAGCTCCAGAGAAGTGGAAGAGAAATCCAGAAGACCTTCCTGTTTCCATGCAACAAA
TATACTCCAATCTGAAGTATAACGTGTCTGTGTTGAATACTAAATCAAACAGAACGTGGTCC
CAGTGTGTGACCAACCACACGCTGGTGCTCACCTGGCTGGAGCCGAACACTCTTTACTGCGT
ACACGTGGAGTCCTTCGTCCCAGGGCCCCCTCGCCGTGCTCAGCCTTCTGAGAAGCAGTGTG
CCAGGACTTTGAAAGATCAATCATCAGAGTTCAAGGCTAAAATCATCTTCTGGTATGTTTTG
CCCATATCTATTACCGTGTTTCTTTTTTCTGTGATGGGCTATTCCATCTACCGATATATCCA
CGTTGGCAAAGAGAAACACCCAGCAAATTTGATTTTGATTTATGGAAATGAATTTGACAAAA
GATTCTTTGTGCCTGCTGAAAAAATCGTGATTAACCTTTATCACCTCAATATCTCGGATGAT
TCTAAAATTTCTCATCAGGATATGAGTTTACTGGGAAAAAGCAGTGATGTATCCAGCCTTAA
TGATCCTCAGCCCAGCGGGAACCTGAGGCCCCCTCAGGAGGAAGAGGAGGTGAAACATTTAG
GGTATGCTTCGCATTTGATGGAAATTTTTTGTGACTCTGAAGAAAACACGGAAGGTACTTCT
CTCACCCAGCAAGAGTCCCTCAGCAGAACAATACCCCCGATAAAACAGTCATTGAATATGA
ATATGATGTCAGAACCACTGACATTTGTGCGGGGCTGAAGAGCAGGAGCTCAGTTTGCAGG
AGGAGGTGTCCACACAAGGAACATTATTGGAGTCGCAGGCAGCGTTGGCAGTCTTGGGCCCCG
CAAACGTTACAGTACTCATAACCCCTCAGCTCCAAGACTTAGACCCCTGGCGCAGGAGCA
CACAGACTCGGAGGAGGGGCCGAGGAAGAGCCATCGACGACCCTGGTCGACTGGGATCCCC
AAACTGGCAGGCTGTGTATTCTTCGCTGTCCAGCTTCGACCAGGATTCAGAGGGCTGCGAG
CCTTCTGAGGGGGATGGGCTCGGAGAGGAGGGTCTTCTATCTAGACTCTATGAGGAGCCGGC
TCCAGACAGGCCACCAGGAGAAAATGAAACCTATCTCATGCAATTCATGGAGGAATGGGGGT
TATATGTGCAGATGGAAAACATGATGCCAACACTTCCTTTTGCCTTTTGTTCCTGTGCAAC
AAGTGAGTCACCCCTTTGATCCCAGCCATAAAGTACCTGGGATGAAAGAAGTTTTTTCCAGT
TTGTCAGTGTCTGTGAGAATTACTTATTTCTTTTCTCTATTCTCATAGCACGTGTGTGATTG
GTTTCATGCATGTAGGTCTCTTAACAATGATGGTGGGCCTCTGGAGTCCAGGGGCTGGCCGGT
TGTTCTATGCAGAGAAAGCAGTCAATAAATGTTTGCCAGACTGGGTGCAGAAATTTATTCAGG
TGGGTGT

FIGURE 62

</usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA62815

<subunit 1 of 1, 442 aa, 1 stop

<MW: 49932, pI: 4.55, NX(S/T): 5

MSYNGLHQRVFKELKLLTLCSSISSQIGPPEVALTTDEKSISVVLTAPEKWKRNPEDLPVSMQ
QIYSNLKYNVSVLNTKSNRTWSQCVTNHTLVLTWLEPNTLYCVHVESFVPGPPRAQPSEKQ
CARTLKDQSSEFKAKIIFWYVLPISITVFLEFSVMGYSIYRYIHVGKEKHPANLILYGNED
KRFFVPAEKIVINFITLNISSDDSKISHQDMSLLGKSSDVSSLNDPQPSGNLRPPQEEEEVKH
LGYASHLMEIFCDSEENTEGTSLTQQESLSRTIPDKTVIEYEYDVRTTDICAGPEEQELSL
QEEVSTQGTLLSQAALAVLGPQTLQYSYTPQLQDLDPQAQHTDSEEGPEEEPPSTTLVDWD
PQTGRCLCIPSLSSFDQDSEGCEPSEGDGLGEEGLLSRLYEPPAPDRPPGENETYLMQFMEEW
GLYVQMEN

Important features:**Signal peptide:**

amino acids 1-28

Transmembrane domain:

amino acids 140-163

N-glycosylation sites.

amino acids 71-74, 80-83, 89-92, 204-207, 423-426

FIGURE 63

CGGACGCGTGGGCGGACGCGTGGGCGGACGCGTGGGTCTCTGCGGGGAGACGCCAGCCTGCG
TCTGCC**ATG**GGGCTCGGGTTGAGGGGCTGGGGACGTCCTCTGCTGACTGTGGCCACCGCCCT
GATGCTGCCCCGTGAAGCCCCCGCAGGCTCCTGGGGGGCCAGATCATCGGGGGCCACGAGG
TGACCCCCCACTCCAGGCCCTACATGGCATCCGTGCGCTTCGGGGGGCCAACATCACTGCGGA
GGCTTCCTGCTGCGAGCCCGCTGGGTGGTCTCGGCCGCCCACTGCTTCAGCCACAGAGACCT
CCGCACTGGCCTGGTGGTGCTGGGCGCCACGTCCTGAGTACTGCGGAGCCCACCCAGCAGG
TGTTTGGCATCGATGCTCTCACCACGCACCCCGACTACCACCCCATGACCCACGCCAACGAC
ATCTGCCTGCTGCGGCTGAACGGCTCTGCTGTCTGGGCCCTGCAGTGGGGCTGCTGAGGCT
GCCAGGGAGAAGGGCCAGGCCCCCACAGCGGGGACACGGTGCCGGGTGGCTGGCTGGGGCT
TCGTGTCTGACTTTGAGGAGCTGCCGCCTGGACTGATGGAGGCCAAGGTCCGAGTGCTGGAC
CCGGACGTCTGCAACAGCTCCTGGAAGGGCCACCTGACACTTACCATGCTCTGCACCCGCGAG
TGGGGACAGCCACAGACGGGGCTTCTGCTCGGCCGACTCCGGAGGGGCCCTGGTGTGCAGGA
ACCGGGCTCACGGCCTCGTTTTCCTTCTCGGGCCTCTGGTGCGGCGACCCCAAGACCCCGAC
GTGTACACGCAGGTGTCCGCCTTTGTGGCCTGGATCTGGGACGTGGTTTCGGCGGAGCAGTCC
CCAGCCCGGGCCCCCTGCCTGGGACCACCAGGCCCCCAAGGAGAAGCCGCC**TGA**GCCACAACCT
TGCGGCATGCAAATGAGATGGCCGCTCCAGGCCTGGAATGTTCCGTGGCTGGGCCCCACGGG
AAGCCTGATGTTTCAGGGTTGGGGTGGGACGGGCAGCGGTGGGGCACACCCATTCCACATGCA
AAGGGCAGAAGCAAACCCAGTAAAATGTAACTGACAAAAAAAAAAAAAAAAAAAAAGAAA

FIGURE 64

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA62845
><subunit 1 of 1, 283 aa, 1 stop
><MW: 30350, pI: 9.66, NX(S/T): 2
MGLGLRGWGRPLLTVATALMLPVKPPAGSWGAIIGGHEVTPHSRPYMASVRFGGQHHCGGF
LLRARWVVSAAHCFSHRDLRTGLVVLGAHVLSTAEP TQQVFGIDALTTHPDYHPMTHANDIC
LLRLNGSAVLGPAVGLLRLPGRRARPPTAGTRCRVAGWGFVSDFEELPPGLMEAKVRVLDPD
VCNSSWKGH LTLTMLCTRSGDSHRRGFCSADSGGPLVCRNRAHGLVSFSGLWCGDPKTPDVY
TQVSAFVAWIWDVVRSSPQPGPLPGTTRPPGEAA
```

Signal peptide:

amino acids 1-30

bioRxiv preprint doi: <https://doi.org/10.1101/000000>; this version posted January 1, 2014. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.

FIGURE 65

GAGCTACCCAGGCGGCTGGTGTGTCAGCAAGCTCCGCGCCGACTCCGGACGCCTGACGCCTGA
CGCCTGTCCCCGGGCCCGGCATGAGCCGCTACCTGCTGCCGCTGTCTGGCGCTGGGCACGGTAG
CAGGCGCCGCGCTGCTGCTCAAGGACTATGTACCGGTGGGGCTTGCCCCAGCAAGGCCACC
ATCCCTGGGAAGACGGTCATCGTGACGGGCGCCAACACAGGCATCGGGAAGCAGACCGCCTT
GGA ACTGGCCAGGAGAGGAGGCAACATCATCCTGGCCTGCCGAGACATGGAGAAGTGTGAGG
CGGCAGCAAAGGACATCCGCGGGGAGACCCTCAATCACCATGTCAACGCCCGGCACCTGGAC
TTGGCTTCCCTCAAGTCTATCCGAGAGTTTGCAGCAAAGATCATTGAAGAGGAGGAGCGAGT
GGACATTCTAATCAACAACGCGGGTGTGATGCGGTGCCCCACTGGACCACCGAGGACGGCT
TCGAGATGCAGTTTGGCGTTAACCACCTGGGTCACTTTCTCTTGACAAACTTGCTGCTGGAC
AAGCTGAAAGCCTCAGCCCCCTTCGCGGATCATCAACCTCTCGTCCCTGGCCATGTTGCTGG
GCACATAGACTTTGACGACTTGA ACTGGCAGACGAGGAAGTATAACACCAAAGCCGCCTACT
GCCAGAGCAAGCTCGCCATCGTCCTCTTCACCAAGGAGCTGAGCCGGCGGCTGCAAGGCTCT
GGTGTGACTGTCAACGCCCTGCACCCCGGCGTGGCCAGGACAGAGCTGGGCAGACACACGGG
CATCCATGGCTCCACCTTCTCCAGCACCACTCGGGCCCATCTTCTGGCTGCTGGTCAAGA
GCCCCGAGCTGGCCGCCAGCCCAGCACATACCTGGCCGTGGCGGAGGAACTGGCGGATGTT
TCCGGAAAGTACTTCGATGGACTCAAACAGAAGGCCCGGCCCGAGGCTGAGGATGAGGA
GGTGGCCCGGAGGCTTTGGGCTGAAAGTGCCCGCCTGGTGGGCTTAGAGGCTCCCTCTGTGA
GGGAGCAGCCCCTCCCCAGATAACCTCTGGAGCAGATTTGAAAGCCAGGATGGCGCCTCCAG
ACCGAGGACAGCTGTCCGCCATGCCCGCAGCTTCCTGGCACTACCTGAGCCGGGAGACCCAG
GACTGGCGGCCGCCATGCCCGCAGTAGGTTCTAGGGGGCGGTGCTGGCCGCAGTGGACTGGC
CTGCAGGTGAGCACTGCCCCGGGCTCTGGCTGGTTCCGTCTGCTCTGCTGCCAGCAGGGGAG
AGGGGCCATCTGATGCTTCCCCTGGGAATCTAAACTGGGAATGGCCGAGGAGGAAGGGGCTC
TGTGCACTTGCAGGCCACGTCAGGAGAGCCAGCGGTGCCTGTCTGGGGAGGGTTCCAAGGTGC
TCCGTGAAGAGCATGGGCAAGTTGTCTGACACTTGGTGGATTCTTGGGTCCCTGTGGGACCT
TGTGCATGCATGGTCCTCTCTGAGCCTTGGTTTCTTCAGCAGTGAGATGCTCAGAATAACTG
CTGTCTCCCATGATGGTGTGGTACAGCGAGCTGTTGTCTGGCTATGGCATGGCTGTGCCGGG
GGTGT TTGCTGAGGGCTTCCTGTGCCAGAGCCCAGCCAGAGAGCAGGTGCAGGTGTCATCCC
GAGTTCAGGCTCTGCACGGCATGGAGTGGGAACCCACACAGCTGCTGCTACAGGACCTGGGA
TTGCCTGGGACTCCACCTTCCTATCAATTCTCATGGTAGTCCAAACTGCAGACTCTCAAAC
TTGCTCATTT

FIGURE 66

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA64842
><subunit 1 of 1, 331 aa, 1 stop
><MW: 35932, pI: 8.45, NX(S/T): 1
MSRYLLPLSALGTVAGAAVLLKDYVTGGACPSKATIPGKTVIVTGANTGIGKQTALELARRG
GNIILACRDMEKCEAAAKDIRGETLNHHVNARHLDLASLKSIREFAAKIIEEEEERVDILINN
AGVMRCPHWTTEDGFEMQFGVNHLGHFLLTNLLLDKLGKASAPSRIINLSSLAHVAGHIDFDD
LNWQTRKYNTKAAYCQSKLAIVLFTKELSRRLQSGSVTVNALHPGVARTELGRHTGIHGSTF
SSTTLGPIFWLLVKSPELAAQPSTYLAVAEELADVSGKYFDGLKQKAPAPEAEDEEVARRLW
AESARLVGLEAPSVREQPLPR
```

Signal peptide:

amino acids 1-17

Downloaded from www.oxfordjournals.org/ by guest on June 11, 2015

FIGURE 67

GAAGTTCGCGAGCGCTGGC**ATG**TGGTCCTGGGGCGCGGCTGGCGGCGCTGCTGGCGGTGCTG
 GCGCTCGGGACAGGAGACCCAGAAAGGGCTGCGGCTCGGGGCGACACGTTCTCGGCGCTGAC
 CAGCGTGGCGCGCGCCCTGGCGCCCGAGCGCCGGCTGCTGGGGCTGCTGAGGCGGTACCTGC
 GCGGGGAGGAGGCGCGGCTGCGGGACCTGACTAGATTCTACGACAAGGTACTTTCTTTGCAT
 GAGGATTCAACAACCCCTGTGGCTAACCCCTCTGCTTGCAATTTACTCTCATCAAACGCCTGCA
 GTCTGACTGGAGGAATGTGGTACATAGTCTGGAGGCCAGTGAGAACATCCGAGCTCTGAAGG
 ATGGCTATGAGAAGGTGGAGCAAGACCTTCCAGCCTTTGAGGACCTTGAGGGAGCAGCAAGG
 GCCCTGATGCGGCTGCAGGACGTGTACATGCTCAATGTGAAAGGCCTGGCCCCGAGGTGTCTT
 TCAGAGAGTCACTGGCTCTGCCATCACTGACCTGTACAGCCCCAAACGGCTCTTTTCTCTCA
 CAGGGGATGACTGCTTCCAAGTTGGCAAGGTGGCCTATGACATGGGGGATTATTACCATGCC
 ATTCCATGGCTGGAGGAGGCTGTCACTCTTCCGAGGATCTTACGGAGAGTGGAAGACAGA
 GGATGAGGCAAGTCTAGAAGATGCCTTGGATCACTTGGCCTTTGCTTATTTCCGGGCAGGAA
 ATGTTTCGTGTGCCCTCAGCCTCTCTCGGGAGTTTCTTCTCTACAGCCCAGATAATAAGAGG
 ATGGCCAGGAATGTCTTGAAATATGAAAGGCTCTTGGCAGAGAGCCCCAACACGTGGTAGC
 TGAGGCTGTCATCCAGAGGCCCAATATACCCACCTGCAGACCAGAGACACCTACGAGGGGC
 TATGTCTAGACCCTGGGTTCCCAGCCCCTCTCTACCAGATCCCTAGCCTCTACTGTTCCCTAT
 GAGACCAATTCCAACGCCTACCTGCTGCTCCAGCCCATCCGGAAGGAGGTCTACACCTGGA
 GCCCTACATTGCTCTCTACCATGACTTCGTCTAGTCACTCAGAGGCTCAGAAAATTAGAGAAC
 TTGCAGAACCATGGCTACAGAGGTCAGTGGTGGCATCAGGGGAGAAGCAGTTACAAGTGGAG
 TACCGCATCAGCAAAAGTGCCTGGCTGAAGGACACTGTTGACCCAAAACCTGGTGACCCTCAA
 CCACCGCATTGCTGCCCTCACAGGCCTTGATGTCCGGCCTCCCTATGCAGAGTATCTGCAGG
 TGGTGAACCTATGGCATCGGAGGACACTATGAGCCTCACTTTGACCATGCTACGTACCAAGC
 AGCCCCCTCTACAGAATGAAGTCAGGAAACCGAGTTGCAACATTTATGATCTATCTGAGCTC
 GGTGGAAGCTGGAGGAGCCACAGCCTTCATCTATGCCAACCTCAGCGTGCCCTGTGGTTAGGA
 ATGCAGCACTGTTTTGGTGGAACTGCACAGGAGTGGTGAAGGGGACAGTGACACACTTCAT
 GCTGGCTGTCCTGTCCTGGTGGGAGATAAGTGGGTGGCCAACAAGTGGATACATGAGTATGG
 ACAGGAATTCCGCAGACCCTGCAGCTCCAGCCCTGAAGAC**TGA**ACTGTTGGCAGAGAGAAGC
 TGGTGGAGTCCCTGTGGCTTTCCAGAGAAGCCAGGAGCCAAAAGCTGGGGTAGGAGAGGAGAA
 AGCAGAGCAGCCTCCTGGAAGAAGGCCTTGTCAGCTTTGTCTGTGCCTCGCAAATCAGAGGC
 AAGGGAGAGGTTGTTACCAGGGGACACTGAGAATGTACATTTGATCTGCCCCAGCCACGGAA
 GTCAGAGTAGGATGCACAGTACAAAGGAGGGGGAGTGGAGGCCTGAGAGGGAAGTTTCTGG
 AGTTCAGATACTCTCTGTTGGGAACAGGACATCTCAACAGTCTCAGGTTTCGATCAGTGGGTC
 TTTTGGCACTTTGAACCTTGACCACAGGGACCAAGAAGTGGCAATGAGGACACCTGCAGGAG
 GGGCTAGCCTGACTCCAGAACTTTAAGACTTTCTCCCCACTGCCTTCTGCTGCAGCCCAAG
 CAGGGAGTGTCCCCCTCCAGAAAGCATATCCAGATGAGTGGTACATTATATAAGGATTTTTT
 TTTAAGTTGAAAACAACCTTTCTTTTCTTTTGTATGATGGTTTTTTAACACAGTCATTAAAA
 ATGTTTATAAATCAAAA

FIGURE 68

MGPGARLAALLAVLALGTGDPERAAARGDTFSALTSVARALAPERRLGLLRRYLRGEEARL
RDLTRFYDKVLSLHEDSTTPVANPLLAFTLIKRLQSDWRNVVHSLEASENIRALKDGYEKVE
QDLPAFEDLEGAARALMRLQDVYMLNVKGLARGVFQRTGSAITDLYSPKRLFSLTGDDCFQ
VGKVAYDMGDYYHAIPWLEEAVSLFRGSYGEWKTEDEASLEDALDHLAFAYFRAGNVSCALS
LSREFLLYSPDNKRMARNVLKYERLLAESPNHVVAEAVIQRPNIPHLQTRDTYEGLCQTLGS
QPTLYQIPSLYCSYETNSNAYLLLQPIRKEVIHLEPYIALYHDFVSDSEAQKIRELAEPWLQ
RSVVASGEKQLQVEYRISKSAWLKDTVDPKLVTLNHRIAALTGLDVRPPYAEYLQVVNYGIG
GHYEPHFDHATSPSSPLYRMKSGNRVATFMIYLSSVEAGGATAFIYANLSVPVVRNAALFWW
NLHRSGEGDSDTLHAGCPVLVGDKWVANKWIHEYGQEFRRPCSSSPED

Signal peptide:

amino acids 1-19

FIGURE 69

GAGATAGGGAGTCTGGGTTTAAGTTCCCTGCTCCATCTCAGGAGAGCCCTTGGTCCCACCCCTAG
GAAGCCACCAGACTCCACGGTGTGGGGCCAATCAGGTGGAATCGGCCCTGGCAGGTGGGGCC
ACGAGCGCTGGCTGAGGGACCGAGCCGGAGAGCCCCGGAGCCCCCGTAACCCGCGCGGGGGAG
CGCCCAGGATGCCGCGCGGGGGACTCGGAGCAGGTGCGCTACTGCGCGCGCTTCTCCTACCTC
TGGCTCAAGTTTTTCACTTATCATCTATTCCACCGTGTTCTGGCTGATTGGGGCCCTGGTCCT
GTCTGTGGGCATCTATGCAGAGGTTGAGCGGCAGAAATATAAAACCCCTTGAAAGTGCCTTCC
TGGCTCCAGCCATCATCCTCATCCTCCTGGGCGTCGTCATGTTTCATGGTCTCCTTCATTGGT
GTGCTGGCGTCCCTCCGTGACAACCTGTACCTTCTCCAAGCATTCATGTACATCCTTGGGAT
CTGCCTCATCATGGAGCTCATTGGTGGCGTGGTGGCCTTGACCTTCCGGAACCAGACCATTG
ACTTCCTGAACGACAACATTTCGAAGAGGAATTGAGAACTACTATGATGATCTGGACTTCAA
AACATCATGGACTTTGTTCAGAAAAAGTTCAAGTGCTGTGGCGGGGAGGACTACCGAGATTG
GAGCAAGAATCAGTACCACGACTGCAGTGCCCTGGACCCCTGGCCTGTGGGGTGCCCTACA
CCTGCTGCATCAGGAACACGACAGAAGTTGTCAACACCATGTGTGGCTACAAAACCTATCGAC
AAGGAGCGTTTCAGTGTGCAGGATGTCATCTACGTGCGGGGCTGCACCAACGCCGTGATCAT
CTGGTTTCATGGACAACCTACACCATCATGGCGTGCATCCTCCTGGGCATCCTGCTTCCCCAGT
TCCTGGGGGTGCTGCTGACGCTGCTGTACATCACCCGGGTGGAGGACATCATCATGGAGCAC
TCTGTCACTGATGGGCTCCTGGGGCCCCGGTGCCAAGCCCAGCGTGGAGGCGGCAGGCACGGG
ATGCTGCTTGTGCTACCCCAATTAGGGGCCCAGCCTGCCATGGCAGCTCCAACAAGGACCGTC
TGGGATAGCACCTCTCAGTCAACATCGTGGGGCTGGACAGGGCTGCGGCCCTCTGCCACA
CTCAGTACTGACCAAAGCCAGGGCTGTGTGTGCCTGTGTGTAGGTCCCACGGCCTCTGCCTC
CCCAGGGAGCAGAGCCTGGGCCTCCCCTAAGAGGCTTTCCCCGAGGCAGCTCTGGAATCTGT
GCCCACCTGGGGCCTGGGGAACAAGGCCCTCCTTTCTCCAGGCCTGGGCTACAGGGGAGGGA
GAGCCTGAGGCTCTGCTCAGGGCCCATTTTCATCTCTGGCAGTGCCTTGGCGGTGGTATTCAA
GGCAGTTTTTGTAGCACCTGTAATTGGGGAGAGGGAGTGTGCCCTCGGGGCAGGAGGGAAGG
GCATCTGGGGAAGGGCAGGAGGGAAGAGCTGTCCATGCAGCCACGCCATGGCCAGGTGGC
CTCTTCTCAGCCTCCCAGGTGCCTTGAGCCCTCTTGCAAGGGCGGCTGCTTCCTTGAGCCTA
GTTTTTTTTTACGTGATTTTTGTAAACATTCATTTTTTTGTACAGATAACAGGAGTTTCTGAC
TAATCAAAGCTGGTATTTCCCCGCATGTCTTATTCTTGCCCTTCCCCCAACCAGTTTGTTAA
TCAAACAATAAAAAACATGTTTTGTTTTGTTTTTAAAAA

FIGURE 70

></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA64863

><subunit 1 of 1, 294 aa, 1 stop

><MW: 33211, pI: 5.35, NX(S/T): 3

MPRGDSEQVRYCARFSYLWLKFSLLIYSTVFWLIGALVLSVGIYAEVERQKYKTLES AFLAP
AIILILLGVVMFMVSFIGVLASLRDNLVLLQAFMYILGICLIMELIGGVVALTFRNQ TIDFL
NDNIRRG IENYYDDLDFKNIMDFVQKKFKCCGGEDYRDWSKNQYHDCSAPG PLACGV PYTCC
IRNTTEVVNTMCGYKTIDKERFSVQDVIYVRGCTNAVIIWFMDNYTIMACILLGILLPQFLG
VLLTLLYITRVEDIIMEHSVTDGLLGPGAKPSVEAAGTGCCLCYPN

Signal peptide:

amino acids 1-44

Transmembrane domains:

amino acids 22-42, 57-85, 93-116, 230-257

An analysis of the Dayhoff database (version 35.45 SwissProt 35), using a WU-BLAST2 sequence alignment analysis of the full-length sequence shown in Figure 198 (SEQ ID NO:338), revealed some homology between the PRO1555 amino acid sequence and the following Dayhoff sequences: YKA4_CAEEL, AB014541_1, HVSX99518_2, SSU63019_1, GEN14286, MMU68267_1, XP2_XENLA, ICP4_HSV11, P_W40200, and AE001360_1.

5 Clone DNA73744-1665 was deposited with the ATCC on October 6, 1998, and is assigned ATCC deposit no. 203322.

EXAMPLE 103: Isolation of cDNA clones Encoding Human PRO1485

10 A consensus DNA sequence was assembled relative to other EST sequences using phrap as described in Example 1 above. This consensus sequence is designated herein "DNA44791". Based on the DNA44791 consensus sequence, oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO1485.

PCR primers (2 forward and 2 reverse) were synthesized:

15 forward PCR primer 1: 5'CCCTCCAAGGATGACAAAGGCGC 3' (SEQ ID NO:341);

forward PCR primer 2: 5'GGTCAGCAGCTTTCTTGCCCTAAATCAGG 3' (SEQ ID NO:342);

reverse PCR primer 1: 5'ATCTCAGGCGGCATCCTGTCAGCC 3' (SEQ ID NO:343); and

reverse PCR primer 2: 5'GTGGATGCCTGCAAGAAGGTTGGG 3' (SEQ ID NO:344).

20 Additionally, a synthetic oligonucleotide hybridization probe was constructed from the consensus DNA44791 sequence which had the following nucleotide sequence:

hybridization probe 5'AGCTTTCTTGCCCTAAATCAGGCCAGCCTCATCAGTCGCTGTGAC 3' (SEQ ID NO:345)

25 In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pair identified above. A positive library was then used to isolate clones encoding the PRO1485 gene using the probe oligonucleotide and one of the PCR primers. RNA for construction of the cDNA libraries was isolated from human testis.

DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO1485 (designated herein as DNA73746-1654 [Figure 199, SEQ ID NO:339]; and the derived protein sequence for PRO1485.

30 The entire coding sequence of PRO1485 is shown in Figure 199 (SEQ ID NO:339). Clone DNA73746-1654 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 151-153 and an apparent stop codon at nucleotide positions 595-597 of SEQ ID NO:339. The predicted polypeptide precursor is 148 amino acids long. The signal peptide is at about amino acids 1-18 of SEQ ID NO:340. The lysozyme C signature, CAAX box, and an N-glycosylation site are shown in Figure 200. Clone
35 DNA73746-1654 has been deposited with ATCC and is assigned ATCC deposit no. 203411. The full-length PRO1485 protein shown in Figure 200 has an estimated molecular weight of about 16,896 daltons and a pI of about 6.05.

An analysis of the Dayhoff database (version 35.45 SwissProt 35), using a WU-BLAST2 sequence alignment analysis of the full-length sequence shown in Figure 200 (SEQ ID NO:340), revealed sequence identity

between the PRO1485 amino acid sequence and the following Dayhoff sequences: LYC_PHACO, P_R76684, 2HFL_Y, JC2144, JC5544, JC5555, JC5369, LYC2_PIG, P_R12113, and JC5380.

EXAMPLE 104: Isolation of cDNA clones Encoding Human PRO1564

A consensus DNA sequence was assembled relative to other EST sequences using phrap as described in Example 1 above. This consensus sequence is herein designated DNA67213. Based on the DNA67213 consensus sequence, oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO1564.

PCR primers (forward and reverse) were synthesized:

forward PCR primer (67213.f1) 5'-GGAGAGGTGGTGGCCATGGACAG-3' (SEQ ID NO:348)

reverse PCR primer (67213.r1) 5'-CTGTCACTGCAAGGAGCCAACACC-3' (SEQ ID NO:349)

Additionally, a synthetic oligonucleotide hybridization probe was constructed from the consensus DNA67213 sequence which had the following nucleotide sequence

hybridization probe (67213.p1)

5'-TATGTCGCTGCGAGGTGGTGAAAACCTCGAACTGTCTTTCAAGGC-3' (SEQ ID NO:350)

In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pair identified above. A positive library was then used to isolate clones encoding the PRO1564 gene using the probe oligonucleotide and one of the PCR primers. RNA for construction of the cDNA libraries was isolated from human breast carcinoma tissue.

DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO1564 (designated herein as DNA73760-1672 [Figure 201, SEQ ID NO:346]; and the derived protein sequence for PRO1564.

The entire nucleotide sequence of DNA73760-1672 is shown in Figure 201 (SEQ ID NO:346). Clone DNA73760-1672 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 462-464 and ending at the stop codon at nucleotide positions 2379-2381 (Figure 201). The predicted polypeptide precursor is 639 amino acids long (Figure 202). The full-length PRO1564 protein shown in Figure 202 has an estimated molecular weight of about 73,063 daltons and a pI of about 6.84. Analysis of the full-length PRO1564 sequence shown in Figure 202 (SEQ ID NO:347) evidences the presence of the following: a signal peptide from about amino acid 1 to about amino acid 28, a transmembrane domain from about amino acid 11 to about amino acid 36, potential N-glycosylation sites from about amino acid 107 to about amino acid 110 and from about amino acid 574 to about amino acid 577, a tyrosine kinase phosphorylation site from about amino acid 50 to about amino acid 57, potential N-myristoylation sites from about amino acid 158 to about amino acid 163, from about amino acid 236 to about amino acid 241, from about amino acid 262 to about amino acid 267, from about amino acid 270 to about amino acid 275, from about amino acid 380 to about amino acid 385 and from about amino acid 513 to about amino acid 518, an amidation site from about amino acid 110 to about amino acid 113 and a prokaryotic membrane lipoprotein lipid attachment site from about amino acid 15 to about amino acid 25. Clone DNA73760-1672 has been deposited with ATCC on October 6, 1998 and is assigned ATCC deposit no.203314.

An analysis of the Dayhoff database (version 35.45 SwissProt 35), using a WU-BLAST2 sequence

alignment analysis of the full-length sequence shown in Figure 202 (SEQ ID NO:347), evidenced significant homology between the PRO1564 amino acid sequence and the following Dayhoff sequences: MMU73819_1, HSY08564_1, P_W34470, P_R66402, PAGT_HUMAN, CEGLY5B_1, CEGLY6A_1, CEGLY6B_1, AP000006_308 and E69322.

5 EXAMPLE 105: Isolation of cDNA clones Encoding Human PRO1755

Use of the signal sequence algorithm described in Example 3 above allowed identification of an EST cluster sequence from the LIFESEQ® database, designated EST Cluster No. 141872. This EST cluster sequence was then compared to a variety of ESTs from the databases listed above to identify existing homologies. The homology search was performed using the computer program BLAST or BLAST2 (Altschul et al., Methods in
10 Enzymology 266:460-480 (1996)). Those comparisons resulting in a BLAST score of 70 (or in some cases 90) or greater that did not encode known proteins were clustered and assembled into a consensus DNA sequence with the program "phrap" (Phil Green, University of Washington, Seattle, Washington). The consensus sequence obtained therefrom is herein designated "DNA55731".

15 In light of the sequence homology between the DNA55731 sequence and a sequence contained within Incyte EST no. 257323, the EST clone was purchased and the cDNA insert was obtained and sequenced. Incyte clone 257323 was derived from a library constructed using RNA isolated from the hNT2 cell line (Stratagene library no. STR9372310), which was derived from a human teratocarcinoma that exhibited properties characteristic of a committed neuronal precursor at an early stage of development. The sequence of this cDNA
20 insert is shown in Figure 203 and is herein designated "DNA76396-1698". Alternatively, the DNA76396-1698 sequence can be obtained by preparing oligonucleotide probes and primers and isolating the sequence from an appropriate library (e.g. STR9372310).

The full length clone shown in Figure 203 contained a single open reading frame with an apparent translational initiation site at nucleotide positions 58 to 60 and ending at the stop codon found at nucleotide
25 positions 886 to 888 (Figure 203; SEQ ID NO:351). The predicted polypeptide precursor (Figure 204, SEQ ID NO:352) is 276 amino acids long. PRO1755 has a calculated molecular weight of approximately 29,426 daltons and an estimated pI of approximately 9.40. Additional features include: a signal peptide sequence at about amino acids 1-31; a transmembrane domain at about amino acids 178-198; a cAMP and cGMP-dependent protein kinase phosphorylation site at about amino acids 210-213; potential N-myristoylation sites at about amino
30 acids 117-122, 154-149, and 214-219; and a cell attachment sequence at about amino acids 149-151.

An analysis of the Dayhoff database (version 35.45 SwissProt 35), using a WU-BLAST2 sequence alignment analysis of the full-length sequence shown in Figure 204 (SEQ ID NO:352), revealed some homology between the PRO1755 amino acid sequence and the following Dayhoff sequences: APG-BRANA, P_R37743, NAU88587_1, YHL1_EBV, P_W31855, CET10B10_4, AF039404_1, PRP1_HUMAN, AF038575_1, and
35 AF053091_1.

Clone DNA76396-1698 was deposited with the ATCC on November 17, 1998, and is assigned ATCC deposit no. 203471.

EXAMPLE 106: Isolation of cDNA clones Encoding Human PRO1757

Use of the signal sequence algorithm described in Example 3 above allowed identification of three EST sequences from the Incyte database, designated Incyte EST clones no. 2007947, 2014962 and 1912034. These EST sequences were then clustered and assembled into a consensus DNA sequence with the program "phrap" (Phil Green, University of Washington, Seattle, Washington). The consensus sequence obtained therefrom is herein designated as DNA56054.

In light of the sequence homology between the DNA56054 sequence and a sequence contained within the Incyte EST clone no. 2007947, the Incyte EST clone no. 2007947 was purchased and the cDNA insert was obtained and sequenced. The sequence of this cDNA insert is shown in Figure 205 and is herein designated as DNA76398-1699.

Clone DNA76398-1699 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 59-61 and ending at the stop codon at nucleotide positions 422-424 (Figure 205). The predicted polypeptide precursor is 121 amino acids long (Figure 206). The full-length PRO1757 protein shown in Figure 206 has an estimated molecular weight of about 12,073 daltons and a pI of about 4.11. Analysis of the full-length PRO1757 sequence shown in Figure 206 (SEQ ID NO:354) evidences the presence of the following: a signal peptide from about amino acid 1 to about amino acid 19, a transmembrane domain from about amino acid 91 to about amino acid 110, a glycosaminoglycan attachment site from about amino acid 44 to about amino acid 47, a cAMP- and cGMP-dependent protein kinase phosphorylation site from about amino acid 116 to about amino acid 119 and a potential N-myristylation site from about amino acid 91 to about amino acid 96. Clone DNA76398-1699 has been deposited with ATCC on November 17, 1998 and is assigned ATCC deposit no. 203474.

An analysis of the Dayhoff database (version 35.45 SwissProt 35), using a WU-BLAST2 sequence alignment analysis of the full-length sequence shown in Figure 206 (SEQ ID NO:354), evidenced significant homology between the PRO1757 amino acid sequence and the following Dayhoff sequences: JQ0964, COLL_HSVS7, HSU70136_1, AF003473_1, D89728_1, MTF1_MOUSE, AF029777_1, HSU88153_1 and P_W05321.

EXAMPLE 107: Isolation of cDNA clones Encoding Human PRO1758

Use of the signal sequence algorithm described in Example 3 above allowed identification of an EST cluster sequence from the LIFESEQ® database, designated EST cluster No. 20926. This EST cluster sequence was then compared to a variety of expressed sequence tag (EST) from the databases mentioned above, to identify existing homologies. The homology search was performed using the computer program BLAST or BLAST2 (Altschul et al., Methods in Enzymology 266:460-480 (1996)). Those comparisons resulting in a BLAST score of 70 (or in some cases 90) or greater that did not encode known proteins were clustered and assembled into a consensus DNA sequence with the program "phrap" (Phil Green, University of Washington, Seattle, Washington). The consensus sequence obtained therefrom is herein designated DNA56260.

In light of the sequence homology between the DNA56260 sequence and a sequence contained within EST no. 2936330 from the LIFESEQ® database, the EST clone, which originated from a library constructed from thymus tissue of a fetus that died from anencephalus, was purchased and the cDNA insert was obtained and sequenced. The sequence of this cDNA insert is shown in Figure 207 and is herein designated as DNA76399-1700.

The full length clone shown in Figure 207 contained a single open reading frame with an apparent translational initiation site at nucleotide positions 78 to 80 and ending at the stop codon found at nucleotide positions 549-551 (Figure 207; SEQ ID NO:355). The predicted polypeptide precursor (Figure 208, SEQ ID NO:356) is 157 amino acids long. PRO1758 has a calculated molecular weight of approximately 17,681 daltons and an estimated pI of approximately 7.65. Additional features include: a signal peptide from about amino acids 1-15; a potential N-glycosylation site at about amino acids 24-27; a cAMP- and cGMP-dependent protein kinase phosphorylation site at about amino acids 27-30; a casein kinase II phosphorylation site at about amino acids 60-63; potential N-myristoylation sites at about amino acids 17-22, 50-55, 129-134, and 133-138; a cell attachment sequence at about amino acids 153-155; and a cytochrome c family heme-binding site signature at about amino acids 18-23.

An analysis of the Dayhoff database (version 35.45 SwissProt 35), using a WU-BLAST2 sequence alignment analysis of the full-length sequence shown in Figure 208 (SEQ ID NO:356), revealed significant homology between the PRO1758 amino acid sequence and Dayhoff sequence no AC005328_2. Homology was also found between the PRO1758 amino acid sequence and Dayhoff sequence no. CELC46F2_1.

Clone DNA76399-1700 was deposited with the ATCC on November 17, 1998 and is assigned ATCC deposit no. 203472.

EXAMPLE 108: Isolation of cDNA clones Encoding Human PRO1575

A consensus DNA sequence was assembled relative to other EST sequences using phrap as described in Example 1 above. This consensus sequence is designated herein as "DNA35699". Based on the DNA35699 consensus sequence, oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO1575.

PCR primers (forward and reverse) were synthesized:

forward PCR primers: CCAGCAGTGCCCATACTCCATAGC (35699.f1; SEQ ID NO:359);
TGACGAGTGGGATACACTGC (35699.f2; SEQ ID NO:360)

reverse PCR primer: GCTCTACGGAACTTCTGCTGTGG (35699.r1; SEQ ID NO:361)

Additionally, a synthetic oligonucleotide hybridization probe was constructed from the consensus DNA35699 sequence which had the following nucleotide sequence:

hybridization probe: ATTCCCAGGCGTGTCAATTTGGGATCAGCACTGATTCTGAGGTTCTGACAC
(35699.p1; SEQ ID NO:362)

In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pair identified above. A positive library was then used to isolate clones encoding the PRO1575 gene using the probe oligonucleotide and one of the PCR primers. RNA for construction of the cDNA libraries was isolated from human pancreatic tissue.

DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO1575 (designated herein as DNA76401-1683 [Figure 209, SEQ ID NO:357]; and the derived protein sequence for PRO1575.

The entire coding sequence of PRO1575 is shown in Figure 209 (SEQ ID NO:357). Clone DNA76401-1683 contains a single open reading frame with an apparent translational initiation site at nucleotide positions

22-24 and an apparent stop codon at nucleotide positions 841-843. The predicted polypeptide precursor is 273 amino acids long. The full-length PRO1575 protein shown in Figure 210 has an estimated molecular weight of about 30,480 daltons and a pI of about 4.60. Additional features include: a signal peptide at about amino acids 1-20; a transmembrane domain at about amino acids 143-162; a potential N-glycosylation site at about amino acids 100-103; and potential N-myristoylation sites at about amino acids 84-89, 103-108, 154-159, and 201-206.

5 An analysis of the Dayhoff database (version 35.45 SwissProt 35), using a WU-BLAST2 sequence alignment analysis of the full-length sequence shown in Figure 210 (SEQ ID NO:358), revealed significant homology between the PRO1575 amino acid sequence and Dayhoff sequence A12005_1. Homology was also revealed between the PRO1575 amino acid sequence and the following additional Dayhoff sequences: P_P80615; P_R25297; P_R51696; A47300; PDI_DROME; P_R49829; P_R63807; DMALPADAP_1; and DRZNF6_1.

10 Clone DNA76401-1683 was deposited with the ATCC on October 20,1998, and is assigned ATCC deposit no. 203360.

EXAMPLE 109: Isolation of cDNA clones Encoding Human PRO1787

15 A consensus DNA sequence was assembled relative to other EST sequences using phrap to form an assembly as described in Example 1 above. This consensus sequence is designated herein "DNA45123". Based on homology of DNA45123 to Incyte EST 3618549 identified in the assembly, as well as other discoveries and information provided herein, the clone including this EST was purchased and sequenced. DNA sequencing of the clone gave the full-length DNA sequence for PRO1787 and the derived protein sequence for PRO1787.

20 The entire coding sequence of PRO1787 is included in Figure 211 (SEQ ID NO:363). Clone DNA76510-2504 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 163-165 and an apparent stop codon at nucleotide positions 970-972 of SEQ ID NO:363. The approximate locations of the signal peptide, transmembrane domain, N-glycosylation sites, N-myristoylation sites and a kinase phosphorylation site are indicated in Figure 212. The predicted polypeptide precursor is 269 amino acids long. Clone DNA76510-2504 has been deposited with the ATCC and is assigned ATCC deposit no. 203477. The full-length PRO1787 protein shown in Figure 212 has an estimated molecular weight of about 29,082 daltons and a pI of about 9.02.

25 An analysis of the Dayhoff database (version 35.45 SwissProt 35), using a WU-BLAST2 sequence alignment analysis of the full-length sequence shown in Figure 212 (SEQ ID NO:364), revealed sequence identity between the PRO1787 amino acid sequence and the following Dayhoff sequences: MYP0_RAT, MYP0_HUMAN, MYP0_BOVIN, GEN12838, HSSCN2B2_1, AF007783_1, HSU90716_1, P_W42015, XLU43330_1 and AF060231_1.

EXAMPLE 110: Isolation of cDNA clones Encoding Human PRO1781

35 Initial DNA sequences referred to herein as DNA58070 and DNA56340 were identified using a yeast screen, in a human SK-Lu-1 adenocarcinoma cell line cDNA library that preferentially represents the 5' ends of the primary cDNA clones. These sequences were clustered and assembled into a consensus DNA sequence using the computer program "phrap" (Phil Green, University of Washington, Seattle, Washington). The consensus sequence is designated herein as "DNA59575".

Based on the DNA59575 consensus sequence, the following oligonucleotides, were synthesized for use

as probes to isolate a clone of the full-length coding sequence for PRO1781 from a human fetal lung cDNA library: TGGAAAAGAAGTCTGGTCAGAAGGTTTAGG (SEQ ID NO:367), CATTGGCTTCATTCTCCTGCTCTG (SEQ ID NO:368), AAAACCTCAGAACAACCTCATTTTGCACC (SEQ ID NO:369) and GTCTACCATGGTTGCTCTTGCCAAATTGTGGGAAGCAGGG (SEQ ID NO:370).

The full length DNA76522-2500 clone shown in Figure 213 contained a single open reading frame with an apparent translational initiation site at nucleotide positions 21 to 23 and ending at the stop codon found at nucleotide positions 1141-1143 (Figure 213; SEQ ID NO:365). The predicted polypeptide precursor (Figure 214, SEQ ID NO:366) is 373 amino acids long. PRO1781 has a calculated molecular weight of approximately 41,221 daltons and an estimated pI of approximately 8.54. Additional features include: a possible signal peptide at about amino acids 1-19; a transmembrane domain at about amino acids 39-60; a tyrosine phosphorylation site at about amino acids 228-236; potential N-myristoylation sites at about amino acids 16-21, 17-22, 43-48, 45-50, 47-52, 49-54, 53-58, 58-63, 59-64, 62-67, 126-131, and 142-147; amidation sites at about amino acids 22-25 and 280-283; and a prokaryotic membrane lipoprotein lipid attachment site at about amino acids 12-22.

An analysis of the Dayhoff database (version 35.45 SwissProt 35), using a WU-BLAST2 sequence alignment analysis of the full-length sequence shown in Figure 214 (SEQ ID NO:366), revealed some homology between the PRO1781 amino acid sequence and the following Dayhoff sequences: CEY4510D_5, AP000001_146, P_R10676, DAC_STRSQ, CEC40H5_5, P_R35204, KPU58495_1, KPN16781_1, AF010403_1, and AF056116_14.

Clone DNA76522-2500 was deposited with the ATCC on November 17, 1998, and is assigned ATCC deposit no. 203469.

EXAMPLE 111: Isolation of cDNA clones Encoding Human PRO1556

Use of the signal sequence algorithm described in Example 3 above allowed identification of an EST cluster sequence from the LIFESEQ® database, designated EST Cluster No. 103158, and also referred to herein as "DNA10398". This EST cluster sequence was then compared to a variety of expressed sequence tag (EST) databases which included public EST databases (e.g., GenBank) and the LIFESEQ® database, to identify existing homologies. The homology search was performed using the computer program BLAST or BLAST2 (Altschul et al., Methods in Enzymology 266:460-480 (1996)). Those comparisons resulting in a BLAST score of 70 (or in some cases 90) or greater that did not encode known proteins were clustered and assembled into a consensus DNA sequence with the program "phrap" (Phil Green, University of Washington, Seattle, Washington). The consensus sequence obtained therefrom is herein designated DNA56417.

In light of the sequence homology between the DNA56417 sequence and a sequence contained within Incyte EST no. 959332, EST no. 959332 was purchased and the cDNA insert was obtained and sequenced. The sequence of this cDNA insert is shown in Figure 215 and is herein designated as DNA76529-1666.

The full length clone shown in Figure 215 contained a single open reading frame with an apparent translational initiation site at nucleotide positions 85 to 87 and ending at the stop codon found at nucleotide positions 892 to 894 (Figure 215; SEQ ID NO:371). The predicted polypeptide precursor (Figure 216, SEQ ID NO:372) is 269 amino acids long. PRO1556 has a calculated molecular weight of approximately 28,004 daltons and an estimated pI of approximately 5.80. Additional features include: a signal peptide sequence at about amino acids 1-24; transmembrane domains at about amino acids 11-25 and 226-243; a potential N-

glycosylation site at about amino acids 182-185, potential cAMP- and cGMP-dependent protein kinase phosphorylation site at about amino acids 70-73; and potential N-myristoylation sites at about amino acids 29-34, 35-39, 117-122, 121-126, 125-130, 154-159, 166-171, 241-246, 246-251, 247-252, 249-254, 250-255, 251-256, 252-257, 253-258, 254-259, 255-260, 256-261, 257-262, and 259-264.

5 An analysis of the Dayhoff database (version 35.45 SwissProt 35), using a WU-BLAST2 sequence alignment analysis of the full-length sequence shown in Figure 216 (SEQ ID NO:372), revealed some homology between the PRO1556 amino acid sequence and the following Dayhoff sequences: T8F5_4, R23B_MOUSE, CANS_HUMAN, P_W41640, DSU51091_1, TP2B_CHICK, DVU20660_1, S43296, P_R23962, and BRN1_HUMAN.

10 Clone DNA76529-1666 was deposited with the ATCC on October 6, 1998, and is assigned ATCC deposit no. 203315.

EXAMPLE 112: Isolation of cDNA clones Encoding Human PRO1759

15 Use of the signal sequence algorithm described in Example 3 above allowed identification of an EST cluster sequence from the Incyte database, designated DNA10571. This EST cluster sequence was then compared to a variety of expressed sequence tag (EST) databases which included public EST databases (e.g., GenBank) and a proprietary EST DNA database (LIFESEQ®, Incyte Pharmaceuticals, Palo Alto, CA) to identify existing homologies. One or more of the ESTs was derived from pooled eosinophils of allergic asthmatic patients. The homology search was performed using the computer program BLAST or BLAST2 (Altshul et al., Methods in Enzymology 266:460-480 (1996)). Those comparisons resulting in a BLAST score of 70 (or in some cases 90) or greater that did not encode known proteins were clustered and assembled into a consensus DNA sequence with the program "phrap" (Phil Green, University of Washington, Seattle, Washington). The consensus sequence obtained therefrom is herein designated DNA57313.

20 In light of the sequence homology between the DNA57313 sequence and the Incyte EST 2434255, the clone including this EST was purchased and the cDNA insert was obtained and sequenced. The sequence of this cDNA insert is shown in Figure 217 and is herein designated as DNA76531-1701.

25 The full length clone shown in Figure 217 contained a single open reading frame with an apparent translational initiation site at nucleotide positions 125-127 and ending at the stop codon found at nucleotide positions 1475-1477 (Figure 217; SEQ ID NO:373). The approximate locations of the signal peptide and transmembrane domains are indicated in Figure 218, whereas the approximate locations for N-myristoylation sites, a lipid attachment site, an amidation site and a kinase phosphorylation site are indicated in Figure 218. The predicted polypeptide precursor (Figure 218, SEQ ID NO:374) is 450 amino acids long. PRO1759 has a calculated molecular weight of approximately 49,765 daltons and an estimated pI of approximately 8.14. Clone DNA76531-1701 was deposited with the ATCC on November 17, 1998 and is assigned ATCC deposit no. 203465.

35 An analysis of the Dayhoff database (version 35.45 SwissProt 35), using a WU-BLAST2 sequence alignment analysis of the full-length sequence shown in Figure 218 (SEQ ID NO:374), revealed sequence identity between the PRO1759 amino acid sequence and the following Dayhoff sequences: OPDE_PSEAE, TH11_TRYBB, S67684, RGT2_YEAST, S68362, ATSUGTRPR_1, P_W17836 (Patent application WO9715668-A2), F69587, A48076, and A45611.

EXAMPLE 113: Isolation of cDNA clones Encoding Human PRO1760

Use of the signal sequence algorithm described in Example 3 above allowed identification of an EST cluster sequence from the Incyte database. This EST cluster sequence was then compared to a variety of expressed sequence tag (EST) databases which included public EST databases (e.g., GenBank) and a proprietary EST DNA database (LIFESEQ®, Incyte Pharmaceuticals, Palo Alto, CA) to identify existing homologies. One or more of the ESTs was derived from a prostate tumor library. The homology search was performed using the computer program BLAST or BLAST2 (Altschul et al., Methods in Enzymology 266:460-480 (1996)). Those comparisons resulting in a BLAST score of 70 (or in some cases 90) or greater that did not encode known proteins were clustered and assembled into a consensus DNA sequence with the program "phrap" (Phil Green, University of Washington, Seattle, Washington). The consensus sequence obtained therefrom is herein designated DNA58798.

In light of the sequence homology between DNA58798 sequence and the Incyte EST 3358745, the clone including this EST was purchased and the cDNA insert was obtained and sequenced. The sequence of this cDNA insert is shown in Figure 219 and is herein designated as DNA76532-1702.

The full length clone shown in Figure 219 contained a single open reading frame with an apparent translational initiation site at nucleotide positions 60-62 and ending at the stop codon found at nucleotide positions 624-626 (Figure 219; SEQ ID NO:375). The predicted polypeptide precursor (Figure 220, SEQ ID NO:376) is 188 amino acids long. Motifs are further indicated in Figure 220. PRO1760 has a calculated molecular weight of approximately 21,042 daltons and an estimated pI of approximately 5.36. Clone DNA76532-1702 was deposited with the ATCC on November 17, 1998 and is assigned ATCC deposit no. 203473.

An analysis of the Dayhoff database (version 35.45 SwissProt 35), using a WU-BLAST2 sequence alignment analysis of the full-length sequence shown in Figure 220 (SEQ ID NO:376), revealed sequence identity between the PRO1760 amino acid sequence and the following Dayhoff sequences: CELT07F12_2, T22J18_16, ATF1C12_3, APE3_YEAST, P_W22471, SAU56908_1, SCPA_STRPY, ATAC00423817, SAPURCLUS_2 and AF041468_9.

EXAMPLE 114: Isolation of cDNA clones Encoding Human PRO1561

A consensus DNA sequence was assembled relative to other EST sequences using phrap and repeated cycles of BLAST and phrap to extend a sequence as far as possible using the EST sequences discussed above as described in Example 1 above. This consensus sequence is herein designated DNA40630. Based on the DNA40630 consensus sequence, oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO1561.

PCR primers (forward and reverse) were synthesized:

forward PCR primer (40630.f1) 5'-CTGCCTCCACTGCTCTGTGCTGGG-3' (SEQ ID NO:379)

reverse PCR primer (40630.r1) 5'-CAGAGCAGTGGATGTTCCCCTGGG-3' (SEQ ID NO:380)

Additionally, a synthetic oligonucleotide hybridization probe was constructed from the consensus DNA40630 sequence which had the following nucleotide sequence

hybridization probe (40630.p1)

5'-CTGAACAAGATGGTCAAGCAAGTGACTGGGAAAATGCCCATCCTC-3' (SEQ ID NO:381)

In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pair identified above. A positive library was then used to isolate clones encoding the PRO1561 gene using the probe oligonucleotide and one of the PCR primers. RNA for construction of the cDNA libraries was isolated from human breast tumor tissue.

DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO1561 (designated herein as DNA76538-1670 [Figure 221, SEQ ID NO:377]; and the derived protein sequence for PRO1561.

The entire nucleotide sequence of DNA76538-1670 is shown in Figure 221 (SEQ ID NO:377). Clone DNA76538-1670 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 29-31 and ending at the stop codon at nucleotide positions 377-379 (Figure 221). The predicted polypeptide precursor is 116 amino acids long (Figure 222). The full-length PRO1561 protein shown in Figure 222 has an estimated molecular weight of about 12,910 daltons and a pI of about 6.41. Analysis of the full-length PRO1561 sequence shown in Figure 222 (SEQ ID NO:378) evidences the presence of the following: a signal peptide from about amino acid 1 to about amino acid 17, a transmembrane domain from about amino acid 1 to about amino acid 24, a potential N-glycosylation site from about amino acid 86 to about amino acid 89, potential N-myristoylation sites from about amino acid 20 to about amino acid 25 and from about amino acid 45 to about amino acid 50 and a phospholipase A2 histidine active site from about amino acid 63 to about amino acid 70. Clone DNA76538-1670 has been deposited with ATCC on October 6, 1998 and is assigned ATCC deposit no. 203313.

An analysis of the Dayhoff database (version 35.45 SwissProt 35), using a WU-BLAST2 sequence alignment analysis of the full-length sequence shown in Figure 222 (SEQ ID NO:378), evidenced significant homology between the PRO1561 amino acid sequence and the following Dayhoff sequences: P_R63053, P_R25416, P_R63055, P_P93363, P_R63046, PA2A_VIPAA, P_W58476, GEN13747, PA2X_HUMAN and PA2A_CRODU.

In addition to the above, a sequence homology search evidenced significant homology between the DNA40630 consensus sequence and Incyte EST clone no. 1921092. As such, Incyte EST clone no. 1921092 was purchased and the insert obtained and sequenced, thereby giving rise to the DNA76538-1670 sequence shown in Figure 221 (SEQ ID NO:377).

EXAMPLE 115: Isolation of cDNA clones Encoding Human PRO1567

A cDNA sequence isolated in the amylase screen described in Example 2 above is herein designated DNA47580. The DNA47580 sequence was then compared to a variety of expressed sequence tag (EST) databases which included public EST databases (e.g., GenBank) and a proprietary EST DNA database (LIFESEQ™, Incyte Pharmaceuticals, Palo Alto, CA) to identify existing homologies. The homology search was performed using the computer program BLAST or BLAST2 (Altschul et al., Methods in Enzymology 266:460-480 (1996)). Those comparisons resulting in a BLAST score of 70 (or in some cases 90) or greater that did not encode known proteins were clustered and assembled into consensus DNA sequences with the program "phrap" (Phil Green, University of Washington, Seattle, Washington). The consensus sequence obtained therefrom is herein designated "DNA57246".

In light of the sequence homology between the DNA57246 sequence and EST no. 1793996 from the

LIFESEQ™ database, the clone containing the EST no. 1793996, which originates from a library constructed from prostate tumor tissue, was purchased and the cDNA insert was obtained and sequenced. The sequence of this cDNA insert is shown in Figure 223 (SEQ ID NO:382) and is herein designated as DNA76541-1675.

A full length clone was identified that contained a single open reading frame with an apparent translational initiation site at nucleotide positions 109-111, and a stop signal at nucleotide positions 643-645 (Fig. 223; SEQ ID NO:382). The predicted polypeptide precursor is 178 amino acids long has a calculated molecular weight of approximately 19,600 daltons and an estimated pI of approximately 5.89. Additional features include a signal peptide at about amino acids 1-22; a potential N-glycosylation site at about amino acids 167-170; a protein kinase C phosphorylation site at about amino acids 107-109; and potential N-myristoylation sites at about amino acids 46-51, 72-77, and 120-125.

An analysis of the Dayhoff database (version 35.45 SwissProt 35), using a WU-BLAST2 sequence alignment analysis of the full-length sequence shown in Figure 224 (SEQ ID NO:383), evidenced significant homology between the PRO1567 amino acid sequence and human colon specific gene CSG6 polypeptide designated Dayhoff sequence "P_W06549". Homology was also found between the PRO1567 amino acid sequence and the following additional Dayhoff sequences: HUAC002301_1, P_246880, A49685, SPBP_RAT, S42924, SPBP_MOUSE, I52115, MMU03711_1, and AF041468_31.

Clone DNA76541-1675 has been deposited with the ATCC on October 27, 1998, and is assigned ATCC deposit no. 203409.

EXAMPLE 116: Isolation of cDNA clones Encoding Human PRO1693

A consensus DNA sequence was assembled relative to other EST sequences using phrap as described in Example 1 above. This consensus sequence is herein designated DNA38251. Based on the DNA38251 consensus sequence, oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO1693.

PCR primers (forward and reverse) were synthesized:

forward PCR primer (38251.f1) 5'-CTGGGATCTGAACAGTTTCGGGGC-3' (SEQ ID NO:386)

reverse PCR primer (38251.r1) 5'-GGTCCCCAGGACATGGTCTGTCCC-3' (SEQ ID NO:387)

Additionally, a synthetic oligonucleotide hybridization probe was constructed from the consensus DNA38251 sequence which had the following nucleotide sequence

hybridization probe (38251.pl)

5'-GCTGAGTTTACATTTACGGTCTAACTCCCTGAGAACCATCCCTGTGCG-3' (SEQ ID NO:388)

In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pair identified above. A positive library was then used to isolate clones encoding the PRO1693 gene using the probe oligonucleotide and one of the PCR primers. RNA for construction of the cDNA libraries was isolated from human fetal kidney tissue.

DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO1693 (designated herein as DNA77301-1708 [Figure 225, SEQ ID NO:384]; and the derived protein sequence for PRO1693.

The entire nucleotide sequence of DNA77301-1708 is shown in Figure 225 (SEQ ID NO:384). Clone

DNA77301-1708 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 508-510 and ending at the stop codon at nucleotide positions 2047-2049 (Figure 225). The predicted polypeptide precursor is 513 amino acids long (Figure 226). The full-length PRO1693 protein shown in Figure 226 has an estimated molecular weight of about 58,266 daltons and a pI of about 9.84. Analysis of the full-length PRO1693 sequence shown in Figure 226 (SEQ ID NO:385) evidences the presence of the following: a signal peptide from about amino acid 1 to about amino acid 33, a transmembrane domain from about amino acid 420 to about amino acid 442, potential N-glycosylation sites from about amino acid 126 to about amino acid 129, from about amino acid 357 to about amino acid 360, from about amino acid 496 to about amino acid 499 and from about amino acid 504 to about amino acid 507, a cAMP- and cGMP-dependent protein kinase phosphorylation site from about amino acid 465 to about amino acid 468, a tyrosine kinase phosphorylation site from about amino acid 136 to about amino acid 142 and potential N-myristylation sites from about amino acid 11 to about amino acid 16, from about amino acid 33 to about amino acid 38, from about amino acid 245 to about amino acid 250, from about amino acid 332 to about amino acid 337, from about amino acid 497 to about amino acid 502 and from about amino acid 507 to about amino acid 512. Clone DNA77301-1708 has been deposited with ATCC on October 27, 1998 and is assigned ATCC deposit no. 203407.

An analysis of the Dayhoff database (version 35.45 SwissProt 35), using a WU-BLAST2 sequence alignment analysis of the full-length sequence shown in Figure 226 (SEQ ID NO:385), evidenced significant homology between the PRO1693 amino acid sequence and the following Dayhoff sequences: AB007876_1, ALS_MOUSE, HSCHON03_1, P_R85889, AF062006_1, AB014462_1, A58532, MUSLRRPA_1, AB007865_1 and AF030435_1.

EXAMPLE 117: Isolation of cDNA clones Encoding Human PRO1784

A cDNA sequence isolated in the amylase screen described in Example 2 above is herein designated DNA43862. Based on the DNA43862 sequence, oligonucleotide probes were generated and used to screen a human fetal kidney library prepared as described in paragraph 1 above. The cloning vector was pRK5B (pRK5B is a precursor of pRK5D that does not contain the SfiI site; see, Holmes et al., *Science*, 253:1278-1280 (1991)), and the cDNA size cut was less than 2800 bp.

PCR primers (forward and reverse) were synthesized:

forward PCR primer (f1) 5'-CTTTTCAGTGTACCTCAGCGATCTC-3' (SEQ ID NO:391); and

reverse PCR primer (r1) 5'-CCAAAACATGGAGCAGGAACAGG-3' (SEQ ID NO:392).

Additionally, a synthetic oligonucleotide hybridization probe was constructed from the DNA43862 sequence which had the following nucleotide sequence:

hybridization probe (p1)

5'-CCAGTTGGTGCTCTCGGACCTACCATGCGAAGAAGATGAAATGTGTG-3' (SEQ ID NO:393).

In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pair identified above. A positive library was then used to isolate clones encoding the PRO1784 gene using the probe oligonucleotide and one of the PCR primers.

A full length clone was identified that contained a single open reading frame with an apparent translational initiation site at nucleotide positions 68-70, and a stop signal at nucleotide positions 506-508 (Fig. 227; SEQ ID NO:389). The predicted polypeptide precursor is 146 amino acids long has a calculated molecular

weight of approximately 16,116 daltons and an estimated pI of approximately 4.99. The approximate locations of the signal peptide, transmembrane domain and N-myristoylation site are indicated in Figure 228. Clone DNA77303-2502 has been deposited with the ATCC and is assigned ATCC deposit no. 203479.

An analysis of the Dayhoff database (version 35.45 SwissProt 35), using a WU-BLAST2 sequence alignment analysis of the full-length sequence shown in Figure 228 (SEQ ID NO:390), evidenced sequence identity between the PRO1784 amino acid sequence and the following Dayhoff sequences: RNU87224_1, RNAF000114_1, P_W31947, S18038, AE001300_8, AF039833_1, P_W39833_1, P_W39788, HSU87223_1, NTU06712_1, and P_W31946.

EXAMPLE 118: Isolation of cDNA clones Encoding Human PRO1605

A cDNA clone (DNA77648-1688) encoding a native human PRO1605 polypeptide was identified by a yeast screen, in a human fetal kidney cDNA library that preferentially represents the 5' ends of the primary cDNA clones.

The full-length DNA77648-1688 clone shown in Figure 229 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 425-427 and ending at the stop codon at nucleotide positions 845-847 (Figure 229). The predicted polypeptide precursor is 140 amino acids long (Figure 230). The full-length PRO1605 protein shown in Figure 230 has an estimated molecular weight of about 15,668 daltons and a pI of about 10.14. Analysis of the full-length PRO1605 sequence shown in Figure 230 (SEQ ID NO:395) evidences the presence of the following: a signal peptide from about amino acid 1 to about amino acid 26. Clone DNA77648-1688 has been deposited with ATCC on October 27, 1998 and is assigned ATCC deposit no. 203408.

An analysis of the Dayhoff database (version 35.45 SwissProt 35), using a WU-BLAST2 sequence alignment analysis of the full-length sequence shown in Figure 230 (SEQ ID NO:395), evidenced significant homology between the PRO1605 amino acid sequence and the following Dayhoff sequences: GNT5_HUMAN, P_R48975, P_W22519, MM26SPROT_1, HSU86782_1, CH60_LEPIN, HMCT_HELPY, F65126, HIU08875_1 and P_R41724.

EXAMPLE 119: Isolation of cDNA clones Encoding Human PRO1788

The extracellular domain (ECD) sequences (including the secretion signal sequence, if any) from about 950 known secreted proteins from the Swiss-Prot public database were used to search EST databases. The EST databases included public EST databases (e.g., GenBank), and a proprietary EST database (LIFESEQ®, Incyte Pharmaceuticals, Palo Alto, CA). The search was performed using the computer program BLAST or BLAST2 [Altschul et al., *Methods in Enzymology*, 266:460-480 (1996)] as a comparison of the ECD protein sequences to a 6 frame translation of the EST sequences. Incyte Clone No. 2968304 was identified as a sequence of interest having a BLAST score of 70 or greater that did not encode known proteins. The nucleotide sequence of Incyte Clone No. 2968304 is designated herein as "DNA6612".

In addition, the DNA6612 sequence was extended using repeated cycles of BLAST and phrap (Phil Green, University of Washington, Seattle, Washington) to extend the sequence as far as possible using the sources of EST sequences discussed above. The extended consensus sequence is designated herein as "DNA49648". Based on the DNA49648 consensus sequence, oligonucleotides were synthesized: 1) to identify

by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO1788.

PCR primers (forward and reverse) were synthesized:

forward PCR primer: CCCTGCCAGCCGAGAGCTTCACC (49648.f1; SEQ ID NO:398)

reverse PCR primer: GGTTGGTGGCCGAAAGGTCCAGC (49648.r1; SEQ ID NO:399)

5 Additionally, a synthetic oligonucleotide hybridization probe was constructed from the consensus DNA49648 sequence which had the following nucleotide sequence:

hybridization probe: CAACCCCAAGCTTAACTGGGCAGGAGCTGAGGTGTTTTTCAGGCC (49648.p1; SEQ ID NO:400)

10 In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pair identified above. A positive library was then used to isolate clones encoding the PRO1788 gene using the probe oligonucleotide and one of the PCR primers. RNA for construction of the cDNA libraries was isolated from human fetal kidney tissue.

15 DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO1788 (designated herein as DNA77652-2505 [Figure 231, SEQ ID NO:396]; and the derived protein sequence for PRO1788.

20 The entire coding sequence of PRO1788 is shown in Figure 231 (SEQ ID NO:396). Clone DNA77652-2505 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 64-66 and an apparent stop codon at nucleotide positions 1123-1125. The predicted polypeptide precursor is 353 amino acids long. The full-length PRO1788 protein shown in Figure 232 has an estimated molecular weight of about 37,847 daltons and a pI of about 6.80. Additional features of PRO1788 include: a signal peptide at about amino acids 1-16; transmembrane domains at about amino acids 215-232 and 287-304; potential N-glycosylation sites at about amino acids 74-77 and 137-140; a glycosaminoglycan attachment site at about amino acids 45-48; a tyrosine kinase phosphorylation site at about amino acids 318-325; N-myristoylation sites at about amino acids 13-18, 32-37, 88-93, 214-219, and 223-228; and a leucine zipper pattern at about amino acids 284-305.

25 An analysis of the Dayhoff database (version 35.45 SwissProt 35), using a WU-BLAST2 sequence alignment analysis of the full-length sequence shown in Figure 232 (SEQ ID NO:397), revealed significant homology between the PRO1788 amino acid sequence and the following Dayhoff sequences: AF030435_1; AF062006_1; DMTARTAN_1; GARP_HUMAN; S42799; P_R71294; HSU88879_1; DROWHEELER_1; A58532; and AF068920_1.

30 Clone DNA77652-2505 was deposited with the ATCC on November 17, 1998, and is assigned ATCC deposit no. 203480.

EXAMPLE 120: Isolation of cDNA clones Encoding Human PRO1801

35 A proprietary expressed sequence tag (EST) DNA database (LIFESEQ®, Incyte Pharmaceuticals, Palo Alto, CA) was searched and an EST was identified which showed homology to the IL-19 protein. This EST sequence is Incyte EST clone no. 819592 and is herein designated DNA79293. Based on the DNA79293 sequence, oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO1801.

PCR primers (forward and reverse) were synthesized:

forward PCR primer 5'-CTCCTGTGGTCTCCAGATTTTCAGGCCTA-3' (SEQ ID NO:403)

reverse PCR primer 5'-AGTCCTCCTTAAGATTCTGATGTCAA-3' (SEQ ID NO:404)

RNA for construction of the cDNA libraries was isolated from human fetal kidney tissue. The cDNA libraries used to isolate the cDNA clones were constructed by standard methods using commercially available reagents such as those from Invitrogen, San Diego, CA. The cDNA was primed with oligo dT containing a NotI site, linked with blunt to SalI hemikinased adaptors, cleaved with NotI, sized appropriately by gel electrophoresis, and cloned in a defined orientation into a suitable cloning vector (such as pRKB or pRKD; pRK5B is a precursor of pRK5D that does not contain the SfiI site; see, Holmes et al., *Science*, 253:1278-1280 (1991)) in the unique XhoI and NotI sites.

DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO1801 (designated herein as DNA83500-2506 [Figure 233, SEQ ID NO:401]; and the derived protein sequence for PRO1801.

The entire nucleotide sequence of DNA83500-2506 is shown in Figure 233 (SEQ ID NO:401). Clone DNA83500-2506 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 109-111 and ending at the stop codon at nucleotide positions 892-894 (Figure 233). The predicted polypeptide precursor is 261 amino acids long (Figure 234). The full-length PRO1801 protein shown in Figure 234 has an estimated molecular weight of about 29,667 daltons and a pI of about 8.76. Analysis of the full-length PRO1801 sequence shown in Figure 234 (SEQ ID NO:402) evidences the presence of the following: a signal peptide from about amino acid 1 to about amino acid 42, cAMP- and cGMP-dependent protein kinase phosphorylation sites from about amino acid 192 to about amino acid 195 and from about amino acid 225 to about amino acid 228 and potential N-myristoylation sites from about amino acid 42 to about amino acid 47, from about amino acid 46 to about amino acid 51 and from about amino acid 136 to about amino acid 141. Clone DNA83500-2506 has been deposited with ATCC on October 29, 1998 and is assigned ATCC deposit no. 203391.

An analysis of the Dayhoff database (version 35.45 SwissProt 35), using a WU-BLAST2 sequence alignment analysis of the full-length sequence shown in Figure 234 (SEQ ID NO:402), evidenced significant homology between the PRO1801 amino acid sequence and the following Dayhoff sequences: P_W37935, HGS_B477, P_R32277, IL10_MACFA, P_W46585, P_R39714, P_R71471, P_R10159, IL10_RAT and P_W57201.

EXAMPLE 121: Isolation of cDNA clones Encoding Human UCP4

EST databases, which included public EST databases (e.g., GenBank), and a proprietary EST database (LIFESEQ™, Incyte Pharmaceuticals, Palo Alto, CA), were searched for sequences having homologies to human UCP3. The search was performed using the computer program BLAST or BLAST2 [Altschul et al., *Methods in Enzymology*, 266:460-480 (1996)] as a comparison of the UCP3 protein sequences to a 6 frame translation of the EST sequences. Those comparisons resulting in a BLAST score of 70 (or in some cases, 90) or greater that did not encode known proteins were clustered and assembled into consensus DNA sequences with the program AssemblIGN and MacVector (Oxford Molecular Group, Inc.).

A DNA sequence ("fromDNA") was assembled relative to other EST sequences using AssemblIGN software. In addition, the fromDNA sequence was extended using repeated cycles of BLAST and AssemblIGN

to extend the sequence as far as possible using the sources of EST sequences discussed above. Based on this DNA sequence, oligonucleotides were synthesized to isolate a clone of the full-length coding sequences for UCP4 by PCR. Forward and reverse PCR primers generally range from 20 to 30 nucleotides and are often designed to give a PCR product of about 100-1000 bp in length. The probe sequences are typically 40-55 bp in length. In some cases, additional oligonucleotides are synthesized when the consensus sequence is greater than about 1-1.5kbp.

PCR primers (forward and reverse) were synthesized:

forward PCR primer CGCGGATCCCGTTATCGTCTTGCGCTACTGC (SEQ ID NO:407)

reverse PCR primer GCGGAATTCTTAAATGGACTGACTCCACTCATC (SEQ ID NO:408)

RNA for construction of the cDNA libraries was isolated from brain tissue. The cDNA libraries used to isolated the cDNA clones were constructed by standard methods using commercially available reagents such as those from Invitrogen, San Diego, CA. The cDNA was primed with oligo dT containing a NotI site, linked with blunt to SalI hemikinased adaptors, cleaved with NotI, sized appropriately by gel electrophoresis, and cloned in a defined orientation into a suitable cloning vector (such as pRKB or pRKD; pRK5B is a precursor of pRK5D that does not contain the SfiI site; see, Holmes et al., *Science*, 253:1278-1280 (1991)) in the unique XhoI and NotI sites.

DNA sequencing of the clone isolated by PCR as described above gave the full-length DNA sequence for UCP4 (designated herein as DNA77568-1626 [Figure 235, SEQ ID NO:405] and the derived protein sequence for UCP4.

The entire coding sequence of UCP4 is shown in Figure 235 (SEQ ID NO:405). Clone DNA77568-1626 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 27-29, and an apparent stop codon at nucleotide positions 996-998. (See Figure 235; SEQ ID NO:405). The predicted polypeptide precursor is 323 amino acids long. It is presently believed that UCP4 is a membrane-bound protein and contains at least 6 transmembrane regions. These putative transmembrane regions in the UCP4 amino acid sequence are illustrated in Figure 236. Clone DNA77568-1626, contained in the pcDNA3 vector (Invitrogen) has been deposited with ATCC and is assigned ATCC deposit no. 203134. UCP4 polypeptide is obtained or obtainable by expressing the molecule encoded by the cDNA insert of the deposited ATCC 203134 vector. Digestion of the vector with BamHI and EcoRI restriction enzymes will yield an approximate 972 plus 34 bp insert. The full-length UCP4 protein shown in Figure 236 has an estimated molecular weight of about 36,061 daltons and a pI of about 9.28.

EXAMPLE 122: Isolation of cDNA clones Encoding Human PRO193

A consensus DNA sequence was assembled relative to other EST sequences using phrap as described in Example 1 above. Based on this consensus sequence, oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO193.

A pair of PCR primers (forward and reverse) were synthesized:

forward PCR primer 5'-GTTTGAGGAAGCTGGGATAC-3' (SEQ ID NO:411); and

reverse PCR primer 5'-CCAAACTCGAGCACCTGTTC-3' (SEQ ID NO:412).

Additionally, a synthetic oligonucleotide hybridization probe was constructed from the consensus sequence which

had the following nucleotide sequence:

hybridization probe

5'-ATGGCAGGCTTCCTAGATAATTTTCGTTGGCCAGAATGTG-3' (SEQ ID NO:413).

5 In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pair identified above. A positive library was then used to isolate clones encoding the PRO193 gene using the probe oligonucleotide and one of the PCR primers. RNA for construction of the cDNA libraries was isolated from human retina tissue (LIB94).

10 DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO193 [herein designated as DNA23322-1393] (SEQ ID NO:409) and the derived protein sequence for PRO193.

15 The entire nucleotide sequence of DNA23322-1393 is shown in Figure 237 (SEQ ID NO:409). Clone DNA23322-1393 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 138-140 and ending at the stop codon at nucleotide positions 612-614 (Figure 237). The predicted polypeptide precursor is 158 amino acids long (Figure 238). The full-length PRO193 protein shown in Figure 238 has an estimated molecular weight of about 17,936 and a pI of about 5.32. Clone DNA23322-1393 has been deposited with the ATCC. Regarding the sequence, it is understood that the deposited clone contains the correct sequence, and the sequences provided herein are based on known sequencing techniques.

20 Still analyzing the amino acid sequence of SEQ ID NO:410, transmembrane domains are at about amino acids 23-42, 60-80, 97-117 and 128-148 of SEQ ID NO:410. A cell attachment sequence is at about amino acids 81-83 of SEQ ID NO:410. A peroxidase proximal heme-ligand domain is at about amino acids 81-83 of SEQ ID NO:410. The corresponding nucleotides can be routinely determined given the sequences provided herein.

EXAMPLE 123: Isolation of cDNA clones Encoding Human PRO1130

25 A consensus DNA sequence was assembled relative to other EST sequences using phrap as described in Example 1 above. This consensus sequence is herein designated DNA34360. Based on the DNA34360 consensus sequence, oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO1130.

PCR primers (forward and reverse) were synthesized:

30 forward PCR primer (34360.f1) 5'-GCCATAGTCACGACATGGATG-3' (SEQ ID NO:416)
forward PCR primer (34360.f2) 5'-GGATGGCCAGAGCTGCTG-3' (SEQ ID NO:417)
forward PCR primer (34360.f3) 5'-AAAGTACAAGTGTGGCCTCATCAAGC-3' (SEQ ID NO:418)
reverse PCR primer (34360.r1) 5'-TCTGACTCCTAAGTCAGGCAGGAG-3' (SEQ ID NO:419)
reverse PCR primer (34360.r2) 5'-ATTCTCTCCACAGACAGCTGGTTC-3' (SEQ ID NO:420)

35 Additionally, a synthetic oligonucleotide hybridization probe was constructed from the consensus DNA34360 sequence which had the following nucleotide sequence

hybridization probe (34360.p1)

5'-GTACAAGTGTGGCCTCATCAAGCCCTGCCCAGCCAACTACTTTGCG-3' (SEQ ID NO:421)

In order to screen several libraries for a source of a full-length clone, DNA from the libraries was

screened by PCR amplification with the PCR primer pair identified above. A positive library was then used to isolate clones encoding the PRO1130 gene using the probe oligonucleotide and one of the PCR primers. RNA for construction of the cDNA libraries was isolated from human aortic endothelial cell tissue.

DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO1130 (designated herein as DNA59814-1486 [Figure 239, SEQ ID NO:414]; and the derived protein sequence for PRO1130.

The entire nucleotide sequence of DNA59814-1486 is shown in Figure 239 (SEQ ID NO:414). Clone DNA59814-1486 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 312-314 and ending at the stop codon at nucleotide positions 984-986 (Figure 239). The predicted polypeptide precursor is 224 amino acids long (Figure 240). The full-length PRO1130 protein shown in Figure 240 has an estimated molecular weight of about 24,963 daltons and a pI of about 9.64. Analysis of the full-length PRO1130 sequence shown in Figure 240 (SEQ ID NO:415) evidences the presence of the following: a signal peptide from about amino acid 1 to about amino acid 15, an ATP/GTP-binding site motif A from about amino acid 184 to about amino acid 191 and a potential N-glycosylation site from about amino acid 107 to about amino acid 110. Clone DNA59814-1486 has been deposited with ATCC on October 20, 1998 and is assigned ATCC deposit no. 203359.

An analysis of the Dayhoff database (version 35.45 SwissProt 35), using a WU-BLAST2 sequence alignment analysis of the full-length sequence shown in Figure 240 (SEQ ID NO:415), evidenced significant homology between the PRO1130 amino acid sequence and the following Dayhoff sequences: P_W06547, 216_HUMAN, D87120_1, MMU72677_1, LAU04889_1, and D69319.

EXAMPLE 124: Isolation of cDNA clones Encoding Human PRO1335

A consensus DNA sequence was assembled relative to other EST sequences using phrap as described in Example 1 above. This consensus sequence is herein designated DNA35727. Based on the DNA35727 consensus sequence, oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO1335.

PCR primers (forward and reverse) were synthesized:

forward PCR primer (35727.f1) 5'-GTAAAGTCGCTGGCCAGC-3' (SEQ ID NO:424)

forward PCR primer (35727.f2) 5'-CCCGATCTGCCTGCTGTA-3' (SEQ ID NO:425)

reverse PCR primer (35727.r1) 5'-CTGCACTGTATGGCCATTATTGTG-3' (SEQ ID NO:426)

Additionally, a synthetic oligonucleotide hybridization probe was constructed from the consensus DNA35727 sequence which had the following nucleotide sequence

hybridization probe (35727.p1)

5'-CAGAAACCCATGATACCCTACTGAACACCGAATCCCCTGGAAGCC-3' (SEQ ID NO:427)

In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pair identified above. A positive library was then used to isolate clones encoding the PRO1335 gene using the probe oligonucleotide and one of the PCR primers. RNA for construction of the cDNA libraries was isolated from human retina tissue.

DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO1335 (designated herein as DNA62812-1594 [Figure 241, SEQ ID NO:422]; and the derived protein sequence for PRO1335.

The entire nucleotide sequence of DNA62812-1594 is shown in Figure 241 (SEQ ID NO:422). Clone DNA62812-1594 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 271-273 and ending at the stop codon at nucleotide positions 1282-1284 (Figure 241). The predicted polypeptide precursor is 337 amino acids long (Figure 242). The full-length PRO1335 protein shown in Figure 242 has an estimated molecular weight of about 37,668 daltons and a pI of about 6.27. Analysis of the full-length PRO1335 sequence shown in Figure 242 (SEQ ID NO:423) evidences the presence of the following: a signal peptide from about amino acid 1 to about amino acid 15, a transmembrane domain from about amino acid 291 to about amino acid 310, a potential N-glycosylation site from about amino acid 213 to about amino acid 216 and amino acid sequence blocks having homology to eukaryotic-type carbonic anhydrase proteins from about amino acid 197 to about amino acid 245, from about amino acid 104 to about amino acid 140 and from about amino acid 22 to about amino acid 69. Clone DNA62812-1594 has been deposited with ATCC on September 9, 1998 and is assigned ATCC deposit no. 203248.

An analysis of the Dayhoff database (version 35.45 SwissProt 35), using a WU-BLAST2 sequence alignment analysis of the full-length sequence shown in Figure 242 (SEQ ID NO:423), evidenced significant homology between the PRO1335 amino acid sequence and the following Dayhoff sequences: AF037335_1, I38013, PTPG_MOUSE, CAH2_HUMAN, ICAC, CAH7_HUMAN, CAH3_HUMAN, CAH1_HUMAN, CAH5_HUMAN and P_R41746.

EXAMPLE 125: Isolation of cDNA clones Encoding Human PRO1329

Use of the signal sequence algorithm described in Example 3 above allowed identification of an EST cluster sequence from the LIFESEQ® database, designated Incyte Cluster No. 167544, also referred herein as "DNA10680". This EST cluster sequence was then compared to a variety of expressed sequence tag (EST) databases which included public EST databases (e.g., GenBank) and a proprietary EST DNA database (LIFESEQ®, Incyte Pharmaceuticals, Palo Alto, CA) to identify existing homologies. The homology search was performed using the computer program BLAST or BLAST2 (Altschul et al., Methods in Enzymology 266:460-480 (1996)). Those comparisons resulting in a BLAST score of 70 (or in some cases 90) or greater that did not encode known proteins were clustered and assembled into a consensus DNA sequence with the program "phrap" (Phil Green, University of Washington, Seattle, Washington). One or more of the ESTs was derived from a cDNA library constructed from RNA isolated from synovial membrane tissue removed from the elbow of a female with rheumatoid arthritis. The consensus sequence obtained therefrom is herein designated "DNA58836".

In light of the sequence homology between the DNA58836 sequence and a sequence contained within the Incyte EST clone no. 368774, EST clone 368774 was purchased and the cDNA insert was obtained and sequenced. The sequence of this cDNA insert is shown in Figure 243 and is herein designated as DNA66660-1585.

The full length clone shown in Figure 243 contained a single open reading frame with an apparent

translational initiation site at nucleotide positions 90 to 92 and ending at the stop codon found at nucleotide positions 717 to 719 (Figure 243; SEQ ID NO:428). The predicted polypeptide precursor (Figure 244, SEQ ID NO:429) is 209 amino acids long, with a signal sequence at about amino acids 1-16. PRO1329 has a calculated molecular weight of approximately 21,588 daltons and an estimated pI of approximately 5.50. Clone DNA66660-1585 was deposited with the ATCC on September 22, 1998 and is assigned ATCC deposit no. 203279.

An analysis of the Dayhoff database (version 35.45 SwissProt 35), using a WU-BLAST2 sequence alignment analysis of the full-length sequence shown in Figure 244 (SEQ ID NO:429), revealed some homology between the PRO1329 amino acid sequence and the following Dayhoff sequences: CELK06A9_3, PROA_XANCP, CXU21300_4, MTV037_17, SYN1_RAT, I56542, S60743, BNOLE3_1, AB001573_1, and P_P80671.

EXAMPLE 126: Isolation of cDNA clones Encoding Human PRO1550

Use of the signal sequence algorithm described in Example 3 above allowed identification of an EST sequence from the Merck database, designated CELT15B7_12, also referred herein as "DNA10022". This EST sequence was then compared to a variety of expressed sequence tag (EST) databases which included public and proprietary EST databases (e.g., GenBank and LIFESEQ®) to identify existing homologies. The homology search was performed using the computer program BLAST or BLAST2 (Altschul et al., Methods in Enzymology 266:460-480 (1996)). Those comparisons resulting in a BLAST score of 70 (or in some cases 90) or greater that did not encode known proteins were clustered and assembled into a consensus DNA sequence with the program "phrap" (Phil Green, University of Washington, Seattle, Washington). The consensus sequence obtained therefrom is herein designated "DNA55708".

In light of the sequence homology between the DNA55708 sequence and a sequence contained within Incyte EST no. 3411659, the EST clone 3411659 was purchased and the cDNA insert was obtained and sequenced in its entirety. The sequence of this cDNA insert is shown in Figure 245 and is herein designated as "DNA76393-1664".

The full length clone shown in Figure 245 contained a single open reading frame with an apparent translational initiation site at nucleotide positions 138 to 140 and ending at the stop codon found at nucleotide positions 867 to 869 (Figure 245; SEQ ID NO:430). The predicted polypeptide precursor (Figure 246, SEQ ID NO:431) is 243 amino acids long. Other features of the PRO1550 protein include: a signal sequence at about amino acids 1-30; a hydrophobic domain at about amino acids 195-217; and a potential N-glycosylation site at about amino acids 186-189. PRO1550 has a calculated molecular weight of approximately 26,266 daltons and an estimated pI of approximately 8.43. Clone DNA76393-1664 was deposited with the ATCC on October 6, 1998, and is assigned ATCC deposit no. 203323.

An analysis of the Dayhoff database (version 35.45 SwissProt 35), using a WU-BLAST2 sequence alignment analysis of the full-length sequence shown in Figure 246 (SEQ ID NO:431), revealed some homology between the PRO1550 amino acid sequence and the following Dayhoff sequences: CELF59E12_11; CA24_ASCSU; AF018082_1; CA13_BOVIN; CA54_HUMAN; CA34_HUMAN; HUMCOL7A1X_1; P_W09643; AF053538_1; and HSEMCMXIV2_1.

EXAMPLE 127: Use of PRO as a hybridization probe

The following method describes use of a nucleotide sequence encoding PRO as a hybridization probe.

DNA comprising the coding sequence of full-length or mature PRO as disclosed herein is employed as a probe to screen for homologous DNAs (such as those encoding naturally-occurring variants of PRO) in human tissue cDNA libraries or human tissue genomic libraries.

Hybridization and washing of filters containing either library DNAs is performed under the following high stringency conditions. Hybridization of radiolabeled PRO-derived probe to the filters is performed in a solution of 50% formamide, 5x SSC, 0.1% SDS, 0.1% sodium pyrophosphate, 50 mM sodium phosphate, pH 6.8, 2x Denhardt's solution, and 10% dextran sulfate at 42°C for 20 hours. Washing of the filters is performed in an aqueous solution of 0.1x SSC and 0.1% SDS at 42°C.

DNAs having a desired sequence identity with the DNA encoding full-length native sequence PRO can then be identified using standard techniques known in the art.

EXAMPLE 128: Expression of PRO in *E. coli*

This example illustrates preparation of an unglycosylated form of PRO by recombinant expression in *E. coli*.

The DNA sequence encoding PRO is initially amplified using selected PCR primers. The primers should contain restriction enzyme sites which correspond to the restriction enzyme sites on the selected expression vector. A variety of expression vectors may be employed. An example of a suitable vector is pBR322 (derived from *E. coli*; see Bolivar et al., *Gene*, 2:95 (1977)) which contains genes for ampicillin and tetracycline resistance. The vector is digested with restriction enzyme and dephosphorylated. The PCR amplified sequences are then ligated into the vector. The vector will preferably include sequences which encode for an antibiotic resistance gene, a trp promoter, a polyhis leader (including the first six STII codons, polyhis sequence, and enterokinase cleavage site), the PRO coding region, lambda transcriptional terminator, and an argU gene.

The ligation mixture is then used to transform a selected *E. coli* strain using the methods described in Sambrook et al., *supra*. Transformants are identified by their ability to grow on LB plates and antibiotic resistant colonies are then selected. Plasmid DNA can be isolated and confirmed by restriction analysis and DNA sequencing.

Selected clones can be grown overnight in liquid culture medium such as LB broth supplemented with antibiotics. The overnight culture may subsequently be used to inoculate a larger scale culture. The cells are then grown to a desired optical density, during which the expression promoter is turned on.

After culturing the cells for several more hours, the cells can be harvested by centrifugation. The cell pellet obtained by the centrifugation can be solubilized using various agents known in the art, and the solubilized PRO protein can then be purified using a metal chelating column under conditions that allow tight binding of the protein.

PRO may be expressed in *E. coli* in a poly-His tagged form, using the following procedure. The DNA encoding PRO is initially amplified using selected PCR primers. The primers will contain restriction enzyme sites which correspond to the restriction enzyme sites on the selected expression vector, and other useful

sequences providing for efficient and reliable translation initiation, rapid purification on a metal chelation column, and proteolytic removal with enterokinase. The PCR-amplified, poly-His tagged sequences are then ligated into an expression vector, which is used to transform an *E. coli* host based on strain 52 (W3110 fuhA(tonA) lon galE rpoHts(htpRts) clpP(lacIq). Transformants are first grown in LB containing 50 mg/ml carbenicillin at 30°C with shaking until an O.D.₆₀₀ of 3-5 is reached. Cultures are then diluted 50-100 fold into

5 CRAP media (prepared by mixing 3.57 g (NH₄)₂SO₄, 0.71 g sodium citrate•2H₂O, 1.07 g KCl, 5.36 g Difco yeast extract, 5.36 g Sheffield hycase SF in 500 mL water, as well as 110 mM MPOS, pH 7.3, 0.55% (w/v) glucose and 7 mM MgSO₄) and grown for approximately 20-30 hours at 30°C with shaking. Samples are removed to verify expression by SDS-PAGE analysis, and the bulk culture is centrifuged to pellet the cells. Cell pellets are frozen until purification and refolding.

10 *E. coli* paste from 0.5 to 1 L fermentations (6-10 g pellets) is resuspended in 10 volumes (w/v) in 7 M guanidine, 20 mM Tris, pH 8 buffer. Solid sodium sulfite and sodium tetrathionate is added to make final concentrations of 0.1M and 0.02 M, respectively, and the solution is stirred overnight at 4°C. This step results in a denatured protein with all cysteine residues blocked by sulfitolization. The solution is centrifuged at 40,000 rpm in a Beckman Ultracentrifuge for 30 min. The supernatant is diluted with 3-5 volumes of metal chelate

15 column buffer (6 M guanidine, 20 mM Tris, pH 7.4) and filtered through 0.22 micron filters to clarify. The clarified extract is loaded onto a 5 ml Qiagen Ni-NTA metal chelate column equilibrated in the metal chelate column buffer. The column is washed with additional buffer containing 50 mM imidazole (Calbiochem, Utrol grade), pH 7.4. The protein is eluted with buffer containing 250 mM imidazole. Fractions containing the desired protein are pooled and stored at 4°C. Protein concentration is estimated by its absorbance at 280 nm

20 using the calculated extinction coefficient based on its amino acid sequence.

The proteins are refolded by diluting the sample slowly into freshly prepared refolding buffer consisting of: 20 mM Tris, pH 8.6, 0.3 M NaCl, 2.5 M urea, 5 mM cysteine, 20 mM glycine and 1 mM EDTA. Refolding volumes are chosen so that the final protein concentration is between 50 to 100 micrograms/ml. The refolding solution is stirred gently at 4°C for 12-36 hours. The refolding reaction is quenched by the addition

25 of TFA to a final concentration of 0.4% (pH of approximately 3). Before further purification of the protein, the solution is filtered through a 0.22 micron filter and acetonitrile is added to 2-10% final concentration. The refolded protein is chromatographed on a Poros R1/H reversed phase column using a mobile buffer of 0.1% TFA with elution with a gradient of acetonitrile from 10 to 80%. Aliquots of fractions with A₂₈₀ absorbance are analyzed on SDS polyacrylamide gels and fractions containing homogeneous refolded protein are pooled.

30 Generally, the properly refolded species of most proteins are eluted at the lowest concentrations of acetonitrile since those species are the most compact with their hydrophobic interiors shielded from interaction with the reversed phase resin. Aggregated species are usually eluted at higher acetonitrile concentrations. In addition to resolving misfolded forms of proteins from the desired form, the reversed phase step also removes endotoxin from the samples.

35 Fractions containing the desired folded PRO polypeptide are pooled and the acetonitrile removed using a gentle stream of nitrogen directed at the solution. Proteins are formulated into 20 mM Hepes, pH 6.8 with 0.14 M sodium chloride and 4% mannitol by dialysis or by gel filtration using G25 Superfine (Pharmacia) resins equilibrated in the formulation buffer and sterile filtered.

Many of the PRO polypeptides disclosed herein were successfully expressed as described above.

EXAMPLE 129: Expression of PRO in mammalian cells

This example illustrates preparation of a potentially glycosylated form of PRO by recombinant expression in mammalian cells.

5 The vector, pRK5 (see EP 307,247, published March 15, 1989), is employed as the expression vector. Optionally, the PRO DNA is ligated into pRK5 with selected restriction enzymes to allow insertion of the PRO DNA using ligation methods such as described in Sambrook et al., *supra*. The resulting vector is called pRK5-PRO.

10 In one embodiment, the selected host cells may be 293 cells. Human 293 cells (ATCC CCL 1573) are grown to confluence in tissue culture plates in medium such as DMEM supplemented with fetal calf serum and optionally, nutrient components and/or antibiotics. About 10 μ g pRK5-PRO DNA is mixed with about 1 μ g DNA encoding the VA RNA gene [Thimmappaya et al., *Cell*, 31:543 (1982)] and dissolved in 500 μ l of 1 mM Tris-HCl, 0.1 mM EDTA, 0.227 M CaCl_2 . To this mixture is added, dropwise, 500 μ l of 50 mM HEPES (pH 7.35), 280 mM NaCl, 1.5 mM NaPO_4 , and a precipitate is allowed to form for 10 minutes at 25°C. The precipitate is suspended and added to the 293 cells and allowed to settle for about four hours at 37°C. The culture medium is aspirated off and 2 ml of 20% glycerol in PBS is added for 30 seconds. The 293 cells are then washed with serum free medium, fresh medium is added and the cells are incubated for about 5 days.

15 Approximately 24 hours after the transfections, the culture medium is removed and replaced with culture medium (alone) or culture medium containing 200 $\mu\text{Ci/ml}$ ^{35}S -cysteine and 200 $\mu\text{Ci/ml}$ ^{35}S -methionine. After 20 a 12 hour incubation, the conditioned medium is collected, concentrated on a spin filter, and loaded onto a 15% SDS gel. The processed gel may be dried and exposed to film for a selected period of time to reveal the presence of PRO polypeptide. The cultures containing transfected cells may undergo further incubation (in serum free medium) and the medium is tested in selected bioassays.

25 In an alternative technique, PRO may be introduced into 293 cells transiently using the dextran sulfate method described by Sompariyac et al., *Proc. Natl. Acad. Sci.*, 12:7575 (1981). 293 cells are grown to maximal density in a spinner flask and 700 μ g pRK5-PRO DNA is added. The cells are first concentrated from the spinner flask by centrifugation and washed with PBS. The DNA-dextran precipitate is incubated on the cell pellet for four hours. The cells are treated with 20% glycerol for 90 seconds, washed with tissue culture medium, and re-introduced into the spinner flask containing tissue culture medium, 5 $\mu\text{g/ml}$ bovine insulin and 30 0.1 $\mu\text{g/ml}$ bovine transferrin. After about four days, the conditioned media is centrifuged and filtered to remove cells and debris. The sample containing expressed PRO can then be concentrated and purified by any selected method, such as dialysis and/or column chromatography.

35 In another embodiment, PRO can be expressed in CHO cells. The pRK5-PRO can be transfected into CHO cells using known reagents such as CaPO_4 or DEAE-dextran. As described above, the cell cultures can be incubated, and the medium replaced with culture medium (alone) or medium containing a radiolabel such as ^{35}S -methionine. After determining the presence of PRO polypeptide, the culture medium may be replaced with serum free medium. Preferably, the cultures are incubated for about 6 days, and then the conditioned medium is harvested. The medium containing the expressed PRO can then be concentrated and purified by any selected method.

Epitope-tagged PRO may also be expressed in host CHO cells. The PRO may be subcloned out of the pRK5 vector. The subclone insert can undergo PCR to fuse in frame with a selected epitope tag such as a poly-his tag into a Baculovirus expression vector. The poly-his tagged PRO insert can then be subcloned into a SV40 driven vector containing a selection marker such as DHFR for selection of stable clones. Finally, the CHO cells can be transfected (as described above) with the SV40 driven vector. Labeling may be performed, as described above, to verify expression. The culture medium containing the expressed poly-His tagged PRO can then be concentrated and purified by any selected method, such as by Ni^{2+} -chelate affinity chromatography.

PRO may also be expressed in CHO and/or COS cells by a transient expression procedure or in CHO cells by another stable expression procedure.

Stable expression in CHO cells is performed using the following procedure. The proteins are expressed as an IgG construct (immunoadhesin), in which the coding sequences for the soluble forms (e.g. extracellular domains) of the respective proteins are fused to an IgG1 constant region sequence containing the hinge, CH2 and CH2 domains and/or is a poly-His tagged form.

Following PCR amplification, the respective DNAs are subcloned in a CHO expression vector using standard techniques as described in Ausubel et al., Current Protocols of Molecular Biology, Unit 3.16, John Wiley and Sons (1997). CHO expression vectors are constructed to have compatible restriction sites 5' and 3' of the DNA of interest to allow the convenient shuttling of cDNA's. The vector used expression in CHO cells is as described in Lucas et al., Nucl. Acids Res. 24:9 (1774-1779 (1996), and uses the SV40 early promoter/enhancer to drive expression of the cDNA of interest and dihydrofolate reductase (DHFR). DHFR expression permits selection for stable maintenance of the plasmid following transfection.

Twelve micrograms of the desired plasmid DNA is introduced into approximately 10 million CHO cells using commercially available transfection reagents Superfect® (Quiagen), Dosper® or Fugene® (Boehringer Mannheim). The cells are grown as described in Lucas et al., supra. Approximately 3×10^7 cells are frozen in an ampule for further growth and production as described below.

The ampules containing the plasmid DNA are thawed by placement into water bath and mixed by vortexing. The contents are pipetted into a centrifuge tube containing 10 mLs of media and centrifuged at 1000 rpm for 5 minutes. The supernatant is aspirated and the cells are resuspended in 10 mL of selective media (0.2 μm filtered PS20 with 5% 0.2 μm diafiltered fetal bovine serum). The cells are then aliquoted into a 100 mL spinner containing 90 mL of selective media. After 1-2 days, the cells are transferred into a 250 mL spinner filled with 150 mL selective growth medium and incubated at 37°C. After another 2-3 days, 250 mL, 500 mL and 2000 mL spinners are seeded with 3×10^5 cells/mL. The cell media is exchanged with fresh media by centrifugation and resuspension in production medium. Although any suitable CHO media may be employed, a production medium described in U.S. Patent No. 5,122,469, issued June 16, 1992 may actually be used. A 3L production spinner is seeded at 1.2×10^6 cells/mL. On day 0, the cell number pH is determined. On day 1, the spinner is sampled and sparging with filtered air is commenced. On day 2, the spinner is sampled, the temperature shifted to 33°C, and 30 mL of 500 g/L glucose and 0.6 mL of 10% antifoam (e.g., 35% polydimethylsiloxane emulsion, Dow Corning 365 Medical Grade Emulsion) taken. Throughout the production, the pH is adjusted as necessary to keep it at around 7.2. After 10 days, or until the viability dropped below 70%, the cell culture is harvested by centrifugation and filtering through a 0.22 μm filter. The filtrate was either stored at 4°C or immediately loaded onto columns for purification.

For the poly-His tagged constructs, the proteins are purified using a Ni-NTA column (Qiagen). Before purification, imidazole is added to the conditioned media to a concentration of 5 mM. The conditioned media is pumped onto a 6 ml Ni-NTA column equilibrated in 20 mM Hepes, pH 7.4, buffer containing 0.3 M NaCl and 5 mM imidazole at a flow rate of 4-5 ml/min. at 4°C. After loading, the column is washed with additional equilibration buffer and the protein eluted with equilibration buffer containing 0.25 M imidazole. The highly purified protein is subsequently desalted into a storage buffer containing 10 mM Hepes, 0.14 M NaCl and 4% mannitol, pH 6.8, with a 25 ml G25 Superfine (Pharmacia) column and stored at -80°C.

Immunoadhesin (Fc-containing) constructs are purified from the conditioned media as follows. The conditioned medium is pumped onto a 5 ml Protein A column (Pharmacia) which had been equilibrated in 20 mM Na phosphate buffer, pH 6.8. After loading, the column is washed extensively with equilibration buffer before elution with 100 mM citric acid, pH 3.5. The eluted protein is immediately neutralized by collecting 1 ml fractions into tubes containing 275 μ L of 1 M Tris buffer, pH 9. The highly purified protein is subsequently desalted into storage buffer as described above for the poly-His tagged proteins. The homogeneity is assessed by SDS polyacrylamide gels and by N-terminal amino acid sequencing by Edman degradation.

Many of the PRO polypeptides disclosed herein were successfully expressed as described above.

EXAMPLE 130: Expression of PRO in Yeast

The following method describes recombinant expression of PRO in yeast.

First, yeast expression vectors are constructed for intracellular production or secretion of PRO from the ADH2/GAPDH promoter. DNA encoding PRO and the promoter is inserted into suitable restriction enzyme sites in the selected plasmid to direct intracellular expression of PRO. For secretion, DNA encoding PRO can be cloned into the selected plasmid, together with DNA encoding the ADH2/GAPDH promoter, a native PRO signal peptide or other mammalian signal peptide, or, for example, a yeast alpha-factor or invertase secretory signal/leader sequence, and linker sequences (if needed) for expression of PRO.

Yeast cells, such as yeast strain AB110, can then be transformed with the expression plasmids described above and cultured in selected fermentation media. The transformed yeast supernatants can be analyzed by precipitation with 10% trichloroacetic acid and separation by SDS-PAGE, followed by staining of the gels with Coomassie Blue stain.

Recombinant PRO can subsequently be isolated and purified by removing the yeast cells from the fermentation medium by centrifugation and then concentrating the medium using selected cartridge filters. The concentrate containing PRO may further be purified using selected column chromatography resins.

Many of the PRO polypeptides disclosed herein were successfully expressed as described above.

EXAMPLE 131: Expression of PRO in Baculovirus-Infected Insect Cells

The following method describes recombinant expression of PRO in Baculovirus-infected insect cells.

The sequence coding for PRO is fused upstream of an epitope tag contained within a baculovirus expression vector. Such epitope tags include poly-his tags and immunoglobulin tags (like Fc regions of IgG). A variety of plasmids may be employed, including plasmids derived from commercially available plasmids such as pVL1393 (Novagen). Briefly, the sequence encoding PRO or the desired portion of the coding sequence of

PRO such as the sequence encoding the extracellular domain of a transmembrane protein or the sequence encoding the mature protein if the protein is extracellular is amplified by PCR with primers complementary to the 5' and 3' regions. The 5' primer may incorporate flanking (selected) restriction enzyme sites. The product is then digested with those selected restriction enzymes and subcloned into the expression vector.

5 Recombinant baculovirus is generated by co-transfecting the above plasmid and BaculoGold™ virus DNA (PharMingen) into *Spodoptera frugiperda* ("Sf9") cells (ATCC CRL 1711) using lipofectin (commercially available from GIBCO-BRL). After 4 - 5 days of incubation at 28°C, the released viruses are harvested and used for further amplifications. Viral infection and protein expression are performed as described by O'Reilley et al., Baculovirus expression vectors: A Laboratory Manual, Oxford: Oxford University Press (1994).

10 Expressed poly-his tagged PRO can then be purified, for example, by Ni²⁺-chelate affinity chromatography as follows. Extracts are prepared from recombinant virus-infected Sf9 cells as described by Rupert et al., Nature, 362:175-179 (1993). Briefly, Sf9 cells are washed, resuspended in sonication buffer (25 mL Hepes, pH 7.9; 12.5 mM MgCl₂; 0.1 mM EDTA; 10% glycerol; 0.1% NP-40; 0.4 M KCl), and sonicated twice for 20 seconds on ice. The sonicates are cleared by centrifugation, and the supernatant is diluted 50-fold in loading buffer (50 mM phosphate, 300 mM NaCl, 10% glycerol, pH 7.8) and filtered through a 0.45 μm filter. A Ni²⁺-NTA agarose column (commercially available from Qiagen) is prepared with a bed volume of 5 mL, washed with 25 mL of water and equilibrated with 25 mL of loading buffer. The filtered cell extract is loaded onto the column at 0.5 mL per minute. The column is washed to baseline A₂₈₀ with loading buffer, at which point fraction collection is started. Next, the column is washed with a secondary wash buffer (50 mM phosphate; 300 mM NaCl, 10% glycerol, pH 6.0), which elutes nonspecifically bound protein. After reaching A₂₈₀ baseline again, the column is developed with a 0 to 500 mM Imidazole gradient in the secondary wash buffer. One mL fractions are collected and analyzed by SDS-PAGE and silver staining or Western blot with Ni²⁺-NTA-conjugated to alkaline phosphatase (Qiagen). Fractions containing the eluted His₁₀-tagged PRO are pooled and dialyzed against loading buffer.

25 Alternatively, purification of the IgG tagged (or Fc tagged) PRO can be performed using known chromatography techniques, including for instance, Protein A or protein G column chromatography.

Many of the PRO polypeptides disclosed herein were successfully expressed as described above.

EXAMPLE 132: Preparation of Antibodies that Bind PRO

This example illustrates preparation of monoclonal antibodies which can specifically bind PRO.

30 Techniques for producing the monoclonal antibodies are known in the art and are described, for instance, in Goding, supra. Immunogens that may be employed include purified PRO, fusion proteins containing PRO, and cells expressing recombinant PRO on the cell surface. Selection of the immunogen can be made by the skilled artisan without undue experimentation.

35 Mice, such as Balb/c, are immunized with the PRO immunogen emulsified in complete Freund's adjuvant and injected subcutaneously or intraperitoneally in an amount from 1-100 micrograms. Alternatively, the immunogen is emulsified in MPL-TDM adjuvant (Ribi Immunochemical Research, Hamilton, MT) and injected into the animal's hind foot pads. The immunized mice are then boosted 10 to 12 days later with additional immunogen emulsified in the selected adjuvant. Thereafter, for several weeks, the mice may also be boosted with additional immunization injections. Serum samples may be periodically obtained from the mice

by retro-orbital bleeding for testing in ELISA assays to detect anti-PRO antibodies.

After a suitable antibody titer has been detected, the animals "positive" for antibodies can be injected with a final intravenous injection of PRO. Three to four days later, the mice are sacrificed and the spleen cells are harvested. The spleen cells are then fused (using 35% polyethylene glycol) to a selected murine myeloma cell line such as P3X63AgU.1, available from ATCC, No. CRL 1597. The fusions generate hybridoma cells which can then be plated in 96 well tissue culture plates containing HAT (hypoxanthine, aminopterin, and thymidine) medium to inhibit proliferation of non-fused cells, myeloma hybrids, and spleen cell hybrids.

The hybridoma cells will be screened in an ELISA for reactivity against PRO. Determination of "positive" hybridoma cells secreting the desired monoclonal antibodies against PRO is within the skill in the art.

The positive hybridoma cells can be injected intraperitoneally into syngeneic Balb/c mice to produce ascites containing the anti-PRO monoclonal antibodies. Alternatively, the hybridoma cells can be grown in tissue culture flasks or roller bottles. Purification of the monoclonal antibodies produced in the ascites can be accomplished using ammonium sulfate precipitation, followed by gel exclusion chromatography. Alternatively, affinity chromatography based upon binding of antibody to protein A or protein G can be employed.

EXAMPLE 133: Purification of PRO Polypeptides Using Specific Antibodies

Native or recombinant PRO polypeptides may be purified by a variety of standard techniques in the art of protein purification. For example, pro-PRO polypeptide, mature PRO polypeptide, or pre-PRO polypeptide is purified by immunoaffinity chromatography using antibodies specific for the PRO polypeptide of interest. In general, an immunoaffinity column is constructed by covalently coupling the anti-PRO polypeptide antibody to an activated chromatographic resin.

Polyclonal immunoglobulins are prepared from immune sera either by precipitation with ammonium sulfate or by purification on immobilized Protein A (Pharmacia LKB Biotechnology, Piscataway, N.J.). Likewise, monoclonal antibodies are prepared from mouse ascites fluid by ammonium sulfate precipitation or chromatography on immobilized Protein A. Partially purified immunoglobulin is covalently attached to a chromatographic resin such as CnBr-activated SEPHAROSE™ (Pharmacia LKB Biotechnology). The antibody is coupled to the resin, the resin is blocked, and the derivative resin is washed according to the manufacturer's instructions.

Such an immunoaffinity column is utilized in the purification of PRO polypeptide by preparing a fraction from cells containing PRO polypeptide in a soluble form. This preparation is derived by solubilization of the whole cell or of a subcellular fraction obtained via differential centrifugation by the addition of detergent or by other methods well known in the art. Alternatively, soluble PRO polypeptide containing a signal sequence may be secreted in useful quantity into the medium in which the cells are grown.

A soluble PRO polypeptide-containing preparation is passed over the immunoaffinity column, and the column is washed under conditions that allow the preferential absorbance of PRO polypeptide (*e.g.*, high ionic strength buffers in the presence of detergent). Then, the column is eluted under conditions that disrupt antibody/PRO polypeptide binding (*e.g.*, a low pH buffer such as approximately pH 2-3, or a high concentration of a chaotrope such as urea or thiocyanate ion), and PRO polypeptide is collected.

EXAMPLE 134: Drug Screening

This invention is particularly useful for screening compounds by using PRO polypeptides or binding fragment thereof in any of a variety of drug screening techniques. The PRO polypeptide or fragment employed in such a test may either be free in solution, affixed to a solid support, borne on a cell surface, or located intracellularly. One method of drug screening utilizes eukaryotic or prokaryotic host cells which are stably transformed with recombinant nucleic acids expressing the PRO polypeptide or fragment. Drugs are screened against such transformed cells in competitive binding assays. Such cells, either in viable or fixed form, can be used for standard binding assays. One may measure, for example, the formation of complexes between PRO polypeptide or a fragment and the agent being tested. Alternatively, one can examine the diminution in complex formation between the PRO polypeptide and its target cell or target receptors caused by the agent being tested.

Thus, the present invention provides methods of screening for drugs or any other agents which can affect a PRO polypeptide-associated disease or disorder. These methods comprise contacting such an agent with an PRO polypeptide or fragment thereof and assaying (i) for the presence of a complex between the agent and the PRO polypeptide or fragment, or (ii) for the presence of a complex between the PRO polypeptide or fragment and the cell, by methods well known in the art. In such competitive binding assays, the PRO polypeptide or fragment is typically labeled. After suitable incubation, free PRO polypeptide or fragment is separated from that present in bound form, and the amount of free or uncomplexed label is a measure of the ability of the particular agent to bind to PRO polypeptide or to interfere with the PRO polypeptide/cell complex.

Another technique for drug screening provides high throughput screening for compounds having suitable binding affinity to a polypeptide and is described in detail in WO 84/03564, published on September 13, 1984. Briefly stated, large numbers of different small peptide test compounds are synthesized on a solid substrate, such as plastic pins or some other surface. As applied to a PRO polypeptide, the peptide test compounds are reacted with PRO polypeptide and washed. Bound PRO polypeptide is detected by methods well known in the art. Purified PRO polypeptide can also be coated directly onto plates for use in the aforementioned drug screening techniques. In addition, non-neutralizing antibodies can be used to capture the peptide and immobilize it on the solid support.

This invention also contemplates the use of competitive drug screening assays in which neutralizing antibodies capable of binding PRO polypeptide specifically compete with a test compound for binding to PRO polypeptide or fragments thereof. In this manner, the antibodies can be used to detect the presence of any peptide which shares one or more antigenic determinants with PRO polypeptide.

EXAMPLE 135: Rational Drug Design

The goal of rational drug design is to produce structural analogs of biologically active polypeptide of interest (*i.e.*, a PRO polypeptide) or of small molecules with which they interact, *e.g.*, agonists, antagonists, or inhibitors. Any of these examples can be used to fashion drugs which are more active or stable forms of the PRO polypeptide or which enhance or interfere with the function of the PRO polypeptide *in vivo* (*c.f.*, Hodgson, Bio/Technology, 9: 19-21 (1991)).

In one approach, the three-dimensional structure of the PRO polypeptide, or of an PRO polypeptide-inhibitor complex, is determined by x-ray crystallography, by computer modeling or, most typically, by a combination of the two approaches. Both the shape and charges of the PRO polypeptide must be ascertained

to elucidate the structure and to determine active site(s) of the molecule. Less often, useful information regarding the structure of the PRO polypeptide may be gained by modeling based on the structure of homologous proteins. In both cases, relevant structural information is used to design analogous PRO polypeptide-like molecules or to identify efficient inhibitors. Useful examples of rational drug design may include molecules which have improved activity or stability as shown by Braxton and Wells, Biochemistry, 31:7796-7801 (1992) or which act as inhibitors, agonists, or antagonists of native peptides as shown by Athauda *et al.*, J. Biochem., 113:742-746 (1993).

It is also possible to isolate a target-specific antibody, selected by functional assay, as described above, and then to solve its crystal structure. This approach, in principle, yields a pharmacore upon which subsequent drug design can be based. It is possible to bypass protein crystallography altogether by generating anti-idiotypic antibodies (anti-ids) to a functional, pharmacologically active antibody. As a mirror image of a mirror image, the binding site of the anti-ids would be expected to be an analog of the original receptor. The anti-id could then be used to identify and isolate peptides from banks of chemically or biologically produced peptides. The isolated peptides would then act as the pharmacore.

By virtue of the present invention, sufficient amounts of the PRO polypeptide may be made available to perform such analytical studies as X-ray crystallography. In addition, knowledge of the PRO polypeptide amino acid sequence provided herein will provide guidance to those employing computer modeling techniques in place of or in addition to x-ray crystallography.

EXAMPLE 136: Stimulation of Endothelial Cell Proliferation (Assay 8)

This assay is designed to determine whether PRO polypeptides of the present invention show the ability to stimulate adrenal cortical capillary endothelial cell (ACE) growth. PRO polypeptides testing positive in this assay would be expected to be useful for the therapeutic treatment of conditions or disorders where angiogenesis would be beneficial including, for example, wound healing, and the like (as would agonists of these PRO polypeptides). Antagonists of the PRO polypeptides testing positive in this assay would be expected to be useful for the therapeutic treatment of cancerous tumors.

Bovine adrenal cortical capillary endothelial (ACE) cells (from primary culture, maximum of 12-14 passages) were plated in 96-well plates at 500 cells/well per 100 microliter. Assay media included low glucose DMEM, 10% calf serum, 2 mM glutamine, and 1X penicillin/streptomycin/fungizone. Control wells included the following: (1) no ACE cells added; (2) ACE cells alone; (3) ACE cells plus VEGF (5 ng/ml); and (4) ACE cells plus FGF (5ng/ml). The control or test sample, (in 100 microliter volumes), was then added to the wells (at dilutions of 1 %, 0.1 % and 0.01 %, respectively). The cell cultures were incubated for 6-7 days at 37°C/5 % CO₂. After the incubation, the media in the wells was aspirated, and the cells were washed 1X with PBS. An acid phosphatase reaction mixture (100 microliter; 0.1M sodium acetate, pH 5.5, 0.1 % Triton X-100, 10 mM p-nitrophenyl phosphate) was then added to each well. After a 2 hour incubation at 37°C, the reaction was stopped by addition of 10 microliters 1N NaOH. Optical density (OD) was measured on a microplate reader at 405 nm.

The activity of a PRO polypeptide was calculated as the fold increase in proliferation (as determined by the acid phosphatase activity, OD 405 nm) relative to (1) cell only background, and (2) relative to maximum stimulation by VEGF. VEGF (at 3-10 ng/ml) and FGF (at 1-5 ng/ml) were employed as an activity reference

for maximum stimulation. Results of the assay were considered "positive" if the observed stimulation was \geq 50% increase over background. VEGF (5 ng/ml) control at 1% dilution gave 1.24 fold stimulation; FGF (5 ng/ml) control at 1% dilution gave 1.46 fold stimulation.

The following PRO polypeptides tested positive in this assay: PRO1244, PRO1286 and PRO1303.

5 EXAMPLE 137: Inhibitory Activity in Mixed Lymphocyte Reaction (MLR) Assay (Assay 67)

This example shows that one or more of the polypeptides of the invention are active as inhibitors of the proliferation of stimulated T-lymphocytes. Compounds which inhibit proliferation of lymphocytes are useful therapeutically where suppression of an immune response is beneficial.

10 The basic protocol for this assay is described in Current Protocols in Immunology, unit 3.12; edited by J E Coligan, A M Kruisbeek, D H Marglies, E M Shevach, W Strober, National Institutes of Health, Published by John Wiley & Sons, Inc.

15 More specifically, in one assay variant, peripheral blood mononuclear cells (PBMC) are isolated from mammalian individuals, for example a human volunteer, by leukopheresis (one donor will supply stimulator PBMCs, the other donor will supply responder PBMCs). If desired, the cells are frozen in fetal bovine serum and DMSO after isolation. Frozen cells may be thawed overnight in assay media (37°C, 5% CO₂) and then washed and resuspended to 3x10⁶ cells/ml of assay media (RPMI; 10% fetal bovine serum, 1% penicillin/streptomycin, 1% glutamine, 1% HEPES, 1% non-essential amino acids, 1% pyruvate). The stimulator PBMCs are prepared by irradiating the cells (about 3000 Rads).

20 The assay is prepared by plating in triplicate wells a mixture of:

100:1 of test sample diluted to 1% or to 0.1%,

50 :1 of irradiated stimulator cells, and

50 :1 of responder PBMC cells.

25 100 microliters of cell culture media or 100 microliter of CD4-IgG is used as the control. The wells are then incubated at 37°C, 5% CO₂ for 4 days. On day 5, each well is pulsed with tritiated thymidine (1.0 mCi/well; Amersham). After 6 hours the cells are washed 3 times and then the uptake of the label is evaluated.

30 In another variant of this assay, PBMCs are isolated from the spleens of Balb/c mice and C57B6 mice. The cells are teased from freshly harvested spleens in assay media (RPMI; 10% fetal bovine serum, 1% penicillin/streptomycin, 1% glutamine, 1% HEPES, 1% non-essential amino acids, 1% pyruvate) and the PBMCs are isolated by overlaying these cells over Lympholyte M (Organon Teknika), centrifuging at 2000 rpm for 20 minutes, collecting and washing the mononuclear cell layer in assay media and resuspending the cells to 1x10⁷ cells/ml of assay media. The assay is then conducted as described above.

Any decreases below control is considered to be a positive result for an inhibitory compound, with decreases of less than or equal to 80% being preferred. However, any value less than control indicates an inhibitory effect for the test protein.

35 The following polypeptide tested positive in this assay: PRO1250, PRO1418 and PRO1410.

EXAMPLE 138: Stimulation of Heart Neonatal Hypertrophy (Assay 1)

This assay is designed to measure the ability of PRO polypeptides to stimulate hypertrophy of neonatal heart. PRO polypeptides testing positive in this assay would be expected to be useful for the therapeutic

treatment of various cardiac insufficiency disorders.

Cardiac myocytes from 1-day old Harlan Sprague Dawley rats were obtained. Cells ($180 \mu\text{l}$ at $7.5 \times 10^4/\text{ml}$, serum $<0.1\%$, freshly isolated) are added on day 1 to 96-well plates previously coated with DMEM/F12 + 4% FCS. Test samples containing the test PRO polypeptide are added directly to wells on day 2 in $20 \mu\text{L}$ volumes. Cells are stained with crystal violet after an additional two days and scored visually by the next day.

5 Incubator conditions require $5\% \text{CO}_2$.

Activity reference: phenylephrine at $1\text{--}100 \mu\text{M}$, PGF-2 alpha at $0.1\text{--}1.0 \mu\text{M}$, endothelin-1 at $1\text{--}10 \text{nM}$, CT1 (LIF) at $1\text{--}10 \text{nM}$. No PBS is included, since calcium concentration is critical for assay response. Assay media included: DMEM/F12 (with 2.44 gm bicarbonate), $10 \mu\text{g}/\text{ml}$ transferrin, $1 \mu\text{g}/\text{ml}$ insulin, $1 \mu\text{g}/\text{ml}$ aprotinin, 2 mmol/L glutamine, $100 \text{ U}/\text{ml}$ penicillin G, $100 \mu\text{g}/\text{ml}$ streptomycin. Protein buffer containing mannitol (4%) gave a positive signal (score 3.5) at $1/10$ (0.4%) and $1/100$ (0.04%), but not at $1/1000$ (0.004%). Therefore, the test sample buffer containing mannitol is not run. A secondary assay consists of measuring the ANP levels (ng/ml) by ELISA in conditioned media from the cells. An increase in the ANP message can be measured by PCR from cells after a few hours.

15 Results are assessed by visually observing cell size: a score = 3.5 or greater is considered positive for conditioned media; a score of 3.0 or greater is considered positive for purified protein.

The following purified PRO polypeptide was observed to stimulate neonatal heart hypertrophy in this assay: PRO1246.

20 EXAMPLE 139: Inhibition of Vascular Endothelial Growth Factor (VEGF) Stimulated Proliferation of Endothelial Cell Growth (Assay 9)

The ability of various PRO polypeptides to inhibit VEGF stimulated proliferation of endothelial cells was tested. Polypeptides testing positive in this assay are useful for inhibiting endothelial cell growth in mammals where such an effect would be beneficial, e.g., for inhibiting tumor growth.

25 Specifically, bovine adrenal cortical capillary endothelial cells (ACE) (from primary culture, maximum of $12\text{--}14$ passages) were plated in 96-well plates at $500 \text{ cells}/\text{well}$ per 100 microliter . Assay media included low glucose DMEM, 10% calf serum, 2 mM glutamine, and 1X penicillin/streptomycin/fungizone. Control wells included the following: (1) no ACE cells added; (2) ACE cells alone; (3) ACE cells plus $5 \text{ ng}/\text{ml}$ FGF; (4) ACE cells plus $3 \text{ ng}/\text{ml}$ VEGF; (5) ACE cells plus $3 \text{ ng}/\text{ml}$ VEGF plus $1 \text{ ng}/\text{ml}$ TGF-beta; and (6) ACE cells plus $3 \text{ ng}/\text{ml}$ VEGF plus $5 \text{ ng}/\text{ml}$ LIF. The test samples, poly-his tagged PRO polypeptides (in 100 microliter volumes), were then added to the wells (at dilutions of 1% , 0.1% and 0.01% , respectively). The cell cultures were incubated for $6\text{--}7$ days at $37^\circ\text{C}/5\% \text{CO}_2$. After the incubation, the media in the wells was aspirated, and the cells were washed 1X with PBS. An acid phosphatase reaction mixture (100 microliter ; 0.1M sodium acetate, $\text{pH } 5.5$, 0.1% Triton X-100, 10 mM p-nitrophenyl phosphate) was then added to each well. After a 2 hour incubation at 37°C , the reaction was stopped by addition of $10 \text{ microliters } 1\text{N NaOH}$. Optical density (OD) 35 was measured on a microplate reader at 405 nm .

The activity of PRO polypeptides was calculated as the percent inhibition of VEGF ($3 \text{ ng}/\text{ml}$) stimulated proliferation (as determined by measuring acid phosphatase activity at OD 405 nm) relative to the cells without stimulation. TGF-beta was employed as an activity reference at $1 \text{ ng}/\text{ml}$, since TGF-beta blocks $70\text{--}90\%$ of VEGF-stimulated ACE cell proliferation. The results are indicative of the utility of the PRO polypeptides in

cancer therapy and specifically in inhibiting tumor angiogenesis. Numerical values (relative inhibition) are determined by calculating the percent inhibition of VEGF stimulated proliferation by the PRO polypeptides relative to cells without stimulation and then dividing that percentage into the percent inhibition obtained by TGF- β at 1 ng/ml which is known to block 70-90% of VEGF stimulated cell proliferation. The results are considered positive if the PRO polypeptide exhibits 30% or greater inhibition of VEGF stimulation of endothelial cell growth (relative inhibition 30% or greater).

The following polypeptide tested positive in this assay: PRO1246.

EXAMPLE 140: Human Venous Endothelial Cell Calcium Flux Assay (Assay 68)

This assay is designed to determine whether PRO polypeptides show the ability to stimulate calcium flux in human umbilical vein endothelial cells (HUVEC, Cell Systems). Calcium influx is a well documented response upon binding of certain ligands to their receptors. A test compound that results in a positive response in the present calcium influx assay can be said to bind to a specific receptor and activate a biological signaling pathway in human endothelial cells. This will ultimately lead, for example, to cell division, inhibition of cell proliferation, endothelial tube formation, cell migration, apoptosis, etc.

Human venous umbilical vein endothelial cells (HUVEC, Cell Systems) in growth media (50:50 without glycine, 1% glutamine, 10mM Hepes, 10% FBS, 10 ng/ml bFGF), were plated on 96-well microtiter ViewPlates-96 (Packard Instrument Company Part #6005182) microtiter plates at a cell density of 2×10^4 cells/well. The day after plating, the cells were washed three times with buffer (HBSS plus 10 mM Hepes), leaving 100 μ l/well. Then 100 μ l/well of 8 μ M Fluo-3 (2x) was added. The cells were incubated for 1.5 hours at 37°C/5% CO₂. After incubation, the cells were then washed 3x with buffer (described above) leaving 100 μ l/well. Test samples of the PRO polypeptides were prepared on different 96-well plates at 5x concentration in buffer. The positive control corresponded to 50 μ M ionomycin (5x); the negative control corresponded to Protein 32. Cell plate and sample plates were run on a FLIPR (Molecular Devices) machine. The FLIPR machine added 25 μ l of test sample to the cells, and readings were taken every second for one minute, then every 3 seconds for the next three minutes.

The fluorescence change from baseline to the maximum rise of the curve (Δ change) was calculated, and replicates averaged. The rate of fluorescence increase was monitored, and only those samples which had a Δ change greater than 1000 and a rise within 60 seconds, were considered positive.

The following PRO polypeptides tested positive in this assay: PRO1246 and PRO1561.

EXAMPLE 141: Skin Vascular Permeability Assay (Assay 64)

This assay shows that certain polypeptides of the invention stimulate an immune response and induce inflammation by inducing mononuclear cell, eosinophil and PMN infiltration at the site of injection of the animal. Compounds which stimulate an immune response are useful therapeutically where stimulation of an immune response is beneficial. This skin vascular permeability assay is conducted as follows. Hairless guinea pigs weighing 350 grams or more are anesthetized with ketamine (75-80 mg/Kg) and 5 mg/Kg xylazine intramuscularly (IM). A sample of purified polypeptide of the invention or a conditioned media test sample is injected intradermally onto the backs of the test animals with 100 μ l per injection site. It is possible to have

about 10-30, preferably about 16-24, injection sites per animal. One μ l of Evans blue dye (1% in physiologic buffered saline) is injected intracardially. Blemishes at the injection sites are then measured (mm diameter) at 1 hr and 6 hr post injection. Animals were sacrificed at 6 hrs after injection. Each skin injection site is biopsied and fixed in formalin. The skins are then prepared for histopathologic evaluation. Each site is evaluated for inflammatory cell infiltration into the skin. Sites with visible inflammatory cell inflammation are scored as positive. Inflammatory cells may be neutrophilic, eosinophilic, monocytic or lymphocytic. At least a minimal perivascular infiltrate at the injection site is scored as positive, no infiltrate at the site of injection is scored as negative.

The following polypeptide tested positive in this assay: PRO1283, PRO1325 and PRO1343.

10 EXAMPLE 142: Induction of c-fos in Endothelial Cells (Assay 34)

This assay is designed to determine whether PRO polypeptides show the ability to induce c-fos in endothelial cells. PRO polypeptides testing positive in this assay would be expected to be useful for the therapeutic treatment of conditions or disorders where angiogenesis would be beneficial including, for example, wound healing, and the like (as would agonists of these PRO polypeptides). Antagonists of the PRO polypeptides testing positive in this assay would be expected to be useful for the therapeutic treatment of cancerous tumors.

Human venous umbilical vein endothelial cells (HUVEC, Cell Systems) in growth media (50% Ham's F12 w/o GHT: low glucose, and 50% DMEM without glycine: with NaHCO_3 , 1% glutamine, 10 mM HEPES, 10% FBS, 10 ng/ml bFGF) were plated on 96-well microtiter plates at a cell density of 1×10^4 cells/well. The day after plating, the cells were starved by removing the growth media and treating the cells with 100 μ l/well test samples and controls (positive control = growth media; negative control = Protein 32 buffer = 10 mM HEPES, 140 mM NaCl, 4% (w/v) mannitol, pH 6.8). The cells were incubated for 30 minutes at 37°C, in 5% CO_2 . The samples were removed, and the first part of the bDNA kit protocol (Chiron Diagnostics, cat. #6005-037) was followed, where each capitalized reagent/buffer listed below was available from the kit.

Briefly, the amounts of the TM Lysis Buffer and Probes needed for the tests were calculated based on information provided by the manufacturer. The appropriate amounts of thawed Probes were added to the TM Lysis Buffer. The Capture Hybridization Buffer was warmed to room temperature. The bDNA strips were set up in the metal strip holders, and 100 μ l of Capture Hybridization Buffer was added to each b-DNA well needed, followed by incubation for at least 30 minutes. The test plates with the cells were removed from the incubator, and the media was gently removed using the vacuum manifold. 100 μ l of Lysis Hybridization Buffer with Probes were quickly pipetted into each well of the microtiter plates. The plates were then incubated at 55°C for 15 minutes. Upon removal from the incubator, the plates were placed on the vortex mixer with the microtiter adapter head and vortexed on the #2 setting for one minute. 80 μ l of the lysate was removed and added to the bDNA wells containing the Capture Hybridization Buffer, and pipetted up and down to mix. The plates were incubated at 53°C for at least 16 hours.

On the next day, the second part of the bDNA kit protocol was followed. Specifically, the plates were removed from the incubator and placed on the bench to cool for 10 minutes. The volumes of additions needed were calculated based upon information provided by the manufacturer. An Amplifier Working Solution was

prepared by making a 1:100 dilution of the Amplifier Concentrate (20 fm/ μ l) in AL Hybridization Buffer. The hybridization mixture was removed from the plates and washed twice with Wash A. 50 μ l of Amplifier Working Solution was added to each well and the wells were incubated at 53°C for 30 minutes. The plates were then removed from the incubator and allowed to cool for 10 minutes. The Label Probe Working Solution was prepared by making a 1:100 dilution of Label Concentrate (40 pmoles/ μ l) in AL Hybridization Buffer. After the 10-minute cool-down period, the amplifier hybridization mixture was removed and the plates were washed twice with Wash A. 50 μ l of Label Probe Working Solution was added to each well and the wells were incubated at 53°C for 15 minutes. After cooling for 10 minutes, the Substrate was warmed to room temperature. Upon addition of 3 μ l of Substrate Enhancer to each ml of Substrate needed for the assay, the plates were allowed to cool for 10 minutes, the label hybridization mixture was removed, and the plates were washed twice with Wash A and three times with Wash D. 50 μ l of the Substrate Solution with Enhancer was added to each well. The plates were incubated for 30 minutes at 37°C and RLU was read in an appropriate luminometer.

The replicates were averaged and the coefficient of variation was determined. The measure of activity of the fold increase over the negative control (Protein 32/HEPES buffer described above) value was indicated by chemiluminescence units (RLU). The results are considered positive if the PRO polypeptide exhibits at least a two-fold value over the negative buffer control. Negative control = 1.00 RLU at 1.00% dilution. Positive control = 8.39 RLU at 1.00% dilution.

The following PRO polypeptides tested positive in this assay: PRO1274, PRO1294, PRO1304 and PRO1130.

EXAMPLE 143: Gene Amplification

This example shows that the PRO1295-, PRO1293-, PRO1265-, PRO1303-, PRO1269-, PRO1410-, PRO1317-, PRO1780-, PRO1555-, PRO1755-, PRO1558-, PRO1759- and PRO1788-encoding genes are amplified in the genome of certain human lung, colon and/or breast cancers and/or cell lines. Amplification is associated with overexpression of the gene product, indicating that the polypeptides are useful targets for therapeutic intervention in certain cancers such as colon, lung, breast and other cancers. Therapeutic agents may take the form of antagonists of PRO1295, PRO1293, PRO1265, PRO1303, PRO1269, PRO1410, PRO1317, PRO1780, PRO1555, PRO1755, PRO1558, PRO1759 and PRO1788 polypeptides, for example, murine-human chimeric, humanized or human antibodies against a PRO1295, PRO1293, PRO1265, PRO1303, PRO1269, PRO1410, PRO1317, PRO1780, PRO1555, PRO1755, PRO1558, PRO1759 or PRO1788 polypeptide.

The starting material for the screen was genomic DNA isolated from a variety of cancers. The DNA is quantitated precisely, *e.g.*, fluorometrically. As a negative control, DNA was isolated from the cells of ten normal healthy individuals which was pooled and used as assay controls for the gene copy in healthy individuals (not shown). The 5' nuclease assay (for example, TaqMan™) and real-time quantitative PCR (for example, ABI Prizm 7700 Sequence Detection System™ (Perkin Elmer, Applied Biosystems Division, Foster City, CA)), were used to find genes potentially amplified in certain cancers. The results were used to determine whether the DNA encoding PRO1295, PRO1293, PRO1265, PRO1303, PRO1269, PRO1410, PRO1317, PRO1780, PRO1555, PRO1755, PRO1558, PRO1759 and PRO1788 is over-represented in any of the primary lung or colon cancers or cancer cell lines or breast cancer cell lines that were screened. The primary lung cancers were obtained from individuals with tumors of the type and stage as indicated in Table 7. An explanation of the abbreviations used

for the designation of the primary tumors listed in Table 7 and the primary tumors and cell lines referred to throughout this example has been given hereinbefore.

5 The results of the TaqMan™ are reported in delta (Δ) Ct units. One unit corresponds to 1 PCR cycle or approximately a 2-fold amplification relative to normal, two units corresponds to 4-fold, 3 units to 8-fold amplification and so on. Quantitation was obtained using primers and a TaqMan™ fluorescent probe derived
10 from the PRO1295-, PRO1293-, PRO1265-, PRO1303-, PRO1269-, PRO1410-, PRO1317-, PRO1780-, PRO1555-, PRO1755-, PRO1558-, PRO1759- and PRO1788-encoding gene. Regions of PRO1295, PRO1293, PRO1265, PRO1303, PRO1269, PRO1410, PRO1317, PRO1780, PRO1555, PRO1755, PRO1558, PRO1759 and PRO1788 which are most likely to contain unique nucleic acid sequences and which are least likely to have spliced out introns are preferred for the primer and probe derivation, *e.g.*, 3'-untranslated regions. The sequences for the primers and probes (forward, reverse and probe) used for the PRO1295, PRO1293, PRO1265, PRO1303, PRO1269, PRO1410, PRO1317, PRO1780, PRO1555, PRO1755, PRO1558, PRO1759 and PRO1788 gene amplification analysis were as follows:

- PRO1295 (DNA59218-1559)
forward: 5'-AGGACTTGCCCTCAGGAA-3' (SEQ ID NO:432)
reverse: 5'-CGCAGGACAGTTGTGAAAATA-3' (SEQ ID NO:433)
probe: 5'-ATGACGCTCGTCCAAGGCCAC-3' (SEQ ID NO:434)
- 5 PRO1293 (DNA60618-1557)
forward: 5'-CCCACCTGTACCACCATGT-3' (SEQ ID NO:435)
probe: 5'-ACTCCAGGCACCATCTGTTCTCCC-3' (SEQ ID NO:436)
reverse: 5'-AAGGGCTGGCATTCAAGTU-3' (SEQ ID NO:437)
- 10 PRO1265 (DNA60764-1533)
forward: 5'-TGACCTGGCAAAGGAAGAA-3' (SEQ ID NO:438)
probe: 5'-CAGCCACCCTCCAGTCCAAGG-3' (SEQ ID NO:439)
reverse: 5'-GGGTCGTGTTTTGGAGAGA-3' (SEQ ID NO:440)
- 15 PRO1303 (DNA65409-1566)
forward: 5'-CTGGCCCTCAGAGCACCAAT-3' (SEQ ID NO:441)
probe: 5'-TCCTCCATCACTTCCCCTAGCTCCA-3' (SEQ ID NO:442)
reverse: 5'-CTGGCAGGAGTTAAAGTTCCAAGA-3' (SEQ ID NO:443)
- 20 PRO1269 (DNA66520-1536)
forward: 5'-AAAGGACACCGGGATGTG-3' (SEQ ID NO:444)
probe: 5'-AGCGTACACTCTCTCCAGGCAACCAG-3' (SEQ ID NO:445)
reverse: 5'-CAATTCTGGATGAGGTGGTAGA-3' (SEQ ID NO:446)
- 25 PRO1410 (DNA68874-1622)
forward: 5'-CAGGACTGAGCGCTTGTTTA-3' (SEQ ID NO:447)
probe: 5'-CAAAGCGCCAAGTACCGGACC-3' (SEQ ID NO:448)
reverse: 5'-CCAGACCTCAGCCAGGAA-3' (SEQ ID NO:449)
- 30 PRO1317 (DNA71166-1685)
forward: 5'-CCCTAGCTGACCCCTTCA-3' (SEQ ID NO:450)
reverse: 5'-TCTGACAAGCAGTTTTCTGAATC-3' (SEQ ID NO:451)
probe: 5'-CTCTCCCCCTCCCTTTTCCTTTGTTT-3' (SEQ ID NO:452)
- 35 PRO1780 (DNA71169-1709)
forward: 5'-CTCTGGTGCCACAGTGA-3' (SEQ ID NO:453)
probe: 5'-CCATGCCTGCTCAGCCAAGAA-3' (SEQ ID NO:454)
reverse: 5'-CAGGAAATCTGGAAACCTACAGT-3' (SEQ ID NO:455)

PRO1555 (DNA73744-1665)

forward: 5'-CCTTGAAAAGGACCCAGTTT-3' (SEQ ID NO:456)

probe: 5'-ATGAGTCGCACCTGCTGTTCCC-3' (SEQ ID NO:457)

reverse: 5'-TAGCAGCTGCCCTTGGTA-3' (SEQ ID NO:458)

forward: 5'-AACAGCAGGTGCGACTCATCTA-3' (SEQ ID NO:459)

5 probe: 5'-TGCTAGGCGACGACACCCAGACC-3' (SEQ ID NO:460)

reverse: 5'-TGGACACGTGGCAGTGGA-3' (SEQ ID NO:461)

PRO1755 (DNA76396-1698)

forward: 5'-TCATGGTCTCGTCCCATTC-3' (SEQ ID NO:462)

10 probe: 5'-CACCATTGTGTTTCTCTGTCTCCCCATC-3' (SEQ ID NO:463)

reverse: 5'-CCGGCATCCTTGGAGTAG-3' (SEQ ID NO:464)

PRO1788 (DNA77652-2505)

forward: 5'-TCCCCATTAGCACAGGAGTA-3' (SEQ ID NO:465)

15 probe: 5'-AGGCTCTTGCCTGTCTGCTGCT-3' (SEQ ID NO:466)

reverse: 5'-GCCCAGAGTCCCACTTGT-3' (SEQ ID NO:467)

PRO1558 (DNA71282-1668)

forward: 5'-ACTGCTCCGCTACTACGA -3' (SEQ ID NO:468)

20 probe: 5'-AGGCATCCTCGCCGTCCTCA -3' (SEQ ID NO:469)

reverse: 5'-AAGGCCAAGGTGAGTCCAT -3' (SEQ ID NO:470)

forward: 5'-CGAGTGTGTGCGAAACCTAA -3' (SEQ ID NO:471)

probe: 5'-TCAGGGTCTACATCAGCCTCCTGC -3' (SEQ ID NO:472)

reverse: 5'-AAGGCCAAGGTGAGTCCAT -3' (SEQ ID NO:473)

PRO1759 (DNA76531-1701)

forward: 5'-CCTACTGAGGAGCCCTATGC -3' (SEQ ID NO:474)

probe: 5'-CCTGAGCTGTAACCCCACTCCAGG -3' (SEQ ID NO:231)

reverse: 5'-AGAGTCTGTCCCAGCTATCTTGT -3' (SEQ ID NO:232)

35 The 5' nuclease assay reaction is a fluorescent PCR-based technique which makes use of the 5' exonuclease activity of Taq DNA polymerase enzyme to monitor amplification in real time. Two oligonucleotide primers are used to generate an amplicon typical of a PCR reaction. A third oligonucleotide, or probe, is designed to detect nucleotide sequence located between the two PCR primers. The probe is non-extendible by Taq DNA polymerase enzyme, and is labeled with a reporter fluorescent dye and a quencher fluorescent dye. Any laser-induced emission from the reporter dye is quenched by the quenching dye when the two dyes are located close together as they are on the probe. During the amplification reaction, the Taq DNA polymerase enzyme cleaves the probe in a template-dependent manner. The resultant probe fragments disassociate in solution, and signal from the released reporter dye is free from the quenching effect of the second fluorophore.

One molecule of reporter dye is liberated for each new molecule synthesized, and detection of the unquenched reporter dye provides the basis for quantitative interpretation of the data.

The 5' nuclease procedure is run on a real-time quantitative PCR device such as the ABI Prism 7700TM Sequence Detection. The system consists of a thermocycler, laser, charge-coupled device (CCD) camera and computer. The system amplifies samples in a 96-well format on a thermocycler. During amplification, laser-induced fluorescent signal is collected in real-time through fiber optics cables for all 96 wells, and detected at the CCD. The system includes software for running the instrument and for analyzing the data.

5' Nuclease assay data are initially expressed as Ct, or the threshold cycle. This is defined as the cycle at which the reporter signal accumulates above the background level of fluorescence. The ΔC_t values are used as quantitative measurement of the relative number of starting copies of a particular target sequence in a nucleic acid sample when comparing cancer DNA results to normal human DNA results.

Table 7 describes the stage, T stage and N stage of various primary tumors which were used to screen the PRO1295, PRO1293, PRO1265, PRO1303, PRO1269, PRO1410, PRO1317, PRO1780, PRO1555, PRO1755, PRO1558, PRO1759 and PRO1788 compounds of the invention.

Table 7
Primary Lung and Colon Tumor Profiles

	Primary Tumor	Stage	Other Stage	Dukes Stage	T Stage	N Stage
5	Human lung tumor AdenoCa (SRCC724) [LT1]	IIA			T1	N1
	Human lung tumor SqCCa (SRCC725) [LT1a]	IIB			T3	N0
	Human lung tumor AdenoCa (SRCC726) [LT2]	IB			T2	N0
	Human lung tumor AdenoCa (SRCC727) [LT3]	IIIA			T1	N2
	Human lung tumor AdenoCa (SRCC728) [LT4]	IB			T2	N0
	Human lung tumor SqCCa (SRCC729) [LT6]	IB			T2	N0
10	Human lung tumor Aden/SqCCa (SRCC730) [LT7]	IA			T1	N0
	Human lung tumor AdenoCa (SRCC731) [LT9]	IB			T2	N0
	Human lung tumor SqCCa (SRCC732) [LT10]	IIB			T2	N1
	Human lung tumor SqCCa (SRCC733) [LT11]	IIA			T1	N1
	Human lung tumor AdenoCa (SRCC734) [LT12]	IV			T2	N0
15	Human lung tumor AdenoSqCCa (SRCC735)[LT13]	IIB			T2	N0
	Human lung tumor SqCCa (SRCC736) [LT15]	IB			T2	N0
	Human lung tumor SqCCa (SRCC737) [LT16]	IB			T2	N0
	Human lung tumor SqCCa (SRCC738) [LT17]	IIB			T2	N1
	Human lung tumor SqCCa (SRCC739) [LT18]	IB			T2	N0
20	Human lung tumor SqCCa (SRCC740) [LT19]	IB			T2	N0
	Human lung tumor LCCa (SRCC741) [LT21]	IIB			T3	N1
	Human lung AdenoCa (SRCC811) [LT22]	1A			T1	N0
	Human colon AdenoCa (SRCC742) [CT2]		M1	D	pT4	N0
	Human colon AdenoCa (SRCC743) [CT3]			B	pT3	N0
25	Human colon AdenoCa (SRCC 744) [CT8]			B	T3	N0
	Human colon AdenoCa (SRCC745) [CT10]		A	pT2	N0	
	Human colon AdenoCa (SRCC746) [CT12]	MO, R1	B	T3	N0	
	Human colon AdenoCa (SRCC747) [CT14]	pMO, RO	B	pT3	pN0	
	Human colon AdenoCa (SRCC748) [CT15]	M1, R2	D	T4	N2	
30	Human colon AdenoCa (SRCC749) [CT16]	pMO	B	pT3	pN0	
	Human colon AdenoCa (SRCC750) [CT17]		C1	pT3	pN1	
	Human colon AdenoCa (SRCC751) [CT1]		MO, R1	B	pT3	N0
	Human colon AdenoCa (SRCC752) [CT4]			B	pT3	M0
	Human colon AdenoCa (SRCC753) [CT5]		G2	C1	pT3	pN0
35	Human colon AdenoCa (SRCC754) [CT6]		pMO, RO	B	pT3	pN0
	Human colon AdenoCa (SRCC755) [CT7]		G1	A	pT2	pN0
	Human colon AdenoCa (SRCC756) [CT9]		G3	D	pT4	pN2
	Human colon AdenoCa (SRCC757) [CT11]		B	T3	N0	
40	Human colon AdenoCa (SRCC758) [CT18]	MO, RO	B	pT3	pN0	

DNA Preparation:

DNA was prepared from cultured cell lines, primary tumors, and normal human blood. The isolation was performed using purification kit, buffer set and protease and all from Qiagen, according to the manufacturer's instructions and the description below.

Cell culture lysis:

Cells were washed and trypsinized at a concentration of 7.5×10^8 per tip and pelleted by centrifuging at 1000 rpm for 5 minutes at 4°C, followed by washing again with 1/2 volume of PBS and recentrifugation. The pellets were washed a third time, the suspended cells collected and washed 2x with PBS. The cells were then suspended into 10 ml PBS. Buffer C1 was equilibrated at 4°C. Qiagen protease #19155 was diluted into 6.25 ml cold ddH₂O to a final concentration of 20 mg/ml and equilibrated at 4°C. 10 ml of G2 Buffer was prepared by diluting Qiagen RNase A stock (100 mg/ml) to a final concentration of 200 µg/ml.

Buffer C1 (10 ml, 4°C) and ddH₂O (40 ml, 4°C) were then added to the 10 ml of cell suspension, mixed by inverting and incubated on ice for 10 minutes. The cell nuclei were pelleted by centrifuging in a Beckman swinging bucket rotor at 2500 rpm at 4°C for 15 minutes. The supernatant was discarded and the nuclei were suspended with a vortex into 2 ml Buffer C1 (at 4°C) and 6 ml ddH₂O, followed by a second 4°C centrifugation at 2500 rpm for 15 minutes. The nuclei were then resuspended into the residual buffer using 200 µl per tip. G2 buffer (10 ml) was added to the suspended nuclei while gentle vortexing was applied. Upon completion of buffer addition, vigorous vortexing was applied for 30 seconds. Quiagen protease (200 µl, prepared as indicated above) was added and incubated at 50°C for 60 minutes. The incubation and centrifugation were repeated until the lysates were clear (*e.g.*, incubating additional 30–60 minutes, pelleting at 3000 x g for 10 min., 4°C).

Solid human tumor sample preparation and lysis:

Tumor samples were weighed and placed into 50 ml conical tubes and held on ice. Processing was limited to no more than 250 mg tissue per preparation (1 tip/preparation). The protease solution was freshly prepared by diluting into 6.25 ml cold ddH₂O to a final concentration of 20 mg/ml and stored at 4°C. G2 buffer (20 ml) was prepared by diluting DNase A to a final concentration of 200 mg/ml (from 100 mg/ml stock). The tumor tissue was homogenated in 19 ml G2 buffer for 60 seconds using the large tip of the polytron in a laminar-flow TC hood in order to avoid inhalation of aerosols, and held at room temperature. Between samples, the polytron was cleaned by spinning at 2 x 30 seconds each in 2L ddH₂O, followed by G2 buffer (50 ml). If tissue was still present on the generator tip, the apparatus was disassembled and cleaned.

Quiagen protease (prepared as indicated above, 1.0 ml) was added, followed by vortexing and incubation at 50°C for 3 hours. The incubation and centrifugation were repeated until the lysates were clear (*e.g.*, incubating additional 30–60 minutes, pelleting at 3000 x g for 10 min., 4°C).

Human blood preparation and lysis:

Blood was drawn from healthy volunteers using standard infectious agent protocols and citrated into 10 ml samples per tip. Quiagen protease was freshly prepared by dilution into 6.25 ml cold ddH₂O to a final concentration of 20 mg/ml and stored at 4°C. G2 buffer was prepared by diluting RNAse A to a final concentration of 200 µg/ml from 100 mg/ml stock. The blood (10 ml) was placed into a 50 ml conical tube and 10 ml C1 buffer and 30 ml ddH₂O (both previously equilibrated to 4°C) were added, and the components mixed by inverting and held on ice for 10 minutes. The nuclei were pelleted with a Beckman swinging bucket rotor at 2500 rpm, 4°C for 15 minutes and the supernatant discarded. With a vortex, the nuclei were suspended into 2 ml C1 buffer (4°C) and 6 ml ddH₂O (4°C). Vortexing was repeated until the pellet was white. The nuclei were then suspended into the residual buffer using a 200 µl tip. G2 buffer (10 ml) was added to the suspended nuclei while gently vortexing, followed by vigorous vortexing for 30 seconds. Quiagen protease was added (200 µl) and incubated at 50°C for 60 minutes. The incubation and centrifugation were repeated until the lysates were clear (*e.g.*, incubating additional 30–60 minutes, pelleting at 3000 x g for 10 min., 4°C).

Purification of cleared lysates:

(1) Isolation of genomic DNA:

Genomic DNA was equilibrated (1 sample per maxi tip preparation) with 10 ml QBT buffer. QF elution buffer was equilibrated at 50°C. The samples were vortexed for 30 seconds, then loaded onto equilibrated tips and drained by gravity. The tips were washed with 2 x 15 ml QC buffer. The DNA was eluted into 30 ml silanized, autoclaved 30 ml Corex tubes with 15 ml QF buffer (50°C). Isopropanol (10.5 ml) was added to each

sample, the tubes covered with parafin and mixed by repeated inversion until the DNA precipitated. Samples were pelleted by centrifugation in the SS-34 rotor at 15,000 rpm for 10 minutes at 4°C. The pellet location was marked, the supernatant discarded, and 10 ml 70% ethanol (4°C) was added. Samples were pelleted again by centrifugation on the SS-34 rotor at 10,000 rpm for 10 minutes at 4°C. The pellet location was marked and the supernatant discarded. The tubes were then placed on their side in a drying rack and dried 10 minutes at 37°C, taking care not to overdry the samples.

After drying, the pellets were dissolved into 1.0 ml TE (pH 8.5) and placed at 50°C for 1-2 hours. Samples were held overnight at 4°C as dissolution continued. The DNA solution was then transferred to 1.5 ml tubes with a 26 gauge needle on a tuberculin syringe. The transfer was repeated 5x in order to shear the DNA. Samples were then placed at 50°C for 1-2 hours.

(2) Quantitation of genomic DNA and preparation for gene amplification assay:

The DNA levels in each tube were quantified by standard A_{260}/A_{280} spectrophotometry on a 1:20 dilution (5 μ l DNA + 95 μ l ddH₂O) using the 0.1 ml quartz cuvettes in the Beckman DU640 spectrophotometer. A_{260}/A_{280} ratios were in the range of 1.8-1.9. Each DNA sample was then diluted further to approximately 200 ng/ml in TE (pH 8.5). If the original material was highly concentrated (about 700 ng/ μ l), the material was placed at 50°C for several hours until resuspended.

Fluorometric DNA quantitation was then performed on the diluted material (20-600 ng/ml) using the manufacturer's guidelines as modified below. This was accomplished by allowing a Hoeffer DyNA Quant 200 fluorometer to warm-up for about 15 minutes. The Hoechst dye working solution (#H33258, 10 μ l, prepared within 12 hours of use) was diluted into 100 ml 1 x TNE buffer. A 2 ml cuvette was filled with the fluorometer solution, placed into the machine, and the machine was zeroed. pGEM 3Zf(+) (2 μ l, lot #360851026) was added to 2 ml of fluorometer solution and calibrated at 200 units. An additional 2 μ l of pGEM 3Zf(+) DNA was then tested and the reading confirmed at 400 +/- 10 units. Each sample was then read at least in triplicate. When 3 samples were found to be within 10% of each other, their average was taken and this value was used as the quantification value.

The fluorometrically determined concentration was then used to dilute each sample to 10 ng/ μ l in ddH₂O. This was done simultaneously on all template samples for a single TaqMan plate assay, and with enough material to run 500-1000 assays. The samples were tested in triplicate with Taqman™ primers and probe both B-actin and GAPDH on a single plate with normal human DNA and no-template controls. The diluted samples were used provided that the CT value of normal human DNA subtracted from test DNA was +/- 1 Ct. The diluted, lot-qualified genomic DNA was stored in 1.0 ml aliquots at -80°C. Aliquots which were subsequently to be used in the gene amplification assay were stored at 4°C. Each 1 ml aliquot is enough for 8-9 plates or 64 tests.

Gene amplification assay:

The PRO1295, PRO1293, PRO1265, PRO1303, PRO1269, PRO1410, PRO1317, PRO1780, PRO1555, PRO1755, PRO1558, PRO1759 and PRO1788 compounds of the invention were screened in the following primary tumors and the resulting Δ Ct values which are ≥ 1.0 are reported in Table 8.

Table 8

 Δ Ct values in lung and colon primary tumors and cell line models

5	Primary Tumors or Cell lines	PRO 1293	PRO 1269	PRO 1410	PRO 1755	PRO 1780	PRO 1788	PRO 1303	PRO 1555	PRO 1265	PRO 1317	PRO 1295	PRO 1558	PRO 1759
	LT1	---	---	---	---	---	---	---	---	---	1.15	---	---	---
	LT1-a	---	---	---	---	---	---	---	---	---	1.49	---	---	---
10	LT3	---	---	---	---	---	---	---	---	1.04	---	---	---	---
	LT4	---	---	---	---	1.16	---	---	---	---	---	---	---	---
	LT7	---	---	---	---	1.02	---	---	---	---	---	---	---	---
	LT9	---	---	---	---	---	---	---	---	---	1.26	---	---	---
	LT10	---	---	---	---	---	---	---	---	---	1.68	---	---	---
15	LT12	---	---	---	---	---	---	---	---	2.17	---	---	---	---
	LT13	---	---	1.12 1.42	---	---	---	1.42	4.20 4.45	2.24	---	---	---	---
	LT15	---	1.22	2.10 1.82	---	---	---	1.17	1.36 1.15	3.51	1.16	---	---	---
	LT16	---	1.14	1.44 1.45	1.36	---	---	1.42	3.71 3.99	3.32	---	---	---	---
	LT17	---	1.26	---	---	---	---	---	---	1.02	1.74	---	---	---
20	LT18	---	---	---	1.18	---	---	---	---	---	---	---	---	---
	CT2	---	---	2.36	2.35	---	---	---	---	---	---	---	---	---
	CT3	---	---	1.09	---	---	1.35	---	---	---	---	---	---	---
	CT8	---	---	---	1.64	---	1.26	---	---	---	---	---	---	---
	CT10	---	---	1.41	2.05	---	1.37	---	---	---	---	---	---	---
25	CT12	---	---	---	1.15	---	1.24	---	---	---	---	---	---	---
	CT14	---	---	1.46	1.40	---	2.58	---	---	---	---	---	---	---
	CT15	---	---	---	---	---	---	---	1.34 1.62	---	---	---	---	---
	CT16	---	---	---	---	---	---	1.13	1.04 1.05	---	---	---	---	---
	CT17	---	---	---	---	---	---	---	1.16	---	---	---	---	---
30	CT1	---	---	---	---	---	1.09	---	---	---	---	---	---	---
	CT4	---	---	---	---	---	1.22	---	---	---	---	---	---	---
	CT5	---	---	2.14	---	---	---	---	---	---	---	---	---	---
	CT9	---	---	---	---	---	1.52	---	---	---	---	---	---	---
	CT11	---	---	1.29	---	---	---	---	---	---	---	---	---	---
35	A549	---	---	---	---	---	---	1.20	2.17 2.11	---	---	---	---	---

5

10

15

20

25

30

35

Calu-1	---	---	---	---	---	---	---	1.39	---	---	---	---	---
Calu-6	---	---	---	---	---	---	---	1.12	---	---	---	---	---
H441	---	---	---	---	---	---	---	2.06	---	---	---	---	---
H460	---	---	---	---	---	---	---	1.88	---	---	---	---	---
SKMES 1	---	---	---	---	---	---	---	1.90	---	---	---	---	---
SW620	---	---	---	---	---	---	---	2.24	---	---	---	---	---
Colo320	---	---	---	---	---	---	---	2.21 2.24	---	---	---	---	---
HT29	---	---	1.22	---	---	---	---	---	---	---	---	---	---
HCT116	---	---	---	---	---	---	---	2.46 2.66	---	---	---	---	---
LT22	---	---	---	1.26	1.07	---	---	---	---	2.69	---	---	---
HF- 000716	---	---	---	---	---	---	---	2.63 2.73	---	---	---	---	---
HF- 000733	---	---	---	---	---	---	---	2.58 2.71 1.39	---	---	---	---	---
HF- 000611	---	---	---	---	---	---	---	4.99	---	---	---	---	---
HF- 000539	2.33	---	---	---	---	---	---	3.13 2.55	---	---	1.49	---	---
HF- 000575	---	---	---	---	---	---	---	1.32	---	---	---	---	---
HF- 000698	---	---	---	---	---	---	---	---	---	---	1.09	---	---
HF- 000545	---	---	---	---	---	---	---	1.59 1.68	---	---	1.11	---	---
HF- 000631	---	---	---	---	---	---	---	1.37	---	---	1.27	---	---
HF- 000840	1.71	---	---	---	---	---	---	3.63	---	---	1.97	1.39	1.11
HF- 000842	---	---	---	---	---	---	---	1.99	---	---	---	1.24	---
HF- 000795	1.13	---	---	---	---	---	---	---	---	---	---	1.01	1.32
HF- 001294	---	---	---	---	---	---	---	---	---	---	---	1.50	---
HF- 001296	---	---	---	---	---	---	---	---	---	---	---	2.88	1.51
HF- 001299	---	---	---	---	---	---	---	---	---	---	---	1.37	---

PRO1265

PRO1265 (DNA60764-1533) was also reexamined along with selected tumors from the above initial screen with framework mapping. Table 9 indicates the chromosomal mapping of the framework markers that were used in the present example. The framework markers are located approximately every 20 megabases and were used to control aneuploidy.

5 PRO1265 was also reexamined with epicenter mapping. The markers indicated in Table 10 are located in close proximity (in the genome) to DNA60764-1533, and are used to assess the relative amplification in the immediate vicinity of Chromosome 19 wherein the molecule is located. The distance between individual markers is measured in centirays (cR), which is a radiation breakage unit approximately equal to a 1% chance of a breakage between two markers. One cR is very roughly equivalent to 20 kilobases. The marker SHGC-33698
10 is closest to DNA60764-1533.

Table 9
Framework Markers Along Chromosome 19

Map Position on Chromosome 19	Stanford Human Genome Center Marker Name
S12	AFMa107xc9
S50	SHGC-31335
S105	SHGC-34102
S155	SHGC-16175

Table 10
Epicenter Markers Along Chromosome 19 used for DNA60764-1533

Map Position on Chromosome 19	Stanford Human Genome Center Marker Name	Distance to next Marker (cR)
DNA34353	—	maps to S158
DNA40620	—	maps to S160
DNA54002	—	maps to S160
S160	SHGC-34723	21
DNA60764	—	—
S161	SHGC-30929	15
S162	SHGC-10328	17
S163	AFMa115wg5	—

The ΔC_t values of the above described framework markers along Chromosome 19 relative to PRO1265 are indicated for selected tumors in Table 11.

Table 11

Amplification of framework markers relative to DNA60764-1533 (ΔC_t)

Tumor	Framework Markers				
	S12	DNA60764-1533	S50	S105	S155
LT1	0.16	0.06	-0.42	0.11	-1.56
LT1a	0.05	-0.27	0.17	0.40	0.00
LT2	0.48	0.41	0.52	0.13	-0.36
LT3	0.27	0.83	0.11	0.50	1.04
LT4	0.48	0.67	0.20	0.56	-0.35
LT6	0.72	0.74	0.32	0.35	0.24
LT7	0.82	0.85	0.95	0.95	0.75
LT9	0.72	0.61	0.19	0.64	-0.35
LT10	0.82	0.98	0.62	0.53	0.32
LT11	0.13	0.25	0.55	-0.34	0.70
LT12	0.04	0.60	0.21	-0.17	2.17
LT13	-0.06	0.57	-0.30	-0.05	2.24
LT15	-0.03	-0.77	0.12	-0.04	3.51
LT16	0.46	1.37	0.51	0.23	3.32
LT17	0.37	0.74	0.21	0.22	1.02
LT18	0.39	0.57	0.11	0.16	0.52
LT22	0.79	0.76	-0.05	0.16	0.59

DISCUSSION AND CONCLUSION:PRO1269 (DNA66520-1536):

The ΔC_t values for DNA66520-1536 in a variety of tumors are reported above. A ΔC_t of >1 was typically used as the threshold value for amplification scoring, as this represents a doubling of gene copy. The above data indicates that significant amplification of nucleic acid DNA66520-1536 encoding PRO1269 occurred in primary lung tumors: LT15, LT16 and LT17. Because amplification of DNA66520-1536 occurs in various lung tumors, it is highly probable to play a significant role in tumor formation or growth. As a result, antagonists (*e.g.*, antibodies) directed against the protein encoded by DNA66520-1536 (PRO1269) would be expected to have utility in cancer therapy.

PRO1410 (DNA68874-1622):

The ΔC_t values for DNA68874-1622 in a variety of tumors are reported above. A ΔC_t of >1 was typically used as the threshold value for amplification scoring, as this represents a doubling of gene copy. the

above data indicates that significant amplification of nucleic acid DNA68874-1622 encoding PRO1410 occurred: (1) in primary lung tumors: LT13, LT15 and LT16; (2) in primary colon tumors: CT2, CT3, CT5, CT10, CT11, and CT14; and (3) in colon cell line HT29. Because amplification of DNA68874-1622 occurs in various lung and colon tumors, it is highly probable to play a significant role in tumor formation or growth. As a result, antagonists (*e.g.*, antibodies) directed against the protein encoded by DNA68874-1622 (PRO1410) would be expected to have utility in cancer therapy.

PRO1755 (DNA76396-1698):

The ΔC_t values for DNA76396-1698 in a variety of tumors are reported above. A ΔC_t of >1 was typically used as the threshold value for amplification scoring, as this represents a doubling of gene copy. The above data indicates that significant amplification of nucleic acid DNA76396-1698 encoding PRO1755 occurred: (1) in primary lung tumors: LT16, LT18 and LT22; and (2) in primary colon tumors: CT2, CT8, CT10, CT12, and CT14. Because amplification of DNA76396-1698 occurs in various lung and colon tumors, it is highly probable to play a significant role in tumor formation or growth. As a result, antagonists (*e.g.*, antibodies) directed against the protein encoded by DNA76396-1698 (PRO1755) would be expected to have utility in cancer therapy.

PRO1780 (DNA71169-1709):

The ΔC_t values for DNA71169-1709 in a variety of tumors are reported above. A ΔC_t of >1 was typically used as the threshold value for amplification scoring, as this represents a doubling of gene copy. The above data indicates that significant amplification of nucleic acid DNA71169-1709 encoding PRO1780 occurred in primary lung tumors: LT4, LT7 and LT22. Because amplification of DNA71169-1709 occurs in various lung tumors, it is highly probable to play a significant role in tumor formation or growth. As a result, antagonists (*e.g.*, antibodies) directed against the protein encoded by DNA71169-1709 (PRO1780) would be expected to have utility in cancer therapy.

PRO1788 (DNA77652-2505):

The ΔC_t values for DNA77652-2505 in a variety of tumors are reported above. A ΔC_t of >1 was typically used as the threshold value for amplification scoring, as this represents a doubling of gene copy. The above data indicates that significant amplification of nucleic acid DNA77652-2505 encoding PRO1788 occurred in primary colon tumors: CT1, CT3, CT4, CT8, CT9, CT10, CT12, and CT14. Because amplification of DNA77652-2505 occurs in various colon tumors, it is highly probable to play a significant role in tumor formation or growth. As a result, antagonists (*e.g.*, antibodies) directed against the protein encoded by DNA77652-2505 (PRO1788) would be expected to have utility in cancer therapy.

PRO1295 (DNA59218-1559):

The ΔC_t values for DNA59218-1559 in a variety of tumors are reported above. A ΔC_t of >1 was typically used as the threshold value for amplification scoring, as this represents a doubling of gene copy. The above data indicates that significant amplification of nucleic acid DNA59218-1559 encoding PRO1295 occurred: (1) in primary lung tumors: HF-000631 and HF-000840; (2) colon tumor centers: HF-000539 and HF-000698;

and (3) in breast tumor center HF-000545. Because amplification of DNA59218-1559 occurs in various tumors, it is highly probable to play a significant role in tumor formation or growth. As a result, antagonists (*e.g.*, antibodies) directed against the protein encoded by DNA59218-1559 (PRO1295) would be expected to have utility in cancer therapy.

5 PRO1293 (DNA60618-1557):

The ΔC_t values for DNA60618-1557 in a variety of tumors are reported above. A ΔC_t of > 1 was typically used as the threshold value for amplification scoring, as this represents a doubling of gene copy. The above data indicates that significant amplification of nucleic acid DNA60618-1557 encoding PRO1293 occurred: (1) in primary lung tumor HF-000840; and (2) in colon tumor centers: HF-000539 and HF-000795. Because amplification of DNA60618-1557 occurs in various lung and colon tumors, it is highly probable to play a significant role in tumor formation or growth. As a result, antagonists (*e.g.*, antibodies) directed against the protein encoded by DNA60618-1557 (PRO1293) would be expected to have utility in cancer therapy.

15 PRO1303 (DNA65409-1566):

The ΔC_t values for DNA65409-1566 in a variety of tumors are reported above. A ΔC_t of > 1 was typically used as the threshold value for amplification scoring, as this represents a doubling of gene copy. The above data indicates that significant amplification of nucleic acid DNA65409-1566 encoding PRO1303 occurred: (1) in primary lung tumors: LT13, LT15 and LT16; (2) in lung cell line A549; and (3) in colon tumor CT16. Because amplification of DNA65409-1566 occurs in various tumors, it is highly probable to play a significant role in tumor formation or growth. As a result, antagonists (*e.g.*, antibodies) directed against the protein encoded by DNA65409-1566 (PRO1566) would be expected to have utility in cancer therapy.

20 PRO1555 (DNA73744-1665):

The ΔC_t values for DNA73744-1665 in a variety of tumors are reported above. A ΔC_t of > 1 was typically used as the threshold value for amplification scoring, as this represents a doubling of gene copy. The above data indicates that significant amplification of nucleic acid DNA73744-1665 encoding PRO1555 occurred: (1) in primary lung tumors: LT13, LT15, LT16, HF-000631, HF-000840, and HF-000842; (2) in lung cell lines: A549, Calu-1, Calu-6, H441, H460, and SKMES1; (3) in primary colon tumors: CT15, CT16, CT17, and colon tumor centers HF-000539 and HF-000575; (4) in colon cell lines: SW620, Colo320 and HCT116; (5) in breast tumor center HF-000545; (6) in kidney tumor center HF-000611; and (7) in testis tumor margin HF-000716 and testis tumor center HF-000733. Because amplification of DNA73744-1665 occurs in various tumors, it is highly probable to play a significant role in tumor formation or growth. As a result, antagonists (*e.g.*, antibodies) directed against the protein encoded by DNA73744-1665 (PRO1555) would be expected to have utility in cancer therapy.

35 PRO1265 (DNA60764-1533):

The ΔC_t values for DNA60764-1533 in a variety of lung tumors are reported above. A ΔC_t value of > 1 was typically used as the threshold value for amplification scoring, as this represents a doubling of gene copy. The above data indicates that significant amplification of DNA60765-1533 occurred in primary lung

tumors LT3, LT12, LT13, LT15, LT16 and LT17. The ΔC_t values of these hits are 1.03, 2.17, 2.24, 3.51, 3.32 and 1.02. This represents an increase in gene copy of approximately 2.04, 4.50, 4.72, 11.39, 9.99 and 2.03.

Amplification has also been confirmed framework mapping for DNA60764-1533 in LT16. The reported ΔC_t value was 1.37, which represents a 2.58 fold increase in gene copy relative to normal tissue. Epicenter mapping has also confirmed amplification of DNA60764-1533 in LT12, LT13, LT15, LT16, CT1, CT4, CT5, CT7 and CT11. These tumors report ΔC_t values of 2.35, 2.37, 3.88, 3.32 in the lung tumors and 1.74, 1.86, 3.28, 1.29 and 2.32 in the colon tumors. Relative to normal tissue, this represents an increase in gene copy of approximately 5.10, 5.17, 14.72 and 9.98 in the lung tumors and 3.34, 3.63, 9.71, 2.45 and 4.99 in the colon tumors.

In contrast, the amplification of the closest known framework markers, epicenter markers and the comparison sequences does not occur to a greater extent than that of DNA60764-1533. This strongly suggests that DNA60764-1533 is the gene responsible for the amplification of the particular region in Chromosome 19. Because amplification of DNA60764-1533 occurs in various lung and colon tumors, it is highly probably to play a significant role in tumor formation or growth. As a result, antagonists (*e.g.*, antibodies) directed against the protein encoded by DNA60764-1533 would be expected to have utility in cancer therapy.

PRO1317 (DNA71166-1685):

The ΔC_t values for DNA71166-1685 in a variety of tumors are reported above. A ΔC_t of >1 was typically used as the threshold value for amplification scoring, as this represents a doubling of gene copy. The above data indicates that significant amplification of nucleic acid DNA71166-1685 encoding PRO1317 occurred in primary lung tumors LT1, LT1a, LT9, LT10, LT15, LT17 and LT22. Because amplification of DNA71166-1685 occurs in various tumors, it is likely associated with tumor formation and/or growth. As a result, antagonists (*e.g.*, antibodies) directed against PRO1317 would be expected to be useful in cancer therapy.

Summary

Because amplification of the various DNA's as described above occurs in various tumors, they are likely associated with tumor formation and/or growth. As a result, antagonists (*e.g.*, antibodies) directed against these polypeptides would be expected to be useful in cancer therapy.

EXAMPLE 144: Stimulatory Activity in Mixed Lymphocyte Reaction (MLR) Assay (Assay 24)

This example shows that certain polypeptides of the invention are active as a stimulator of the proliferation of stimulated T-lymphocytes. Compounds which stimulate proliferation of lymphocytes are useful therapeutically where enhancement of an immune response is beneficial. A therapeutic agent may take the form of antagonists of the polypeptide of the invention, for example, murine-human chimeric, humanized or human antibodies against the polypeptide.

The basic protocol for this assay is described in Current Protocols in Immunology, unit 3.12; edited by J E Coligan, A M Kruisbeek, D H Marglies, E M Shevach, W Strober, National Institutes of Health, Published by John Wiley & Sons, Inc.

More specifically, in one assay variant, peripheral blood mononuclear cells (PBMC) are isolated from mammalian individuals, for example a human volunteer, by leukopheresis (one donor will supply stimulator PBMCs, the other donor will supply responder PBMCs). If desired, the cells are frozen in fetal bovine serum and DMSO after isolation. Frozen cells may be thawed overnight in assay media (37°C, 5% CO₂) and then washed and resuspended to 3x10⁶ cells/ml of assay media (RPMI; 10% fetal bovine serum, 1% penicillin/streptomycin, 1% glutamine, 1% HEPES, 1% non-essential amino acids, 1% pyruvate). The stimulator PBMCs are prepared by irradiating the cells (about 3000 Rads).

The assay is prepared by plating in triplicate wells a mixture of:

100:1 of test sample diluted to 1% or to 0.1%,

50 :1 of irradiated stimulator cells, and

50 :1 of responder PBMC cells.

100 microliters of cell culture media or 100 microliter of CD4-IgG is used as the control. The wells are then incubated at 37°C, 5% CO₂ for 4 days. On day 5, each well is pulsed with tritiated thymidine (1.0 mCi/well; Amersham). After 6 hours the cells are washed 3 times and then the uptake of the label is evaluated.

In another variant of this assay, PBMCs are isolated from the spleens of Balb/c mice and C57B6 mice. The cells are teased from freshly harvested spleens in assay media (RPMI; 10% fetal bovine serum, 1% penicillin/streptomycin, 1% glutamine, 1% HEPES, 1% non-essential amino acids, 1% pyruvate) and the PBMCs are isolated by overlaying these cells over Lympholyte M (Organon Teknika), centrifuging at 2000 rpm for 20 minutes, collecting and washing the mononuclear cell layer in assay media and resuspending the cells to 1x10⁷ cells/ml of assay media. The assay is then conducted as described above.

Positive increases over control are considered positive with increases of greater than or equal to 180% being preferred. However, any value greater than control indicates a stimulatory effect for the test protein.

The following PRO polypeptides tested positive in this assay: PRO1246 and PRO1343.

EXAMPLE 145: Mouse Kidney Mesangial Cell Proliferation Assay (Assay 92)

This assay shows that certain polypeptides of the invention act to induce proliferation of mammalian kidney mesangial cells and, therefore, are useful for treating kidney disorders associated with decreased mesangial cell function such as Berger disease or other nephropathies associated with Schönlein-Henoch purpura, celiac disease, dermatitis herpetiformis or Crohn disease. The assay is performed as follows. On day one, mouse kidney mesangial cells are plated on a 96 well plate in growth media (3:1 mixture of Dulbecco's modified Eagle's medium and Ham's F12 medium, 95% fetal bovine serum, 5% supplemented with 14 mM HEPES) and grown overnight. On day 2, PRO polypeptides are diluted at 2 concentrations (1% and 0.1%) in serum-free medium and added to the cells. Control samples are serum-free medium alone. On day 4, 20μl of the Cell Titer 96 Aqueous one solution reagent (Progema) was added to each well and the colorimetric reaction was allowed to proceed for 2 hours. The absorbance (OD) is then measured at 490 nm. A positive in the assay is anything that gives an absorbance reading which is at least 15% above the control reading.

The following polypeptide tested positive in this assay: PRO1265, PRO1244 and PRO1382.

EXAMPLE 146: Induction of Pancreatic β-Cell Precursor Differentiation (Assay 89)

This assay shows that certain polypeptides of the invention act to induce differentiation of pancreatic

β -cell precursor cells into mature pancreatic β -cells and, therefore, are useful for treating various insulin deficient states in mammals, including diabetes mellitus. The assay is performed as follows. The assay uses a primary culture of mouse fetal pancreatic cells and the primary readout is an alteration in the expression of markers that represent either β -cell precursors or mature β -cells. Marker expression is measured by real time quantitative PCR (RTQ-PCR); wherein the marker being evaluated is insulin.

5 The pancreata are dissected from E14 embryos (CD1 mice). The pancreata are then digested with collagenase/dispase in F12/DMEM at 37°C for 40 to 60 minutes (collagenase/dispase, 1.37 mg/ml, Boehringer Mannheim, #1097113). The digestion is then neutralized with an equal volume of 5% BSA and the cells are washed once with RPMI1640. At day 1, the cells are seeded into 12-well tissue culture plates (pre-coated with laminin, 20 μ g/ml in PBS, Boehringer Mannheim, #124317). Cells from pancreata from 1-2 embryos are distributed per well. The culture medium for this primary culture is 14F/1640. At day 2, the media is removed and the attached cells washed with RPMI/1640. Two mls of minimal media are added in addition to the protein to be tested. At day 4, the media is removed and RNA prepared from the cells and marker expression analyzed by real time quantitative RT-PCR. A protein is considered to be active in the assay if it increases the expression of the relevant β -cell marker as compared to untreated controls.

15 14F/1640 is RPMI1640 (Gibco) plus the following:

group A 1:1000

group B 1:1000

recombinant human insulin 10 μ g/ml

Aprotinin (50 μ g/ml) 1:2000 (Boehringer manheim #981532)

20 Bovine pituitary extract (BPE) 60 μ g/ml

Gentamycin 100 ng/ml

Group A : (in 10ml PBS)

Transferrin, 100mg (Sigma T2252)

Epidermal Growth Factor, 100 μ g (BRL 100004)

25 Triiodothyronine, 10 μ l of 5x10⁻⁶ M (Sigma T5516)

Ethanolamine, 100 μ l of 10⁻¹ M (Sigma E0135)

Phosphoethanolamine, 100 μ l of 10⁻¹ M (Sigma P0503)

Selenium, 4 μ l of 10⁻¹ M (Aesar #12574)

Group C : (in 10ml 100% ethanol)

30 Hydrocortisone, 2 μ l of 5X10⁻³ M (Sigma #H0135)

Progesterone, 100 μ l of 1X10⁻³ M (Sigma #P6149)

Forskolin, 500 μ l of 20mM (Calbiochem #344270)

Minimal media:

35 RPMI 1640 plus transferrin (10 μ g/ml), insulin (1 μ g/ml), gentamycin (100 ng/ml), aprotinin (50 μ g/ml) and BPE (15 μ g/ml).

Defined media:

RPMI 1640 plus transferrin (10 μ g/ml), insulin (1 μ g/ml), gentamycin (100 ng/ml) and aprotinin (50 μ g/ml).

The following polypeptides were positive in this assay: PRO1275 and PRO1474.

EXAMPLE 147: Fetal Hemoglobin Induction in an Erythroblastic Cell Line (Assay 107)

This assay is useful for screening PRO polypeptides for the ability to induce the switch from adult hemoglobin to fetal hemoglobin in an erythroblastic cell line. Molecules testing positive in this assay are expected to be useful for therapeutically treating various mammalian hemoglobin-associated disorders such as the various thalassemias. The assay is performed as follows. Erythroblastic cells are plated in standard growth medium at 1000 cells/well in a 96 well format. PRO polypeptides are added to the growth medium at a concentration of 0.2% or 2% and the cells are incubated for 5 days at 37°C. As a positive control, cells are treated with 100µM hemin and as a negative control, the cells are untreated. After 5 days, cell lysates are prepared and analyzed for the expression of gamma globin (a fetal marker). A positive in the assay is a gamma globin level at least 2-fold above the negative control.

The following polypeptides tested positive in this assay: PRO1478, PRO1265, PRO1412, PRO1279, PRO1304, PRO1306, PRO1418, PRO1410 and PRO1575.

EXAMPLE 148: Detection of Polypeptides That Affect Glucose and/or FFA Uptake in Skeletal Muscle (Assay 106)

This assay is designed to determine whether PRO polypeptides show the ability to affect glucose or FFA uptake by skeletal muscle cells. PRO polypeptides testing positive in this assay would be expected to be useful for the therapeutic treatment of disorders where either the stimulation or inhibition of glucose uptake by skeletal muscle would be beneficial including, for example, diabetes or hyper- or hypo-insulinemia.

In a 96 well format, PRO polypeptides to be assayed are added to primary rat differentiated skeletal muscle, and allowed to incubate overnight. Then fresh media with the PRO polypeptide and +/- insulin are added to the wells. The sample media is then monitored to determine glucose and FFA uptake by the skeletal muscle cells. The insulin will stimulate glucose and FFA uptake by the skeletal muscle, and insulin in media without the PRO polypeptide is used as a positive control, and a limit for scoring. As the PRO polypeptide being tested may either stimulate or inhibit glucose and FFA uptake, results are scored as positive in the assay if greater than 1.5 times or less than 0.5 times the insulin control.

The following PRO polypeptides tested positive as either stimulators or inhibitors of glucose and/or FFA uptake in this assay: PRO1130, PRO1275, PRO1418, PRO1555 and PRO1787.

EXAMPLE 149: Detection of PRO Polypeptides That Affect Glucose or FFA Uptake by Primary Rat Adipocytes (Assay 94)

This assay is designed to determine whether PRO polypeptides show the ability to affect glucose or FFA uptake by adipocyte cells. PRO polypeptides testing positive in this assay would be expected to be useful for the therapeutic treatment of disorders where either the stimulation or inhibition of glucose uptake by adipocytes would be beneficial including, for example, obesity, diabetes or hyper- or hypo-insulinemia.

In a 96 well format, PRO polypeptides to be assayed are added to primary rat adipocytes, and allowed to incubate overnight. Samples are taken at 4 and 16 hours and assayed for glycerol, glucose and FFA uptake. After the 16 hour incubation, insulin is added to the media and allowed to incubate for 4 hours. At this time, a sample is taken and glycerol, glucose and FFA uptake is measured. Media containing insulin without the PRO polypeptide is used as a positive reference control. As the PRO polypeptide being tested may either stimulate or inhibit glucose and FFA uptake, results are scored as positive in the assay if greater than 1.5 times or less than 0.5 times the insulin control.

The following PRO polypeptides tested positive as stimulators of glucose and/or FFA uptake in this assay: PRO1265, PRO1283, PRO1279, PRO1303, PRO1306, PRO1325, PRO1565 and PRO1567.

The following PRO polypeptides tested positive as inhibitors of glucose and/or FFA uptake in this assay: PRO1194, PRO1190, PRO1326, PRO1343, PRO1480, PRO1474, PRO1575 and PRO1760.

EXAMPLE 150: Chondrocyte Re-differentiation Assay (Assay 110)

This assay shows that certain polypeptides of the invention act to induce redifferentiation of chondrocytes, therefore, are expected to be useful for the treatment of various bone and/or cartilage disorders such as, for example, sports injuries and arthritis. The assay is performed as follows. Porcine chondrocytes are isolated by overnight collagenase digestion of articular cartilage of metacarpophalangeal joints of 4-6 month old female pigs. The isolated cells are then seeded at 25,000 cells/cm² in Ham F-12 containing 10% FBS and 4 µg/ml gentamycin. The culture media is changed every third day and the cells are then seeded in 96 well plates at 5,000 cells/well in 100µl of the same media without serum and 100 µl of the test PRO polypeptide, 5 nM staurosporin (positive control) or medium alone (negative control) is added to give a final volume of 200 µl/well. After 5 days of incubation at 37°C, a picture of each well is taken and the differentiation state of the chondrocytes is determined. A positive result in the assay occurs when the redifferentiation of the chondrocytes is determined to be more similar to the positive control than the negative control.

The following polypeptide tested positive in this assay: PRO1265, PRO1250, PRO1430, PRO1356, PRO1275, PRO1274, PRO1286, PRO1273, PRO1283, PRO1279, PRO1306, PRO1325, PRO1343, PRO1418, PRO1565, PRO1474, PRO1787, PRO1556 and PRO1801.

EXAMPLE 151: Induction of Pancreatic β-Cell Precursor Proliferation (Assay 117)

This assay shows that certain polypeptides of the invention act to induce an increase in the number of pancreatic β-cell precursor cells and, therefore, are useful for treating various insulin deficient states in mammals, including diabetes mellitus. The assay is performed as follows. The assay uses a primary culture of mouse fetal pancreatic cells and the primary readout is an alteration in the expression of markers that represent either β-cell precursors or mature β-cells. Marker expression is measured by real time quantitative PCR (RTQ-PCR); wherein the marker being evaluated is a transcription factor called Pdx1.

The pancreata are dissected from E14 embryos (CD1 mice). The pancreata are then digested with collagenase/dispase in F12/DMEM at 37°C for 40 to 60 minutes (collagenase/dispase, 1.37 mg/ml, Boehringer Mannheim, #1097113). The digestion is then neutralized with an equal volume of 5% BSA and the cells are washed once with RPMI1640. At day 1, the cells are seeded into 12-well tissue culture plates (pre-coated with laminin, 20µg/ml in PBS, Boehringer Mannheim, #124317). Cells from pancreata from 1-2 embryos are

distributed per well. The culture medium for this primary culture is 14F/1640. At day 2, the media is removed and the attached cells washed with RPMI/1640. Two mls of minimal media are added in addition to the protein to be tested. At day 4, the media is removed and RNA prepared from the cells and marker expression analyzed by real time quantitative RT-PCR. A protein is considered to be active in the assay if it increases the expression of the relevant β -cell marker as compared to untreated controls.

5 14F/1640 is RPMI1640 (Gibco) plus the following:

group A 1:1000

group B 1:1000

recombinant human insulin 10 μ g/ml

Aprotinin (50 μ g/ml) 1:2000 (Boehringer manheim #981532)

10 Bovine pituitary extract (BPE) 60 μ g/ml

Gentamycin 100 ng/ml

Group A : (in 10ml PBS)

Transferrin, 100mg (Sigma T2252)

Epidermal Growth Factor, 100 μ g (BRL 100004)

15 Triiodothyronine, 10 μ l of 5×10^{-6} M (Sigma T5516)

Ethanolamine, 100 μ l of 10^{-1} M (Sigma E0135)

Phosphoethalamine, 100 μ l of 10^{-1} M (Sigma P0503)

Selenium, 4 μ l of 10^{-1} M (Aesar #12574)

Group C : (in 10ml 100% ethanol)

20 Hydrocortisone, 2 μ l of 5×10^{-3} M (Sigma #H0135)

Progesterone, 100 μ l of 1×10^{-3} M (Sigma #P6149)

Forskolin, 500 μ l of 20mM (Calbiochem #344270)

Minimal media:

25 RPMI 1640 plus transferrin (10 μ g/ml), insulin (1 μ g/ml), gentamycin (100 ng/ml), aprotinin (50 μ g/ml) and BPE (15 μ g/ml).

Defined media:

RPMI 1640 plus transferrin (10 μ g/ml), insulin (1 μ g/ml), gentamycin (100 ng/ml) and aprotinin (50 μ g/ml).

The following polypeptides tested positive in this assay: PRO1382 and PRO1561.

30

EXAMPLE 152: Proliferation of Rat Utricular Supporting Cells (Assay 54)

35 This assay shows that certain polypeptides of the invention act as potent mitogens for inner ear supporting cells which are auditory hair cell progenitors and, therefore, are useful for inducing the regeneration of auditory hair cells and treating hearing loss in mammals. The assay is performed as follows. Rat UEC-4 utricular epithelial cells are aliquoted into 96 well plates with a density of 3000 cells/well in 200 μ l of serum-containing medium at 33°C. The cells are cultured overnight and are then switched to serum-free medium at 37°C. Various dilutions of PRO polypeptides (or nothing for a control) are then added to the cultures and the cells are incubated for 24 hours. After the 24 hour incubation, 3 H-thymidine (1 μ Ci/well) is added and the cells

are then cultured for an additional 24 hours. The cultures are then washed to remove unincorporated radiolabel, the cells harvested and Cpm per well determined. Cpm of at least 30% or greater in the PRO polypeptide treated cultures as compared to the control cultures is considered a positive in the assay.

The following polypeptides tested positive in this assay: PRO1340.

5 EXAMPLE 153: Chondrocyte Proliferation Assay (Assay 111)

This assay is designed to determine whether PRO polypeptides of the present invention show the ability to induce the proliferation and/or redifferentiation of chondrocytes in culture. PRO polypeptides testing positive in this assay would be expected to be useful for the therapeutic treatment of various bone and/or cartilage disorders such as, for example, sports injuries and arthritis.

10 Porcine chondrocytes are isolated by overnight collagenase digestion of articular cartilage of the metacarpophalangeal joint of 4-6 month old female pigs. The isolated cells are then seeded at 25,000 cells/cm² in Ham F-12 containing 10% FBS and 4 µg/ml gentamycin. The culture media is changed every third day and the cells are reseeded to 25,000 cells/cm² every five days. On day 12, the cells are seeded in 96 well plates at 5,000 cells/well in 100µl of the same media without serum and 100 µl of either serum-free medium (negative control), staurosporin (final concentration of 5 nM; positive control) or the test PRO polypeptide are added to give a final volume of 200 µl/well. After 5 days at 37°C, 20 µl of Alamar blue is added to each well and the plates are incubated for an additional 3 hours at 37°C. The fluorescence is then measured in each well (Ex:530 nm; Em: 590 nm). The fluorescence of a plate containing 200 µl of the serum-free medium is measured to obtain the background. A positive result in the assay is obtained when the fluorescence of the PRO polypeptide treated sample is more like that of the positive control than the negative control.

20 The following PRO polypeptides tested positive in this assay: PRO1265, PRO1412, PRO1347, PRO1279, PRO1410 and PRO1474.

EXAMPLE 154: Inhibition of Heart Neonatal Hypertrophy Induced by LIF+ET-1 (Assay 74)

25 This assay is designed to determine whether PRO polypeptides of the present invention show the ability to inhibit neonatal heart hypertrophy induced by LIF and endothelin-1 (ET-1). A test compound that provides a positive response in the present assay would be useful for the therapeutic treatment of cardiac insufficiency diseases or disorders characterized or associated with an undesired hypertrophy of the cardiac muscle.

30 Cardiac myocytes from 1-day old Harlan Sprague Dawley rats (180 µl at 7.5 x 10⁴/ml, serum <0.1, freshly isolated) are introduced on day 1 to 96-well plates previously coated with DMEM/F12 + 4%FCS. Test PRO polypeptide samples or growth medium alone (negative control) are then added directly to the wells on day 2 in 20 µl volume. LIF + ET-1 are then added to the wells on day 3. The cells are stained after an additional 2 days in culture and are then scored visually the next day. A positive in the assay occurs when the PRO polypeptide treated myocytes are visually smaller on the average or less numerous than the untreated myocytes.

35 The following PRO polypeptides tested positive in this assay: PRO1760.

EXAMPLE 155: Tissue Expression Distribution

Oligonucleotide probes were constructed from some of the PRO polypeptide-encoding nucleotide sequences shown in the accompanying figures for use in quantitative PCR amplification reactions. The

oligonucleotide probes were chosen so as to give an approximately 200-600 base pair amplified fragment from the 3' end of its associated template in a standard PCR reaction. The oligonucleotide probes were employed in standard quantitative PCR amplification reactions with cDNA libraries isolated from different human adult and/or fetal tissue sources and analyzed by agarose gel electrophoresis so as to obtain a quantitative determination of the level of expression of the PRO polypeptide-encoding nucleic acid in the various tissues tested. Knowledge of the expression pattern or the differential expression of the PRO polypeptide-encoding nucleic acid in various different human tissue types provides a diagnostic marker useful for tissue typing, with or without other tissue-specific markers, for determining the primary tissue source of a metastatic tumor, and the like. These assays provided the following results.

	<u>DNA Molecule</u>	<u>Tissues With Significant Expression</u>	<u>Tissues Lacking Significant Expression</u>
10	DNA19902-1669	HUVEC cells, colon tumor	dendritic cells, lymphoblast cells, heart
	DNA23322-1393	uterus, colon tumor, prostate	cartilage
	DNA26846-1397	lymphoblast cells	uterus, heart, cartilage
	DNA56107-1415	spleen, substantia nigra, colon tumor	cartilage
	DNA56406-1704	THP-1 macrophages, uterus, spleen	endothelial cells, prostate, cartilage
15	DNA56529-1647	liver, kidney, brain	adenocarcinoma, lung, bone marrow
	DNA56862-1343	endothelial cells, substantia nigra	colon tumor, lymphoblast cells, uterus
		hippocampus	
	DNA57254-1477	kidney	lung, placenta, brain
	DNA58730-1607	bone marrow, kidney	lung, brain
20	DNA58732-1650	lung, bone marrow	brain, liver
	DNA58828-1519	adenocarcinoma	lung, retina, small intestine
	DNA58852-1637	uterus	colon tumor, heart, brain
	DNA59212-1627	uterus	prostate, cartilage, heart
	DNA59219-1613	spleen, dendrocytes, prostate, uterus	substantia nigra, colon tumor, heart
25	DNA59817-1703	bone marrow	lung, small intestine, placenta
	DNA60278-1530	prostate, colon tumor	uterus, cartilage
	DNA60608-1577	kidney, bone marrow	breast carcinoma, small intestine, lung
	DNA60611-1524	breast carcinoma	lung, small intestine, retina
	DNA60740-1615	breast carcinoma, adenocarcinoma	lung, small intestine, brain
30	DNA62809-1531	THP-1 macrophages	uterus, spleen, brain, colon tumor
	DNA62815-1576	colon tumor, uterus, prostate	spleen, brain, heart, cartilage
	DNA62845-1684	liver, bone marrow	adenocarcinoma, lung, brain
	DNA64849-1604	kidney	lung, pancreas, liver, thyroid
	DNA64863-1573	lung, brain, kidney, bone marrow	liver, pancreas
35	DNA64881-1602	uterus	heart, spleen, brain, endothelial cells
	DNA64902-1667	uterus	prostate, brain, heart, spleen
	DNA64952-1568	lung, brain	pancreas
	DNA65403-1565	spleen, dendrocytes, THP-1 macrophages	endothelial cells, colon tumor, lymphoblasts
40	DNA65408-1578	prostate, spleen, dendrocytes	uterus, heart, substantia nigra
	DNA65423-1595	testis	breast carcinoma, retina, small intestine
	DNA66512-1564	heart, uterus, prostate, cartilage	endothelial cells
	DNA66519-1535	dendrocytes, lymphoblasts, uterus	substantia nigra, prostate, spleen
	DNA66521-1583	uterus, heart, hippocampus	cartilage, dendrocytes, spleen
45	DNA66658-1584	prostate, uterus, hippocampus, spleen	colon tumor, cartilage, heart
	DNA66672-1586	spleen	heart, prostate, brain, uterus
	DNA66674-1599	uterus, prostate	heart, brain, spleen, cartilage, colon tumor
50	DNA68836-1656	kidney	lung, brain, bone marrow, liver
	DNA68871-1638	uterus, colon tumor, prostate	heart, cartilage, brain, spleen
	DNA68880-1676	heart, endothelial cells, brain, uterus	THP-1 macrophages
	DNA68885-1678	uterus, colon tumor, prostate	brain, heart, cartilage, endothelial cells
	DNA71180-1655	brain	lung, bone marrow, liver, kidney

	DNA71184-1634 DNA71234-1651 DNA71277-1636	breast carcinoma, bone marrow, testis kidney, bone marrow prostate, cartilage, heart, uterus	brain, adrenal gland lung, brain, placenta colon tumor, substantia nigra, endothelial cells
5	DNA71286-1687 DNA71883-1660 DNA73492-1671	uterus, prostate, brain, cartilage aortic endothelial cells breast carcinoma, aortic endothelial cells bone marrow	heart lung, retina, small intestine, kidney lung, brain, testis
10	DNA73734-1680 DNA73735-1681 DNA73736-1657 DNA73737-1658 DNA73742-1662	prostate, spleen prostate spleen, substantia nigra, hippocampus, cartilage uterus spleen, uterus, prostate	heart, cartilage, brain, uterus brain, heart, cartilage, spleen prostate, heart, uterus, dendrocytes prostate, heart, spleen, cartilage dendrocytes, colon tumor, endothelial cells
15	DNA73746-1654 DNA73760-1672 DNA76393-1664 DNA76398-1699	prostate breast carcinoma endothelial cells, cartilage, uterus hippocampus, prostate, THP-1 macrophages	uterus, heart, brain, cartilage, spleen retina, brain, kidney, liver, testis brain, prostate heart, uterus, spleen, dendrocytes
20	DNA76399-1700 DNA76522-2500 DNA77301-1708 DNA77648-1688	IM-9 lymphoblasts colon tumor brain retina, breast carcinoma, kidney, liver, bone marrow	prostate, spleen, heart, cartilage, uterus uterus, prostate, brain, heart, cartilage lung, small intestine, kidney, liver brain, lung
25	DNA77568-1626 DNA58727-1474	brain HUVEC, dendrocytes, uterus	lung, liver, placenta, heart substantia nigra, hippocampus, prostate, colon tumor
30	DNA61185-1646 DNA61608-1606 DNA66304-1546	colon tumor, HUVEC colon tumor, dendrocytes, spleen, testis prostate, testis	uterus, dendrocytes, substantia nigra substantia nigra, placenta uterus, brain, heart, colon tumor, adrenal gland
35	DNA71213-1659 DNA62812-1594	brain, spleen, HUVEC, colon tumor heart	prostate, uterus, heart, cartilage placenta, testis, uterus, adrenal gland, bone marrow, prostate bone marrow
40	DNA66660-1585 DNA66669-1597 DNA68866-1644 DNA73730-1679	colon tumor, HUVEC, testis, placenta, uterus heart, placenta, adrenal gland, uterus testis, colon tumor, prostate, spleen, testis, adrenal gland, uterus, prostate, uterus	cartilage, testis, colon tumor, HUVEC, bone marrow, prostate, spleen cartilage, adrenal gland, HUVEC, placenta cartilage, colon tumor, heart, placenta, spleen

Deposit of Material

The following materials have been deposited with the American Type Culture Collection, 10801 University Blvd., Manassas, VA 20110-2209, USA (ATCC):

Table 12

	<u>Material</u>	<u>ATCC Dep. No.</u>	<u>Deposit Date</u>
5	DNA19902-1669	203454	November 3, 1998
	DNA26846-1397	203406	October 27, 1998
	DNA56107-1415	203405	October 27, 1998
	DNA56406-1704	203478	November 17, 1998
	DNA56529-1647	203293	September 29, 1998
10	DNA56531-1648	203286	September 29, 1998
	DNA56862-1343	203174	September 1, 1998
	DNA57254-1477	203289	September 29, 1998
	DNA57841-1522	203458	November 3, 1998
	DNA58727-1474	203171	September 1, 1998
15	DNA58730-1607	203221	September 15, 1998
	DNA58732-1650	203290	September 29, 1998
	DNA58828-1519	203172	September 1, 1998
	DNA58852-1637	203271	September 22, 1998
	DNA59212-1627	203245	September 9, 1998
20	DNA59218-1559	203287	September 29, 1998
	DNA59219-1613	203220	September 15, 1998
	DNA59586-1520	203288	September 29, 1998
	DNA59817-1703	203470	November 17, 1998
	DNA60278-1530	203170	September 1, 1998
25	DNA60608-1577	203126	August 18, 1998
	DNA60611-1524	203175	September 1, 1998
	DNA60618-1557	203292	September 29, 1998
	DNA60740-1615	203456	November 3, 1998
	DNA60764-1533	203452	November 10, 1998
30	DNA60775-1532	203173	September 1, 1998
	DNA61185-1646	203464	November 17, 1998
	DNA61608-1606	203239	September 9, 1998
	DNA62808-1326	203358	October 20, 1998
	DNA62809-1531	203237	September 9, 1998
35	DNA62815-1578	203247	September 9, 1998
	DNA62845-1684	203361	October 20, 1998
	DNA64842-1632	203278	September 22, 1998
	DNA64849-1604	203468	November 17, 1998
	DNA64863-1573	203251	September 9, 1998
40	DNA64881-1602	203240	September 9, 1998
	DNA64883-1526	203253	September 9, 1998
	DNA64885-1529	203457	November 3, 1998
	DNA64886-1601	203241	September 9, 1998
	DNA64888-1542	203249	September 9, 1998
45	DNA64889-1541	203250	September 9, 1998
	DNA64897-1628	203216	September 15, 1998
	DNA64902-1667	203317	October 6, 1998
	DNA64903-1553	203223	September 15, 1998
	DNA64905-1558	203233	September 15, 1998
50	DNA64950-1590	203224	September 15, 1998
	DNA64952-1568	203222	September 15, 1998
	DNA65402-1540	203252	September 9, 1998
	DNA65403-1565	203230	September 15, 1998
	DNA65404-1551	203244	September 9, 1998
55	DNA65405-1547	203476	November 17, 1998
	DNA65406-1567	203219	September 15, 1998

	DNA65408-1578	203217	September 15, 1998
	DNA65409-1566	203232	September 15, 1998
	DNA65410-1569	203231	September 15, 1998
	DNA65423-1595	203227	September 15, 1998
	DNA66304-1546	203321	October 6, 1998
5	DNA66511-1411	203228	September 15, 1998
	DNA66512-1564	203218	September 15, 1998
	DNA66519-1535	203236	September 15, 1998
	DNA66520-1536	203226	September 15, 1998
	DNA66521-1583	203225	September 15, 1998
10	DNA66526-1616	203246	September 9, 1998
	DNA66658-1584	203229	September 15, 1998
	DNA66659-1593	203269	September 22, 1998
	DNA66663-1598	203268	September 22, 1998
	DNA66669-1597	203272	September 22, 1998
15	DNA66672-1586	203265	September 22, 1998
	DNA66674-1599	203281	September 22, 1998
	DNA66675-1587	203282	September 22, 1998
	DNA67962-1649	203291	September 29, 1998
	DNA68836-1656	203455	November 3, 1998
20	DNA68864-1629	203276	September 22, 1998
	DNA68866-1644	203283	September 22, 1998
	DNA68871-1638	203280	September 22, 1998
	DNA68874-1622	203277	September 22, 1998
	DNA68880-1676	203319	October 6, 1998
25	DNA68885-1570	203311	October 6, 1998
	DNA71166-1685	203355	October 20, 1998
	DNA71169-1709	203467	November 17, 1998
	DNA71180-1655	203403	October 27, 1998
	DNA71184-1634	203266	September 22, 1998
30	DNA71213-1659	203401	October 27, 1998
	DNA71234-1651	203402	October 27, 1998
	DNA71277-1636	203285	September 22, 1998
	DNA71282-1668	203312	October 6, 1998
	DNA71286-1604	203357	October 20, 1998
35	DNA71883-1660	203475	November 17, 1998
	DNA73401-1633	203273	September 22, 1998
	DNA73492-1671	203324	October 6, 1998
	DNA73727-1673	203459	November 3, 1998
	DNA73730-1679	203320	October 6, 1998
40	DNA73734-1680	203363	October 20, 1998
	DNA73735-1681	203356	October 20, 1998
	DNA73736-1657	203466	November 17, 1998
	DNA73737-1658	203412	October 27, 1998
	DNA73739-1645	203270	September 22, 1998
45	DNA73742-1662	203316	October 6, 1998
	DNA73744-1665	203322	October 6, 1998
	DNA73746-1654	203411	October 27, 1998
	DNA73760-1672	203314	October 6, 1998
	DNA76396-1698	203471	November 17, 1998
50	DNA76398-1699	203474	November 17, 1998
	DNA76399-1700	203472	November 17, 1998
	DNA76401-1683	203360	October 20, 1998
	DNA76510-2504	203477	November 17, 1998
	DNA76522-2500	203469	November 17, 1998
55	DNA76529-1666	203315	October 6, 1998
	DNA76531-1701	203465	November 17, 1998
	DNA76532-1702	203473	November 17, 1998
	DNA76538-1670	203313	October 6, 1998

	DNA76541-1675	203409	October 27, 1998
	DNA77301-1708	203407	October 27, 1998
	DNA77303-2502	203479	November 17, 1998
	DNA77648-1688	203408	October 27, 1998
	DNA77652-2505	203480	November 17, 1998
5	DNA83500-2506	203391	October 29, 1998
	DNA77568-1626	203134	August 18, 1998
	DNA23322-1393	203400	October 27, 1998
	DNA59814-1486	203359	October 20, 1998
	DNA62812-1594	203248	September 9, 1998
10	DNA66660-1585	203279	September 22, 1998
	DNA76393-1664	203323	October 6, 1998

These deposit were made under the provisions of the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purpose of Patent Procedure and the Regulations thereunder (Budapest Treaty). This assures maintenance of a viable culture of the deposit for 30 years from the date of deposit. The deposits will be made available by ATCC under the terms of the Budapest Treaty, and subject to an agreement between Genentech, Inc. and ATCC, which assures permanent and unrestricted availability of the progeny of the culture of the deposit to the public upon issuance of the pertinent U.S. patent or upon laying open to the public of any U.S. or foreign patent application, whichever comes first, and assures availability of the progeny to one determined by the U.S. Commissioner of Patents and Trademarks to be entitled thereto according to 35 USC §122 and the Commissioner's rules pursuant thereto (including 37 CFR §1.14 with particular reference to 886 OG 638).

The assignee of the present application has agreed that if a culture of the materials on deposit should die or be lost or destroyed when cultivated under suitable conditions, the materials will be promptly replaced on notification with another of the same. Availability of the deposited material is not to be construed as a license to practice the invention in contravention of the rights granted under the authority of any government in accordance with its patent laws.

The foregoing written specification is considered to be sufficient to enable one skilled in the art to practice the invention. The present invention is not to be limited in scope by the construct deposited, since the deposited embodiment is intended as a single illustration of certain aspects of the invention and any constructs that are functionally equivalent are within the scope of this invention. The deposit of material herein does not constitute an admission that the written description herein contained is inadequate to enable the practice of any aspect of the invention, including the best mode thereof, nor is it to be construed as limiting the scope of the claims to the specific illustrations that it represents. Indeed, various modifications of the invention in addition to those shown and described herein will become apparent to those skilled in the art from the foregoing description and fall within the scope of the appended claims.